CLINIMIX E- leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection
CLINIMIX E- leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection
Baxter Healthcare Company

HIGHLIGHTS OF PRESCRIBING INFORMATION
These highlights do not include all the information needed to use CLINIMIX E safely and effectively. See full prescribing information for CLINIMIX E.

CLINIMIX E (amino acids with electrolytes in dextrose with calcium) injection, for intravenous use
Initial U.S. Approval: 1997

RECENT MAJOR CHANGES

Dosage and Administration, Instructions for Use (2.3, 2.7) 04/2021
Warnings and Precautions (5.6) 04/2021
Dosage and Administration (2.1, 2.3, 2.4, 2.6, 2.8) 09/2020

INDICATIONS AND USAGE

CLINIMIX E is indicated as a source of calories, protein, and electrolytes for patients requiring parenteral nutrition when oral or enteral nutrition is not possible, insufficient, or contraindicated. CLINIMIX E may be used to treat negative nitrogen balance in patients. (1)

DOSAGE AND ADMINISTRATION

See full prescribing information for information on preparation, administration, instructions for use, dosing considerations, including the recommended dosage in adults and pediatrics, and dosage modifications in patients with kidney disease. (2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8)

DOSAGE FORMS AND STRENGTHS

CLINIMIX E injection is available in multiple strengths. See full prescribing information for detailed description of each formulation. (3, 11)

CONTRAINDICATIONS

- Concomitant treatment with ceftriaxone in neonates (28 days of age or younger). (4)
- Known hypersensitivity to one or more amino acids or dextrose. (4)
- Inborn errors of amino acid metabolism. (4)
- Patients with pulmonary edema or acidosis due to low cardiac output. (4)

WARNINGS AND PRECAUTIONS

- Pulmonary Embolism due to Pulmonary Vascular Precipitates: if signs of pulmonary distress occur, stop the infusion and initiate a medical evaluation. (5.1)
- Precipitation with Ceftriaxone: do not administer ceftriaxone simultaneously with CLINIMIX E via a Y-site. (4, 5.2, 8.4)
- Hypersensitivity Reactions: monitor for signs and symptoms and discontinue infusion if reactions occur. (5.3)
- Risk of Infections, Refeeding Complications, and Hyperglycemia or Hyperosmolar Hyperglycemic State: monitor for signs and symptoms; monitor laboratory parameters. (5.4, 5.5, 5.6)
- Vein Damage and Thrombosis: solutions with osmolarity of ≥ 900 mOsm/L must be infused through a central catheter. (2.2, 5.7)
- Hepatobiliary Disorders: monitor liver function parameters and ammonia levels. (5.8)
- Aluminum Toxicity: increased risk in patients with impaired kidney function, including preterm infants. (5.9, 8.4)
- Parenteral Nutrition Associated Liver Disease: increased risk in patients who receive parenteral nutrition for extended periods of time, especially preterm infants; monitor liver function tests, if abnormalities occur consider discontinuation or dosage reduction. (5.10, 8.4)
- Electrolyte Imbalance and Fluid Overload: patients with cardiac insufficiency or kidney disease may require adjustment of fluid, protein and electrolyte content. (5.11, 8.4)

ADVERSE REACTIONS

Adverse reactions include diuresis, extravasation, glycosuria, hyperglycemia, and hyperosmolar coma. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Baxter Healthcare Corporation at 1-866-888-2472 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

USE IN SPECIFIC POPULATIONS

Pediatric Use: increased risk of hypoglycemia/hyperglycemia: monitor serum glucose concentrations. (8.4)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 4/2021

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE
2 DOSAGE AND ADMINISTRATION
2.1 Preparation Prior to Administration
2.2 Important Administration Instructions
2.3 Instructions for Use
2.4 Preparation and Addition of Lipid Emulsion
2.5 Dosing Considerations
2.6 Recommended Dosage in Adults
2.7 Dosage Modifications in Patients with Kidney Disease
2.8 Recommended Dosage in Pediatric Patients
2.9 Discontinuation of CLINIMIX E injection
3 DOSAGE FORMS AND STRENGTHS
4 CONTRAINDICATIONS
5 WARNINGS AND PRECAUTIONS
5.1 Pulmonary Embolism due to Pulmonary Vascular Precipitates

5.2 Precipitation with Ceftriaxone
5.3 Hypersensitivity Reactions
5.4 Risk of Infections
5.5 Refeeding Syndrome
5.6 Hyperglycemia or Hyperosmolar Hyperglycemic State
5.7 Vein Damage and Thrombosis
5.8 Hepatobiliary Disorders
5.9 Aluminum Toxicity
5.10 Risk of Parenteral Nutrition Associated Liver Disease
5.11 Electrolyte Imbalance and Fluid Overload
5.12 Monitoring/Laboratory Tests
6 ADVERSE REACTIONS
7 DRUG INTERACTIONS
7.1 Drugs that Can Cause Hyperkalemia
8 USE IN SPECIFIC POPULATIONS
8.1 Pregnancy
8.2 Lactation
8.4 Pediatric Use
8.5 Geriatric Use
10 OVERDOSAGE
11 DESCRIPTION
12 CLINICAL PHARMACOLOGY
12.1 Mechanism of Action
12.3 Pharmacokinetics
16 HOW SUPPLIED/STORAGE AND HANDLING
17 PATIENT COUNSELING INFORMATION
* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

CLINIMIX E is indicated as a source of calories, protein, and electrolytes for patients requiring parenteral nutrition when oral or enteral nutrition is not possible, insufficient, or contraindicated. CLINIMIX E may be used to treat negative nitrogen balance in patients.

2 DOSAGE AND ADMINISTRATION

2.1 Preparation Prior to Administration

- CLINIMIX E is available in a three port container configuration and a two port container configuration.
    - *Three Port Container:* the ports consist of one medication port, one additive port and one outlet port. Additives can be introduced to the container through the medication port and lipids through the additive port on the three port container.
    - *Two Port Container:* the ports consist of one medication port and one outlet port. Additives, including lipids, can be introduced to the container through the medication port on the two port container.
- Tear protective overwrap at slit and remove solution container. Small amounts of moisture may be found on the solution container from water permeating from inside the container. The amount of permeated water is insufficient to affect the solution significantly. If larger amounts of water are found, the container should be checked for tears or leaks.
- Inspect the container prior to activation. Some opacity of the plastic due to moisture absorption during the sterilization process may be observed. This is normal and does not affect the solution quality or safety. The opacity will diminish gradually. Evaluate the following:
    - If the outlet or additive port protectors are damaged, detached, or not present, discard container as solution path sterility may be impaired.
    - Check to ensure seal between chambers is intact, solutions are contained in separate chambers, and the content of the individual chambers is clear, colorless or slightly yellow. Discard if the seal is broken or if the solution is bright yellow or yellowish brown.
    - Check for minute leaks by separately squeezing each chamber. If external leaks or leakage between the chambers are found, discard solution as sterility or stability may be impaired.
- Lipids and/or additives can be introduced to the container after opening seal between chambers. Because additives may be incompatible, evaluate all additions to the plastic container for compatibility. Activate chambers of container prior to introduction of additives. Mix thoroughly when additives have been introduced. Supplemental medication may be added with a 19 to 22 gauge needle through the medication port.
- Calcium and phosphate ratios must be considered. Excess addition of calcium and phosphate, especially in the form of mineral salts, may result in the formation of calcium phosphate precipitates *[see Warnings and Precautions (5.1)].*
- Inspect the container to ensure precipitates have not formed during the mixing or

addition of additives. A slight yellow color does not alter the quality and efficacy of this product. If lipid has been added, ensure the emulsion has not separated. Separation of the emulsion can be visibly identified by a yellowish streaking or the accumulation of yellowish droplets in the mixed emulsion. Discard the admixture if any of the above are observed.

2.2 Important Administration Instructions

- Set the vent to the closed position on a vented intravenous administration set to prevent air embolism.
- Use a dedicated line without any connections to avoid air embolism.
- CLINIMIX E is for intravenous infusion only into a central or peripheral vein. The choice of a central or peripheral venous route should depend on the osmolarity of the final infusate. Solutions with osmolarity of 900 mOsm/L or greater must be infused through a central catheter *[see Warnings and Precautions (5.7)].*

    - For central vein infusion only: CLINIMIX E 4.25/10, 5/15, 5/20, 8/10, 8/14
    - For central or peripheral vein infusion: CLINIMIX E 2.75/5 and 4.25/5

- The solution should be inspected for precipitates before admixing, after admixing, and again before administration.
- Use a 0.22 micron filter for administration of CLINIMIX E. If a lipid is also administered, use a 1.2 micron filter.
- If lipid emulsion is added, do not use administration sets and lines that contain di-2-ethylhexyl phthalate (DEHP). Administration sets that contain polyvinyl chloride (PVC) components have DEHP as a plasticizer.
- Ceftriaxone must not be administered simultaneously with calcium-containing intravenous solutions such as CLINIMIX E via a Y-site. However, in patients other than neonates, ceftriaxone and CLINIMIX E may be administered sequentially if the infusion lines are thoroughly flushed between infusions with a compatible fluid *[see Contraindications (4), Warnings and Precautions (5.2)]*.

2.3 Instructions for Use

1. Open by tearing protective overwrap at slit and remove solution container. The two port container includes an oxygen-absorbing sachet. Discard the oxygen-absorbing sachet after removal from the overwrap.
2. To proceed with activation, the container should be at room temperature. Lay the room temperature container onto a flat surface. Grasp the container firmly on each side of the top of the container (**Figure 1**).
3. Starting from the top, using some pressure, slowly roll the container to open seal between chambers as shown in **Figure 2**. Do not pull or rip the seal apart. The seal must be completely opened towards the port side of the container. The upper section of the seal towards the hanger side can remain unbroken.
4. Mix the contents thoroughly by inverting the container upside down to ensure a homogenous admixture (**Figure 3**).
5. Once the container is mixed, check for leaks.
6. Make additions (if prescribed).
   Because additives may be incompatible, evaluate all additions to the container for compatibility and stability of the resulting preparation. Consult with pharmacist, if available. Questions about compatibility may be directed to Baxter. If it is deemed advisable to introduce additives, use aseptic technique. For information on adding lipid emulsions *see Dosage and Administration (2.4)*.

    a. Prepare medication port.
    b. Using syringe with 19 to 22 gauge needle, puncture resealable medication port and inject.
    c. Mix solution and medication thoroughly (**Figure 3**). For high density medication (high specific gravity), such as potassium chloride, squeeze ports while ports are upright and mix thoroughly.

7. Inspect final solution for discoloration and particulate matter. Check for leaks.
8. Spike and hang container.

    a. Suspend container from eyelet support.
    b. Twist off protector from outlet port at bottom of container (**Figure 4**).
    c. Attach administration set. Refer to complete directions accompanying set.

   For single dose only. Discard unused portion.

Figures 1 – 4 (Three Port Container):



Figures 1 – 4 (Two Port Container):



Instructions on Storage

Storage After Removal of Overwrap:

Once removed from the protective overwrap, mixed (peel seal activated) or unmixed (peel seal intact), CLINIMIX E solutions may be stored under refrigeration for up to 9 days.

Storage Once any Additive is Added:

Use promptly after mixing. Any storage with additives should be under refrigeration and limited to a brief period of time, less than 24 hours. After removal from refrigeration, use promptly and complete the infusion within 24 hours. Any remaining mixture must be discarded.

Protect the activated parenteral nutrition solution from light.

2.4 Preparation and Addition of Lipid Emulsion

Three Port Container

1. Prior to adding lipid emulsion, mix amino acid and dextrose injection as shown in **Figures 1-3**.
2. Prepare lipid emulsion transfer set following instructions provided.
3. Attach transfer set to lipid emulsion container using aseptic technique.

4. Twist off protector on the additive port of the container.
5. Attach the transfer set to the exposed additive port.
6. Open clamp on transfer set.
7. After completing transfer, use appropriate plastic clamp or metal ferrule to seal off additive port tube.
8. Remove transfer set.
9. Mix contents of container thoroughly. Inspect final solution for discoloration and particulate matter. Check for leaks.

Two Port Container

1. Prior to adding lipid emulsion, mix amino acid and dextrose injection as shown in Figures 1-3.
2. Prepare lipid emulsion transfer set following instructions provided.
3. Attach transfer set to lipid emulsion container using aseptic technique.
4. Prepare medication port.
5. Using a 19 to 22 gauge needle, puncture resealable medication port.
6. Open clamp on transfer set and transfer lipid emulsion.
7. Remove needle.
8. Mix contents of container thoroughly. Inspect final solution for discoloration and particulate matter. Check for leaks.

Storage Once Lipids are Added:

Use promptly after mixing. Any storage with additives should be under refrigeration and limited to a brief period of time, no longer than 24 hours. After removal from refrigeration, use promptly and complete the infusion within 24 hours. Any mixture remaining must be discarded.

2.5 Dosing Considerations

- The dosage of CLINIMIX E should be individualized based on the patient's clinical condition (ability to adequately metabolize amino acids and dextrose), body weight and nutritional/fluid requirements, as well as additional energy given orally/enterally to the patient. Prior to initiating CLINIMIX E the following patient information should be reviewed: all concomitant medications, gastrointestinal function and laboratory data such as electrolytes (including magnesium, calcium, and phosphorus), glucose, urea/creatinine, liver panel, complete blood count and triglyceride level (if adding lipid emulsion). Refer to the complete prescribing information of lipid emulsion for dosing information.
- CLINIMIX E formulations have varying concentrations of protein, carbohydrate and a standard concentration of electrolytes; thus infusion rates to achieve requirements will vary. Protein, caloric, fluid and electrolyte requirements all need to be taken into consideration when determining individual patient dosage needs.
- The dosage selection is based only on the recommended protein requirements. The maximum dextrose infusion rates and calorie and fluid requirements must also be considered when determining the clinically appropriate infusion rate for patients.
- CLINIMIX E meets the total nutritional requirements for protein and dextrose in stable patients, and can be individualized to meet specific needs with the addition of nutrients.
- Total daily fluid requirements can be met beyond the volume of amino acids solution by supplementing with non-carbohydrate or carbohydrate-containing electrolyte solutions. In many patients, provision of adequate calories in the form of hypertonic dextrose may require the administration of exogenous insulin to prevent hyperglycemia and glycosuria.
- Prior to administration of CLINIMIX E correct severe fluid, electrolyte and acid-base disorders.
- Monitor levels of serum potassium during therapy. It may be necessary to add additional potassium to the CLINIMIX E admixture.
- Lipid emulsion administration should be considered with prolonged use (more than 5 days) of CLINIMIX E in order to prevent essential fatty acid deficiency (EFAD). Serum lipids should be monitored for evidence of EFAD in patients maintained on fat-free parenteral nutrition. See prescribing information of lipid emulsion.
- The flow rate should be increased gradually. The flow rate must be adjusted taking into account the dose being administered, the daily volume intake, and the duration of the infusion.

2.6 Recommended Dosage in Adults

The recommended daily nutritional requirements for protein and dextrose compared to the amount of nutrition provided by CLINIMIX E are shown in Table 1. As indicated on an individual basis, maintenance vitamins, additional electrolytes, trace elements and other components (including lipids) should be administered as required to prevent deficiencies and complications from developing.
The maximum infusion rates in adult patients are show in Table 2.

In addition to meeting protein needs, the administration rate should be governed, especially during the first few day of therapy, by the patient's tolerance to dextrose. Daily intake of amino acids and dextrose should be increased gradually to the maximum required dose as indicated by frequent determinations of blood glucose levels.

Table 1: Nutritional Comparison – Adult Patients

	Recommended CLINIMIX E Adult Dosage						
	CLINIMIX E 2.75/5	**CLINIMIX E 4.25/5**	**CLINIMIX E 4.25/10**	**CLINIMIX E 5/15**	**CLINIMIX E 5/20**	**CLINIMIX E 8/10**	**CLINIMIX E 8/14**

Fluid (mL/kg/day)	29 to 40	19 to 40	19 to 40	16 to 40	16 to 40	10 to 25	10 to 25
Protein[*](g/kg/day) (Nitrogen g/kg/day)	0.8 to 1.1 (0.13 to 0.18)	0.8 to 1.7 (0.13 to 0.27)	0.8 to 1.7 (0.13 to 0.27)	0.8 to 2 (0.13 to 0.32)	0.8 to 2 (0.13 to 0.32)	0.8 to 2 (0.13 to 0.32)	0.8 to 2 (0.13 to 0.32)
Dextrose (g/kg/day)	1.45 to 2	0.95 to 2	1.9 to 4	2.4 to 6	3.2 to 8	1 to 2.5	1.4 to 3.5

* Protein is provided as amino acids. When infused intravenously amino acids are metabolized and utilized as the building blocks of protein.

Table 2: Maximum Infusion Rate in Adult Patients

		Maximum Infusion Rates in Adults Patients						
		CLINIMIX E 2.75/5	**CLINIMIX E 4.25/5**	**CLINIMIX E 4.25/10**	**CLINIMIX E 5/15**	**CLINIMIX E 5/20**	**CLINIMIX E 8/10**	**CLINIMIX E 8/14**
Maximum Infusion Rate (mL/kg/hour)		3.6	2.4	2.4	1.67	1.25	1.3	1.3
Corresponding infusion rate	**Amino Acid (g/kg/hour)**	0.1[*]	0.1[*]	0.1[*]	0.08	0.06	0.1[*]	0.1[*]
	Dextrose (g/kg/hour)	0.18	0.12	0.24	0.25[*]	0.25[*]	0.13	0.18

* Rate limiting factor

2.7 Dosage Modifications in Patients with Kidney Disease

Prior to administration, correct severe fluid or electrolyte imbalances. Closely monitor serum electrolyte levels and adjust the volume of CLINIMIX E administered as required *[see Warnings and Precautions (5.11)]*.

Chronic kidney disease patients with less than nephrotic range proteinuria require 0.8 g of protein/kg/day. Chronic kidney disease patients with nephrotic range proteinuria require 0.8 g of protein/kg/day plus 1 g of protein for each gram of proteinuria. Patients needing dialysis should receive from 1.2 g of protein/kg/day up to a maximum of 2.5 g of protein/kg/day depending on the nutritional status and the dialysis modality. Serum electrolyte levels should be closely monitored. The CLINIMIX E dosage can be adjusted based on the severity of kidney disease, supplementing protein as indicated. If required, additional amino acids may be added to the CLINIMIX E container or infused separately. Compatibility of additions should be evaluated by a pharmacist and questions may be directed to Baxter.

2.8 Recommended Dosage in Pediatric Patients

The dosage and constant infusion rate of intravenous dextrose must be selected with caution in pediatric patients, particularly neonates and low weight infants, because of the increased risk of hyperglycemia/hypoglycemia *[see Use in Specific Populations (8.4)]*. Frequent monitoring of serum glucose concentrations is required when dextrose is prescribed to pediatric patients, particularly neonates and low birth weight infants. The infusion rate and volume should be determined by the consulting physician experienced in pediatric intravenous fluid therapy.

In pediatric patients, CLINIMIX E is dosed on the basis of protein provided as amino acids. The recommended dosage, by age group is provided in **Tables 3 - 6**. Infusion rates are based on protein and do not take carbohydrates, fluid or electrolytes into consideration.

This product does not contain the amino acids cysteine and taurine, considered conditionally essential for neonates and infants. If possible, these amino acids should be added to this product if used in this pediatric population.

Table 3: Preterm and Term Infants Less than 1 Month of Age

	Recommended CLINIMIX E Dosage in Preterm and Term Infants Less than 1 Month of Age						
	CLINIMIX E 2.75/5	**CLINIMIX E 4.25/5**	**CLINIMIX E 4.25/10**	**CLINIMIX E 5/15**	**CLINIMIX E 5/20**	**CLINIMIX E 8/10**	**CLINIMIX E 8/14**
Infusion Rate Range (mL/kg/hr)	4.5 to 6	2.9 to 3.9	2.9 to 3.9	2.5 to 3.3	2.5 to 3.3	1.6 to 2.1	1.6 to 2.1
Fluid (mL/kg/day)	108 to 144	70 to 94	70 to 94	60 to 79	60 to 79	38.4 to 50	38.4 to 50
Protein[*](g/kg/day) (Nitrogen g/kg/day)	3 to 4 (0.48 to 0.64)	3 to 4 (0.48 to 0.64)	3 to 4 (0.48 to 0.64)	3 to 4 (0.48 to 0.64)	3 to 4 (0.48 to 0.64)	3 to 4 (0.48 to 0.64)	3 to 4 (0.48 to 0.64)
Dextrose (g/kg/day)	5.4 to 7.2	3.5 to 4.7	7 to 9.4	9 to 11.9	12 to 15.8	3.8 to 5	5.4 to 7

* Protein is provided as amino acids. When infused intravenously amino acids are metabolized and utilized as the building blocks of protein.

Table 4: Pediatric Patients 1 Month to Less than 1 Year of Age

	Recommended CLINIMIX E Dosage in Pediatric Patients 1 Month to Less than 1 Year of Age						
	CLINIMIX E 2.75/5	**CLINIMIX E 4.25/5**	**CLINIMIX E 4.25/10**	**CLINIMIX E 5/15**	**CLINIMIX E 5/20**	**CLINIMIX E 8/10**	**CLINIMIX E 8/14**
Infusion Rate	3 to 4.5	2 to 2.9	2 to 2.9	1.7 to 2.5	1.7 to 2.5	1 to 1.6	1 to 1.6

Range (mL/kg/hr)	3 to 4.5	2 to 2.9	2 to 2.9	1.7 to 2.5	1.7 to 2.5	1 to 1.6	1 to 1.6
Fluid (mL/kg/day)	72 to 108	48 to 70	48 to 70	41 to 60	41 to 60	24 to 38.4	24 to 38.4
Protein* (g/kg/day) (Nitrogen g/kg/day)	2 to 3 (0.32 to 0.48)	2 to 3 (0.32 to 0.48)	2 to 3 (0.32 to 0.48)	2 to 3 (0.32 to 0.48)	2 to 3 (0.32 to 0.48)	2 to 3 (0.32 to 0.48)	2 to 3 (0.32 to 0.48)
Dextrose (g/kg/day)	3.6 to 5.4	2.4 to 3.5	4.8 to 7	6.1 to 9	8.2 to 12	2.4 to 3.8	3.4 to 5.4

* Protein is provided as amino acids. When infused intravenously amino acids are metabolized and utilized as the building blocks of protein.

Table 5: Pediatric Patients 1 Year to Less than 11 Years of Age

	Recommended CLINIMIX E Dosage in Pediatric Patients 1 Year to Less than 11 Years of Age						
	CLINIMIX E 2.75/5	**CLINIMIX E 4.25/5**	**CLINIMIX E 4.25/10**	**CLINIMIX E 5/15**	**CLINIMIX E 5/20**	**CLINIMIX E 8/10**	**CLINIMIX E 8/14**
Infusion Rate Range (mL/kg/hr)	1.5 to 3	1 to 2	1 to 2	0.8 to 1.7	0.8 to 1.7	0.5 to 1	0.5 to 1
Fluid (mL/kg/day)	36 to 72	24 to 48	24 to 48	19 to 41	19 to 41	12 to 24	12 to 24
Protein* (g/kg/day) (Nitrogen g/kg/day)	1 to 2 (0.16 to 0.32)	1 to 2 (0.16 to 0.32)	1 to 2 (0.16 to 0.32)	1 to 2 (0.16 to 0.32)	1 to 2 (0.16 to 0.32)	1 to 2 (0.16 to 0.32)	1 to 2 (0.16 to 0.32)
Dextrose (g/kg/day)	1.8 to 3.6	1.2 to 2.4	2.4 to 4.8	2.9 to 6.1	3.8 to 8.2	1.2 to 2.4	1.7 to 3.4

* Protein is provided as amino acids. When infused intravenously amino acids are metabolized and utilized as the building blocks of protein.

Table 6: Pediatric Patients 11 Years to 17 Years of Age

	Recommended CLINIMIX E Dosage in Pediatric Patients 11 Years to 17 Years of Age						
	CLINIMIX E 2.75/5	**CLINIMIX E 4.25/5**	**CLINIMIX E 4.25/10**	**CLINIMIX E 5/15**	**CLINIMIX E 5/20**	**CLINIMIX E 8/10**	**CLINIMIX E 8/14**
Infusion Rate Range (mL/kg/hr)	1.2 to 2.3	0.8 to 1.5	0.8 to 1.5	0.7 to 1.3	0.7 to 1.3	0.4 to 0.8	0.4 to 0.8
Fluid (mL/kg/day)	29 to 55	19 to 36	19 to 36	17 to 31	17 to 31	9.6 to 19.2	9.6 to 19.2
Protein* (g/kg/day) (Nitrogen g/kg/day)	0.8 to 1.5 (0.13 to 0.24)	0.8 to 1.5 (0.13 to 0.24)	0.8 to 1.5 (0.13 to 0.24)	0.8 to 1.5 (0.13 to 0.24)	0.8 to 1.5 (0.13 to 0.24)	0.8 to 1.5 (0.13 to 0.24)	0.8 to 1.5 (0.13 to 0.24)
Dextrose (g/kg/day)	1.4 to 2.8	1 to 1.8	1.9 to 3.6	2.5 to 4.7	3.4 to 6.2	1 to 1.9	1.4 to 2.7

* Protein is provided as amino acids. When infused intravenously amino acids are metabolized and utilized as the building blocks of protein.

2.9 Discontinuation of CLINIMIX E injection

To reduce the risk of hypoglycemia after discontinuation, a gradual decrease in flow rate in the last hour of infusion should be considered.

3 DOSAGE FORMS AND STRENGTHS

CLINIMIX E injection is available in 1000 mL and 2000 mL dual chamber containers. The individual chambers contain essential and nonessential amino acids with electrolytes and dextrose with calcium. Table 7 describes the individual components of CLINIMIX E.

Table 7: INGREDIENTS PER 100mL OF CLINIMIX E

Strength of CLINIMIX E		CLINIMIX E 2.75/5 sulfite-free (2.75% Amino Acid in 5% Dextrose) Injection	CLINIMIX E 4.25/5 sulfite-free (4.25% Amino Acid in 5% Dextrose) Injection	CLINIMIX E 4.25/10 sulfite-free (4.25% Amino Acid in 10% Dextrose) Injection	CLINIMIX E 5/15 sulfite-free (5% Amino Acid in 15% Dextrose) Injection	CLINIMIX 5/20 sulfite-free (5% Amino Acid in 20% Dextrose) Injection	CLINIMIX E 8/10 sulfite-free (8% Amino Acid in 10% Dextrose) Injection	CLINIMIX E 8/14 sulfite-free (8% Amino Acid in 14% Dextrose) Injection
	Dextrose Hydrous, USP (g/100 mL)	5	5	10	15	20	10	14
	Amino Acids (g/100 mL)	2.75	4.25	4.25	5	5	8	8
	Total Nitrogen (mg/100 mL)	454	702	702	826	826	1320	1320
Essential Amino Acids (mg/100 mL)	Leucine	201	311	311	365	365	584	584
	Isoleucine	165	255	255	300	300	480	480
	Valine	160	247	247	290	290	464	464
	Lysine (added as the hydrochloride salt)	159	247	247	290	290	464	464
	Phenylalanine	154	238	238	280	280	448	448

	Histidine	132	204	204	240	240	384	384
	Threonine	116	179	179	210	210	336	336
	Methionine	110	170	170	200	200	320	320
	Tryptophan	50	77	77	90	90	144	144
Nonessential Amino Acids (mg/100 mL)	Alanine	570	880	880	1035	1035	1656	1656
	Arginine	316	489	489	575	575	920	920
	Glycine	283	438	438	515	515	824	824
	Proline	187	289	289	340	340	544	544
	Serine	138	213	213	250	250	400	400
	Tyrosine	11	17	17	20	20	32	32
Electrolytes (mg/100 mL)	Sodium Acetate Trihydrate, USP	217	297	297	340	340	0	0
	Dibasic Potassium Phosphate, USP	261	261	261	261	261	261	261
	Sodium Chloride, USP	112	77	77	59	59	205	205
	Magnesium Chloride, USP	51	51	51	51	51	51	51
	Calcium Chloride Dihydrate, USP	33	33	33	33	33	33	33
Electrolyte Profile (mEq/L)*	Sodium	35	35	35	35	35	35	35
	Potassium	30	30	30	30	30	30	30
	Magnesium	5	5	5	5	5	5	5
	Calcium	4.5 (2.2 mmol/L)	4.5 (2.2 mmol/L)	4.5 (2.2 mmol/L)	4.5 (2.2 mmol/L)	4.5 (2.2 mmol/L)	4.5 (2.2 mmol/L)	4.5 (2.2 mmol/L)
	Acetate†	51	70	70	80	80	83	83
	Chloride‡	39	39	39	39	39	76	76
	Phosphate (as $HPO_4^{=}$)	30 (15 mmol/L)	30 (15 mmol/L)	30 (15 mmol/L)	30 (15 mmol/L)	30 (15 mmol/L)	30 (15 mmol/L)	30 (15 mmol/L)
	pH§ (Range)	6.0 (4.5 to 7.0)	6.0 (4.5 to 7.0)	6.0 (4.5 to 7.0)	6.0 (4.5 to 7.0)	6.0 (4.5 to 7.0)	6.0 (4.5 to 7.0)	6.0 (4.5 to 7.0)
	Osmolarity (mOsmol/L) (calc)	665	815	1070	1395	1650	1450	1650
Caloric Content (kcal/L)	From Dextrose	170	170	340	510	680	343	477
	From Amino Acids	110	170	170	200	200	320	320
	TOTAL (Dextrose and Amino Acids)	280	340	510	710	880	663	797

* Balanced by ions from amino acids.
† Derived from glacial acetic acid (for pH adjustment) and sodium acetate.
‡ Contributed by calcium chloride, lysine hydrochloride, magnesium chloride, sodium chloride, and hydrochloric acid.
§ pH of sulfite-free amino acid injection with electrolytes in the outlet port chamber was adjusted with glacial acetic acid and pH of dextrose injection port chamber was adjusted with hydrochloric acid.

4 CONTRAINDICATIONS

The use of CLINIMIX E is contraindicated in:

1. Neonates (28 days of age or younger) receiving concomitant treatment with ceftriaxone, even if separate infusion lines are used, due to the risk of fatal ceftriaxone calcium salt precipitation in the neonate's bloodstream *[see Warnings and Precautions (5.2), Use in Specific Populations (8.4)]*.
2. Patients with known hypersensitivity to one or more amino acids or dextrose *[see Warnings and Precautions (5.3)]*.
3. Patients with inborn errors of amino acid metabolism due to risk of severe metabolic and neurologic complications.
4. Patients with pulmonary edema or acidosis due to low cardiac output.

5 WARNINGS AND PRECAUTIONS

5.1 Pulmonary Embolism due to Pulmonary Vascular Precipitates

Pulmonary vascular precipitates causing pulmonary vascular emboli and pulmonary distress have been reported in patients receiving parenteral nutrition. In some cases, fatal outcomes due to pulmonary embolism have occurred. Patients, especially those with hypophosphatemia, may require the addition of phosphate. To prevent hypocalcemia, calcium supplementation should always accompany phosphate administration. Excessive addition of calcium and phosphate increases the risk of the formation of calcium phosphate precipitates. Precipitates have been reported even in the absence of phosphate salt in the solution. Precipitation following passage through an in-line filter and suspected in vivo precipitate formation has also been reported. If signs of pulmonary distress occur, stop the infusion and initiate a medical evaluation. In addition to inspection of the solution *[see Dosage and Administration (2.1, 2.2, 2.3, 2.4)]*, the

infusion set and catheter should also periodically be checked for precipitates.

5.2 Precipitation with Ceftriaxone

Precipitation of ceftriaxone-calcium can occur when ceftriaxone is mixed with calcium-containing parenteral nutrition solutions, such as CLINIMIX E, in the same intravenous administration line. Do not administer ceftriaxone simultaneously with CLINIMIX E via a Y-site.

Deaths have occurred in neonates (less than 28 days of age) who received concomitant intravenous calcium-containing solutions with ceftriaxone resulting from calcium-ceftriaxone precipitates in the lungs and kidneys, even when separate infusion lines were used. CLINIMIX E is contraindicated in neonates receiving ceftriaxone *[see Contraindications (4),Use in Specific Populations (8.4)].*

In patients older than 28 days (including adults), ceftriaxone and CLINIMIX E may be administered sequentially if the infusion lines are thoroughly flushed between infusions with a compatible fluid.

5.3 Hypersensitivity Reactions

Hypersensitivity/infusion reactions including anaphylaxis have been reported with CLINIMIX E. Stop infusion immediately and treat patient accordingly if any signs or symptoms of a hypersensitivity reaction develop. Signs or symptoms may include: hypotension, hypertension, peripheral cyanosis, tachycardia, dyspnea, vomiting, nausea, urticaria, rash, pruritus, erythema, hyperhidrosis, pyrexia, and chills.

5.4 Risk of Infections

Patients who require parenteral nutrition are at high risk of infections because the nutritional components of these solutions can support microbial growth. Infection and sepsis may also occur as a result of the use of intravenous catheters to administer parenteral nutrition.

The risk of infection is increased in patients with malnutrition-associated immunosuppression, hyperglycemia exacerbated by dextrose infusion, long-term use and poor maintenance of intravenous catheters, or immunosuppressive effects of other concomitant conditions, drugs, or other components of the parenteral formulation (e.g., lipid emulsion).

To decrease the risk of infection, ensure aseptic technique in catheter placement and maintenance, as well as aseptic technique in the preparation and administration of the nutritional formula.

Monitor for signs and symptoms (including fever and chills) of early infections, including laboratory test results (including leukocytosis and hyperglycemia) and frequent checks of the parenteral access device and insertion site for edema, redness and discharge.

5.5 Refeeding Syndrome

Refeeding severely undernourished patients may result in refeeding syndrome, characterized by the intracellular shift of potassium, phosphorus, and magnesium as the patient becomes anabolic. Thiamine deficiency and fluid retention may also develop. To prevent these complications, monitor severely undernourished patients and slowly increase nutrient intakes.

5.6 Hyperglycemia or Hyperosmolar Hyperglycemic State

When using CLINIMIX E in patients with diabetes mellitus, impaired glucose tolerance may worsen hyperglycemia. Administration of dextrose at a rate exceeding the patient’s utilization rate may lead to hyperglycemia, coma, and death. Patients with dehydration, resulting in a transient reduction in glomerular filtration rate and pre-renal azotemia, may be at greater risk of developing hyperosmolar hyperglycemic state. Monitor blood glucose levels and treat hyperglycemia to maintain optimum levels while administering CLINIMIX E. Insulin may be administered or adjusted to maintain optimal blood glucose levels during CLINIMIX E administration.

5.7 Vein Damage and Thrombosis

Solutions with osmolarity of 900 mOsm/L or greater must be infused through a central catheter. CLINIMIX E solutions containing more than 5% dextrose have an osmolarity greater than or equal to 900 mOsm/L. CLINIMIX E 4.25/10, 5/15, 5/20, 8/10 and 8/14 are indicated for administration into a central vein only, such as the superior vena cava *[see Dosage and Administration (2.2)]*. The infusion of hypertonic nutrient injections into a peripheral vein may result in vein irritation, vein damage, and/or thrombosis.

CLINIMIX E 2.75/5 and 4.25/5 are indicated for peripheral administration, or may be infused into a central vein *[see Dosage and Administration (2.2)]*. The primary complication of peripheral access is venous thrombophlebitis, which manifests as pain, erythema, tenderness or a palpable cord. Remove the catheter as soon as possible, if thrombophlebitis develops.

5.8 Hepatobiliary Disorders

Hepatobiliary disorders are known to develop in some patients without preexisting liver disease who receive parenteral nutrition, including cholecystitis, cholelithiasis, cholestasis, hepatic steatosis, fibrosis and cirrhosis, possibly leading to hepatic failure. The etiology of these disorders is thought to be multifactorial and may differ between patients.

Increase in blood ammonia levels and hyperammonemia may occur in patients receiving amino acid solutions. In some patients this may indicate hepatic insufficiency or the

presence of an inborn error of amino acid metabolism *[see Contraindications (4)]*.

Monitor liver function parameters and ammonia levels. Patients developing signs of hepatobiliary disorders should be assessed early by a clinician knowledgeable in liver diseases in order to identify possible causative and contributory factors, and possible therapeutic and prophylactic interventions.

5.9 Aluminum Toxicity

CLINIMIX E contains no more than 25 mcg/L of aluminum. The aluminum contained in CLINIMIX E may reach toxic levels with prolonged administration in patients with impaired kidney function.

Preterm infants are at greater risk because their kidneys are immature, and they require large amounts of calcium and phosphate solutions, which contain aluminum. Patients with impaired kidney function, including preterm infants, who receive parenteral levels of aluminum at greater than 4 to 5 mcg/kg/day, accumulate aluminum at levels associated with central nervous system and bone toxicity. Tissue loading may occur at even lower rates of administration.

5.10 Risk of Parenteral Nutrition Associated Liver Disease

Parenteral Nutrition Associated Liver Disease (PNALD) has been reported in patients who receive parenteral nutrition for extended periods of time, especially preterm infants, and can present as cholestasis or steatohepatitis. The exact etiology is unknown and is likely multifactorial. If CLINIMIX E treated patients develop liver test abnormalities consider discontinuation or dosage reduction.

5.11 Electrolyte Imbalance and Fluid Overload

Patients with abnormal renal function due to pre-renal azotemia, renal obstruction, or intrinsic kidney disease may be at increased risk of electrolyte and fluid volume imbalance. Patients with cardiac insufficiency due to left ventricular systolic dysfunction are susceptible to excess fluid accumulation. Use CLINIMIX E with caution in patients with cardiac insufficiency or kidney disease. CLINIMIX E dosage may require adjustment with specific attention to fluid, protein, and electrolyte content in these patients.

Monitor renal function parameters. Patients developing signs of kidney disease should be assessed early by a clinician knowledgeable in kidney disease in order to determine the appropriate CLINIMIX E dosage and other treatment options.

5.12 Monitoring/Laboratory Tests

Monitor fluid and electrolyte status, serum osmolarity, blood glucose, liver and kidney function, blood count and coagulation parameters throughout treatment. In situations of severely elevated electrolyte levels, stop CLINIMIX E until levels have been corrected.

6 ADVERSE REACTIONS

The following serious adverse reactions are discussed in greater detail in other sections of the prescribing information.

- Pulmonary embolism due to pulmonary vascular precipitates *[see Warnings and Precautions (5.1)]*
- Death in neonates due to calcium-ceftriaxone precipitates *[see Warnings and Precautions (5.2)]*
- Hypersensitivity reactions *[see Warnings and Precautions (5.3)]*
- Risk of Infections *[see Warnings and Precautions (5.4)]*
- Refeeding syndrome *[see Warnings and Precautions (5.5)]*
- Hyperglycemia or hyperosmolar hyperglycemic state *[see Warnings and Precautions (5.6)]*
- Vein damage and thrombosis *[see Warnings and Precautions (5.7)]*
- Hepatobiliary disorders *[see Warnings and Precautions (5.8)]*
- Parenteral Nutrition Associated Liver Disease (PNALD) *[see Warnings and Precautions (5.10)]*
- Electrolyte imbalance and fluid overload *[see Warnings and Precautions (5.11)]*

The following adverse reactions from voluntary reports or clinical studies have been reported with CLINIMIX E. Because many of these reactions were reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

- Diuresis
- Extravasation
- Glycosuria
- Hyperglycemia
- Hyperosmolar coma

7 DRUG INTERACTIONS

7.1 Drugs that Can Cause Hyperkalemia

Because of its potassium content, CLINIMIX E should be administered with caution in patients treated with agents or products that can cause hyperkalemia or increase the risk of hyperkalemia, such as potassium sparing diuretics (amiloride, spironolactone, triamterene), with ACE inhibitors, angiotensin II receptor antagonists, or the immunosuppressants tacrolimus and cyclosporine.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are no adequate or well-controlled studies in pregnant women with CLINIMIX E. Additionally, animal reproduction studies have not been conducted with amino acids and electrolytes and dextrose. It is not known whether CLINIMIX E can cause fetal harm when administered to a pregnant woman.

The estimated background risk of major birth defects and miscarriage for the indicated population is unknown. However, the estimated background risk in the U.S. general population of major birth defects is 2 to 4% and of miscarriage is 15 to 20% of clinically recognized pregnancies.

Clinical Considerations

Disease-Associated Maternal and/or Embryo-Fetal Risk

Based on clinical practice guidelines, parenteral nutrition should be considered in cases of severe maternal malnutrition where nutritional requirements cannot be fulfilled by the enteral route because of the risks to the fetus associated with severe malnutrition, such as preterm delivery, low birth weight, intrauterine growth restriction, congenital malformations and perinatal mortality.

8.2 Lactation

Risk Summary

It is not known whether CLINIMIX E is present in human milk. There are no data on the effects of CLINIMIX E on the breastfed infant or on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for CLINIMIX E and any potential adverse effects on the breastfed child from CLINIMIX E or from the underlying maternal condition.

8.4 Pediatric Use

Safety and effectiveness of CLINIMIX E in pediatric patients have not been established by adequate and well-controlled studies. Use of dextrose, amino acid infusions and electrolytes in pediatric patients is based on clinical practice *[see Dosage and Administration (2.8)]*.

Deaths have occurred in neonates (28 days of age or younger) who received concomitant intravenous calcium-containing solutions with ceftriaxone resulting from calcium-ceftriaxone precipitates in the lungs and kidneys, even when separate infusion lines were used. CLINIMIX E is contraindicated in neonates receiving ceftriaxone *[see Contraindications (4), Warnings and Precautions (5.2)].*

Newborns, especially those born premature and with low birth weight, are at increased risk of developing hypo – or hyperglycemia and therefore need close monitoring during treatment with intravenous glucose solutions to ensure adequate glycemic control in order to avoid potential long term adverse effects. Hypoglycemia in the newborn can cause prolonged seizures, coma and brain damage. Hyperglycemia has been associated with intraventricular hemorrhage, late onset bacterial and fungal infection, retinopathy of prematurity, necrotizing enterocolitis, bronchopulmonary dysplasia, prolonged length of hospital stay, and death. Plasma electrolyte concentrations should be closely monitored in the pediatric population as this population may have impaired ability to regulate fluids and electrolytes.

Because of immature renal function, preterm infants receiving prolonged treatment with CLINIMIX E, may be at risk of aluminum toxicity *[see Warnings and Precautions (5.9)]*.

Patients, including pediatric patients, may be at risk for Parenteral Nutrition Associated Liver Disease (PNALD) *[see Warnings and Precautions (5.10)]*.

Hyperammonemia is of special significance in infants (birth to two years). This reaction appears to be related to a deficiency of the urea cycle amino acids of genetic or product origin. It is essential that blood ammonia be measured frequently in infants *[see Warnings and Precautions (5.8)]*.

8.5 Geriatric Use

Clinical studies of CLINIMIX E did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from other younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients.

In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or drug therapy.

10 OVERDOSAGE

An increased infusion rate of CLINIMIX E can cause hyperglycemia, hyperosmolality, and adverse effects on water and electrolyte balance *[see Warnings and Precautions (5.6, 5.11)].*

Severe hyperglycemia and severe dilutional hyponatremia, and their complications, can be fatal.

Discontinue infusion and institute appropriate corrective measures in the event of

overhydration or solute overload during therapy, with particular attention to respiratory and cardiovascular systems.

For current information on the management of poisoning or overdosage, contact the National Poison Control Center at 1-800-222-1222 or www.poison.org.

11 DESCRIPTION

CLINIMIX E sulfite-free (amino acids with electrolytes in dextrose with calcium) injection for intravenous use consists of sterile, nonpyrogenic, hypertonic solutions in a dual chamber container.

The outlet port chamber contains essential and nonessential amino acids with electrolytes. The formulas for the individual electrolytes and amino acids are provided in Table 8**.**

Table 8: Formulas for Electrolytes and Amino Acids

Electrolytes	
Sodium Acetate	$C_2H_3NaO_2 \cdot 3H_2O$
Potassium Phosphate, dibasic	K_2HPO_4
Magnesium Chloride	$MgCl_2 \cdot 6H_2O$
Sodium Chloride	NaCl
Essential Amino Acids	
Leucine	$(CH_3)_2$ $CHCH_2CH$ (NH_2) COOH
Isoleucine	CH_3CH_2CH (CH_3) CH (NH_2) COOH
Valine	$(CH_3)_2$ CHCH (NH_2) COOH
Lysine (added as the hydrochloride salt)	H_2N $(CH_2)_4$ CH (NH_2) COOH
Phenylalanine	(C_6H_5) CH_2 CH (NH_2) COOH
Histidine	$(C_3H_3N_2)$ CH_2CH (NH_2) COOH
Threonine	CH_3CH (OH) CH (NH_2) COO
Methionine	CH_3S $(CH_2)2$ CH (NH_2) COOH
Tryptophan	(C_8H_6N) CH_2 CH (NH_2) COOH
Nonessential Amino Acids	
Alanine	CH_3CH (NH_2) COOH
Arginine	H_2NC (NH) NH $(CH_2)3$ CH (NH_2) COOH
Glycine	H_2NCH_2COOH
Proline	$[(CH_2)_3$ NH CH] COOH
Serine	$HOCH_2CH$ (NH_2) COOH
Tyrosine	$[C_6H_4$ (OH)] CH_2CH (NH_2) COOH

The injection port chamber contains dextrose with calcium. The formula for Calcium Chloride is: $CaCl_2 \cdot 2H_2O$. Dextrose, USP, is chemically designated D-glucose, monohydrate ($C_6H_{12}O_6 \cdot H_2O$) and has the following structure:



Dextrose Hydrous, USP
(D-Glucose monohydrate)

Dextrose is derived from corn.

See Table 7 for composition, pH, osmolarity, ionic concentration and caloric content of the admixed product *[see Dosage Forms and Strengths (3)].*

The dual chamber container is a lipid-compatible plastic container (PL 2401 Plastic).

CLINIMIX E contains no more than 25 mcg/L of aluminum.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

CLINIMIX E is used as a supplement of nutrition in patients, providing macronutrients (amino acids and dextrose) and micronutrients (electrolytes) parenterally.

The amino acids provide the structural units that make up proteins and are used to synthesize proteins and other biomolecules or are oxidized to urea and carbon dioxide as a source of energy.

The administered dextrose is oxidized to carbon dioxide and water, yielding energy.

12.3 Pharmacokinetics

The disposition of infused amino acids, dextrose, and electrolytes are essentially the same as those absorbed from ordinary food.

16 HOW SUPPLIED/STORAGE AND HANDLING

CLINIMIX E (amino acids with electrolytes in dextrose with calcium) injection (sulfite-free) is available in 1000 mL and 2000 mL volumes (See **Table 9**).

Table 9: CLINIMIX E Formulations (per 07-19-00-4607 and BE-30-03-649)

After mixing, the product represents	1000 mL Code and NDC Number	2000 mL Code and NDC Number
CLINIMIX E 2.75/5 sulfite-free (2.75% Amino Acid with Electrolytes in 5% Dextrose with Calcium) Injection	Code 2B7735 NDC 0338-1142-03	
CLINIMIX E 4.25/5 sulfite-free (4.25% Amino Acid with Electrolytes in 5% Dextrose with Calcium) Injection	Code 2B7737 NDC 0338-1144-03	Code 2B7716 NDC 0338-1113-04
CLINIMIX E 4.25/10 sulfite-free (4.25% Amino Acid with Electrolytes in 10% Dextrose with Calcium) Injection	Code 2B7738 NDC 0338-1145-03	Code 2B7717 NDC 0338-1115-04
CLINIMIX E 5/15 sulfite-free (5% Amino Acid with Electrolytes in 15% Dextrose with Calcium) Injection	Code 2B7740 NDC 0338-1147-03	Code 2B7721 NDC 0338-1123-04
CLINIMIX E 5/20 sulfite-free (5% Amino Acid with Electrolytes in 20% Dextrose with Calcium) Injection	Code 2B7741 NDC 0338-1148-03	Code 2B7722 NDC 0338-1125-04
CLINIMIX E 8/10 sulfite-free (8% Amino Acid with Electrolytes in 10% Dextrose with Calcium) Injection	Code EADB9943 NDC 0338-210-06	Code EADB9945 NDC 0338-0214-04
CLINIMIX E 8/14 sulfite-free (8% Amino Acid with Electrolytes in 14% Dextrose with Calcium) Injection	Code EADB9963 NDC 0338-0202-06	Code EADB9965 NDC 0338-0206-04

Table 9: CLINIMIX E Formulations (per BE-30-04-048)

After mixing, the product represents	1000 mL Code and NDC Number	2000 mL Code and NDC Number
CLINIMIX E 2.75/5 sulfite-free (2.75% Amino Acid with Electrolytes in 5% Dextrose with Calcium) Injection	Code 2B7735L NDC 0338-7020-01	/
CLINIMIX E 4.25/5 sulfite-free (4.25% Amino Acid with Electrolytes in 5% Dextrose with Calcium) Injection	Code 2B7737L NDC 0338-7022-01	Code 2B7716L NDC 0338-7024-01
CLINIMIX E 4.25/10 sulfite-free (4.25% Amino Acid with Electrolytes in 10% Dextrose with Calcium)	Code 2B7738L NDC 0338-7026-01	Code 2B7717L NDC 0338-7028-01

Injection		
CLINIMIX E 5/15 sulfite-free (5% Amino Acid with Electrolytes in 15% Dextrose with Calcium) Injection	Code 2B7740L NDC 0338-7030-01	Code 2B7721L NDC 0338-7032-01
CLINIMIX E 5/20 sulfite-free (5% Amino Acid with Electrolytes in 20% Dextrose with Calcium) Injection	Code 2B7741L NDC 0338-7034-01	Code 2B7722L NDC 0338-7036-01
CLINIMIX E 8/10 sulfite-free (8% Amino Acid with Electrolytes in 10% Dextrose with Calcium) Injection	Code EADB9943 NDC 0338-0210-06	Code EADB9945 NDC 0338-0214-04
CLINIMIX E 8/14 sulfite-free (8% Amino Acid with Electrolytes in 14% Dextrose with Calcium) Injection	Code EADB9963 NDC 0338-0202-06	Code EADB9965 NDC 0338-0206-04

Minimize exposure of CLINIMIX E to heat and avoid excessive heat.

Protect from freezing.

Store CLINIMIX E at room temperature (25°C/77°F) (may briefly store at up to 40°C/104°F).

Refrigerated storage is limited to 9 days once the protective overwrap has been opened.

Do not use if the protective overwrap has been previously opened or damaged.

For storage of admixed solutions see *Dosage and Administration (2.3, 2.4)*.

17 PATIENT COUNSELING INFORMATION

Inform patients, caregivers, or home healthcare providers of the following risks of CLINIMIX E:

- Pulmonary embolism due to pulmonary vascular precipitates *[see Warnings and Precautions (5.1)]*
- Death in neonates due to calcium-ceftriaxone precipitates *[see Warnings and Precautions (5.2)]*
- Hypersensitivity reactions *[see Warnings and Precautions (5.3)]*
- Risk of Infections *[see Warnings and Precautions (5.4)]*
- Refeeding syndrome *[see Warnings and Precautions (5.5)]*
- Hyperglycemia or hyperosmolar hyperglycemic state *[see Warnings and Precautions (5.6)]*
- Vein damage and thrombosis *[see Warnings and Precautions (5.7)]*
- Hepatobiliary disorders *[see Warnings and Precautions (5.8)]*
- Aluminum toxicity *[see Warnings and Precautions (5.9)]*
- Parenteral Nutrition Associated Liver Disease (PNALD) *[see Warnings and Precautions (5.10)]*
- Electrolyte imbalance and fluid overload *[see Warnings and Precautions (5.11)]*

Baxter Healthcare Corporation
Deerfield, IL 60015 USA
Baxter and Clinimix E are registered trademarks of Baxter International Inc.
07-19-00-4607 [Applies to Baxter Healthcare Corporation, Jayuya – Puerto Rico manufacturing plant]
BE-30-03-649 [Applies to Baxter SA, Lessines – Belgium manufacturing plant for Clinimix E 8/20 and Clinimix E 8/14]
BE-30-04-048 [Applies to Baxter SA, Lessines – Belgium manufacturing plant for the other Clinimix E formulations]

PACKAGE LABEL - PRINCIPAL DISPLAY PANEL



Container Label

LOT EXP

2B7735 NDC 0338-1142-03

CLINIMIX E
WITH ELECTROLYTES 2.75/5

SULFITE-FREE
(2.75% Amino Acids
with Electrolytes in
5% Dextrose with
Calcium) Injection

500 mL INJECTION PORT CHAMBER
10% Dextrose Injection with Calcium

500 mL OUTLET PORT CHAMBER
5.5% Amino Acid Injection

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESRCIBING INFORMATION FOR INSTRUCTIONS
ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY
AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED
INJECTION

DEXTROSE HYDROUS USP 5 g
ESSENTIAL AMINO ACIDS
LEUCINE 201 mg
ISOLEUCINE 165 mg
VALINE 160 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT) 159 mg
PHENYLALANINE 154 mg
HISTIDINE 132 mg
THREONINE 116 mg
METHIONINE 110 mg
TRYPTOPHAN 50 mg
NONESSENTIAL AMINO ACIDS
ALANINE 570 mg
ARGININE 316 mg
GLYCINE 283 mg
PROLINE 187 mg
SERINE 138 mg
TYROSINE 11 mg
ELECTROLYTES
SODIUM ACETATE TRIHYDRATE USP 217 mg
DIBASIC POTASSIUM PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 112 mg

MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE DIHYDRATE USP 33 mg
mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 51
CHLORIDE 39
PHOSPHATE 30 (15 mmol/L)
BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
STERILE
SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)
AVOID EXCESSIVE HEAT
PROTECT FROM FREEZING
SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 US
MADE IN USA



Container Label

LOT EXP

2B7737 NDC 0338-1144-03

CLINIMIX E
WITH ELECTROLYTES 4.25/5

SULFITE-FREE
(4.25% Amino Acid
with Electrolytes in
5% Dextrose with
Calcium) Injection

500 mL INJECTION PORT CHAMBER
10% Dextrose Injection with Calcium

500 mL OUTLET PORT CHAMBER
8.5% Amino Acid Injection

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESRCIBING INFORMATION FOR INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED
INJECTION

DEXTROSE HYDROUS USP 5 g
ESSENTIAL AMINO ACIDS
LEUCINE 311 mg
ISOLEUCINE 255 mg
VALINE 247 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT) 247 mg
PHENYLALANINE 238 mg
HISTIDINE 204 mg
THREONINE 179 mg
METHIONINE 170 mg
TRYPTOPHAN 77 mg
NONESSENTIAL AMINO ACIDS
ALANINE 880 mg
ARGININE 489 mg
GLYCINE 438 mg
PROLINE 289 mg
SERINE 213 mg
TYROSINE 17 mg
ELECTROLYTES
SODIUM ACETATE TRIHYDRATE USP 297 mg
DIBASIC POTASSIUM PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 77 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE DIHYDRATE USP 33 mg
mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 70
CHLORIDE 39
PHOSPHATE 30 (15 mmol/L)
BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
STERILE
SINGLE DOSE CONTAINER
ROOM TEMPERATURE (25°C/77°F)
AVOID EXCESSIVE HEAT
PROTECT FROM FREEZING
SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 US
MADE IN USA



Container Label

LOT EXP

2B7716 NDC 0338-1113-04

CLINIMIX E
WITH ELECTROLYTES 4.25/5

SULFITE-FREE
(4.25% Amino Acid
with Electrolytes in
5% Dextrose with
Calcium) Injection

1000 mL INJECTION PORT CHAMBER
10% Dextrose Injection with Calcium

1000 mL OUTLET PORT CHAMBER
8.5% Amino Acid Injection

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESRCIBING INFORMATION FOR INSTRUCTIONS
ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE
QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED
INJECTION

DEXTROSE HYDROUS USP 5 g
ESSENTIAL AMINO ACIDS
LEUCINE 311 mg
ISOLEUCINE 255 mg
VALINE 247 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT) 247 mg
PHENYLALANINE 238 mg
HISTIDINE 204 mg
THREONINE 179 mg
METHIONINE 170 mg
TRYPTOPHAN 77 mg
NONESSENTIAL AMINO ACIDS
ALANINE 880 mg
ARGININE 489 mg
GLYCINE 438 mg
PROLINE 289 mg
SERINE 213 mg
TYROSINE 17 mg
ELECTROLYTES
SODIUM ACETATE TRIHYDRATE USP 297 mg
DIBASIC POTASSIUM PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 77 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE DIHYDRATE USP 33 mg
mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 70
CHLORIDE 39
PHOSPHATE 30 (15 mmol/L)
BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
STERILE
SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 US



Container Label

LOT EXP

2B7738 NDC 0338-1145-03

CLINIMIX E
WITH ELECTROLYTES 4.25/10

SULFITE-FREE
(4.25% Amino Acid
with Electrolytes in
10% Dextrose with
Calcium) Injection

500 mL INJECTION PORT CHAMBER
20% Dextrose Injection with Calcium

500 mL OUTLET PORT CHAMBER
8.5% Amino Acid Injection

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESRCIBING INFORMATION FOR INSTRUCTIONS
ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE
QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED
INJECTION

DEXTROSE HYDROUS USP 10 g
ESSENTIAL AMINO ACIDS
LEUCINE 311 mg
ISOLEUCINE 255 mg
VALINE 247 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT) 247 mg
PHENYLALANINE 238 mg
HISTIDINE 204 mg
THREONINE 179 mg
METHIONINE 170 mg
TRYPTOPHAN 77 mg
NONESSENTIAL AMINO ACIDS
ALANINE 880 mg
ARGININE 489 mg
GLYCINE 438 mg
PROLINE 289 mg
SERINE 213 mg
TYROSINE 17 mg
ELECTROLYTES
SODIUM ACETATE TRIHYDRATE USP 297 mg
DIBASIC POTASSIUM PHOSPHATE USP 261 mg

SODIUM CHLORIDE USP 77 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE DIHYDRATE USP 33 mg
mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 70
CHLORIDE 39
PHOSPHATE 30 (15 mmol/L)
BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
STERILE
SINGLE DOSE CONTAINER
ROOM TEMPERATURE (25°C/77°F)
AVOID EXCESSIVE HEAT
PROTECT FROM FREEZING
SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 US



Container Label

LOT EXP

2B7717 NDC 0338-1115-04

CLINIMIX E
WITH ELECTROLYTES 4.25/10

SULFITE-FREE
(4.25% Amino Acid
with Electrolytes in
10% Dextrose with
Calcium) Injection

CENTRAL LINE INFUSION ONLY

1000 mL INJECTION PORT CHAMBER
20% Dextrose Injection with Calcium

1000 mL OUTLET PORT CHAMBER
8.5% Amino Acid Injection

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESRCIBING INFORMATION FOR INSTRUCTIONS
ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE
QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED
INJECTION

DEXTROSE HYDROUS USP 10 g
ESSENTIAL AMINO ACIDS
LEUCINE 311 mg
ISOLEUCINE 255 mg
VALINE 247 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT) 247 mg
PHENYLALANINE 238 mg
HISTIDINE 204 mg
THREONINE 179 mg
METHIONINE 170 mg
TRYPTOPHAN 77 mg
NONESSENTIAL AMINO ACIDS
ALANINE 880 mg
ARGININE 489 mg
GLYCINE 438 mg
PROLINE 289 mg
SERINE 213 mg
TYROSINE 17 mg
ELECTROLYTES
SODIUM ACETATE TRIHYDRATE USP 297 mg
DIBASIC POTASSIUM PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 77 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE DIHYDRATE USP 33 mg
mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 70
CHLORIDE 39
PHOSPHATE 30 (15 mmol/L)
BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
STERILE
SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 US



Container Label

LOT EXP

2B7740 NDC 0338-1147-03

CLINIMIX E
WITH ELECTROLYTES 5/15

SULFITE-FREE
(5% Amino Acid
with Electrolytes in
15% Dextrose with
Calcium) Injection

CENTRAL LINE INFUSION ONLY

500 mL INJECTION PORT CHAMBER
30% Dextrose Injection with Calcium

500 mL OUTLET PORT CHAMBER
10% Amino Acid Injection

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESRCIBING INFORMATION FOR INSTRUCTIONS
ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE
QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED
INJECTION

DEXTROSE HYDROUS USP 15 g
ESSENTIAL AMINO ACIDS
LEUCINE 365 mg
ISOLEUCINE 300 mg
VALINE 290 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT) 290 mg
PHENYLALANINE 280 mg
HISTIDINE 240 mg
THREONINE 210 mg
METHIONINE 200 mg
TRYPTOPHAN 90 mg
NONESSENTIAL AMINO ACIDS
ALANINE 1035 mg
ARGININE 575 mg
GLYCINE 515 mg
PROLINE 340 mg
SERINE 250 mg
TYROSINE 20 mg
ELECTROLYTES
SODIUM ACETATE TRIHYDRATE USP 340 mg
DIBASIC POTASSIUM PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 59 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE DIHYDRATE USP 33 mg
mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 80
CHLORIDE 39
PHOSPHATE 30 (15 mmol/L)
BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
STERILE
SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)
AVOID EXCESSIVE HEAT
PROTECT FROM FREEZING
SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 US



Container Label

Container Label

LOT EXP

2B7721 NDC 0338-1123-04

CLINIMIX E
WITH ELECTROLYTES 5/15

SULFITE-FREE
(5% Amino Acid
with Electrolytes in
15% Dextrose with
Calcium) Injection

1000 mL INJECTION PORT CHAMBER
30% Dextrose Injection with Calcium

1000 mL OUTLET PORT CHAMBER
10% Amino Acid Injection

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESRCIBING INFORMATION FOR INSTRUCTIONS
ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE
QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED
INJECTION

DEXTROSE HYDROUS USP 15 g
ESSENTIAL AMINO ACIDS
LEUCINE 365 mg
ISOLEUCINE 300 mg
VALINE 290 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT) 290 mg
PHENYLALANINE 280 mg
HISTIDINE 240 mg
THREONINE 210 mg
METHIONINE 200 mg
TRYPTOPHAN 90 mg
NONESSENTIAL AMINO ACIDS
ALANINE 1035 mg
ARGININE 575 mg
GLYCINE 515 mg
PROLINE 340 mg
SERINE 250 mg
TYROSINE 20 mg

ELECTROLYTES
SODIUM ACETATE TRIHYDRATE USP 340 mg
DIBASIC POTASSIUM PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 59 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE DIHYDRATE USP 33 mg
mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 80
CHLORIDE 39
PHOSPHATE 30 (15 mmol/L)
BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
STERILE
SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 US



Container Label

LOT EXP

2B7741 NDC 0338-1148-03

CLINIMIX E
WITH ELECTROLYTES 5/20

SULFITE-FREE
(5% Amino Acid
with Electrolytes in
20% Dextrose with
Calcium) Injection

500 mL INJECTION PORT CHAMBER
40% Dextrose Injection with Calcium

500 mL OUTLET PORT CHAMBER
10% Amino Acid Injection

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESRCIBING INFORMATION FOR INSTRUCTIONS
ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED INJECTION

DEXTROSE HYDROUS USP 20 g
ESSENTIAL AMINO ACIDS
LEUCINE 365 mg
ISOLEUCINE 300 mg
VALINE 290 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT) 290 mg
PHENYLALANINE 280 mg
HISTIDINE 240 mg
THREONINE 210 mg
METHIONINE 200 mg
TRYPTOPHAN 90 mg
NONESSENTIAL AMINO ACIDS
ALANINE 1035 mg
ARGININE 575 mg
GLYCINE 515 mg
PROLINE 340 mg
SERINE 250 mg
TYROSINE 20 mg
ELECTROLYTES
SODIUM ACETATE TRIHYDRATE USP 340 mg
DIBASIC POTASSIUM PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 59 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE DIHYDRATE USP 33 mg
mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 80
CHLORIDE 39
PHOSPHATE 30 (15 mmol/L)
BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
STERILE
SINGLE DOSE CONTAINER
ROOM TEMPERATURE (25°C/77°F)
AVOID EXCESSIVE HEAT
PROTECT FROM FREEZING
SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 US



Container Label

LOT EXP

2B7722 NDC 0338-1125-04

CLINIMIX E
WITH ELECTROLYTES 5/20

SULFITE-FREE
(5% Amino Acid
with Electrolytes in
20% Dextrose with
Calcium) Injection

1000 mL INJECTION PORT CHAMBER
40% Dextrose Injection with Calcium

1000 mL OUTLET PORT CHAMBER
10% Amino Acid Injection

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESRCIBING INFORMATION FOR INSTRUCTIONS
ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS

A SLIGHT YELLOW COLOR DOES NOT ALTER THE
QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED
INJECTION

DEXTROSE HYDROUS USP 20 g
ESSENTIAL AMINO ACIDS
LEUCINE 365 mg
ISOLEUCINE 300 mg
VALINE 290 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT) 290 mg
PHENYLALANINE 280 mg
HISTIDINE 240 mg
THREONINE 210 mg
METHIONINE 200 mg
TRYPTOPHAN 90 mg
NONESSENTIAL AMINO ACIDS
ALANINE 1035 mg
ARGININE 575 mg
GLYCINE 515 mg
PROLINE 340 mg
SERINE 250 mg
TYROSINE 20 mg
ELECTROLYTES
SODIUM ACETATE TRIHYDRATE USP 340 mg
DIBASIC POTASSIUM PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 59 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE DIHYDRATE USP 33 mg
mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 80
CHLORIDE 39
PHOSPHATE 30 (15 mmol/L)
BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
STERILE
SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 US



EADB9963 1000 mL
NDC 0338-0202-01
CLINIMIX E
WITH ELECTROLYTES 8/14
SULFITE-FREE
(8% Amino Acids
with Electrolytes in
14% Dextrose with
Calcium) Injection
CENTRAL LINE INFUSION ONLY
360 mL INJECTION PORT CHAMBER
39% Dextrose Injection with Calcium
640 mL OUTLET PORT CHAMBER
12.5% Amino Acid Injection
with Electrolytes
Rx Only
ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE
SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON ACTIVATION
AFTER MIXING THE PRODUCT REPRESENTS 1000 mL
REFRIGERATED STORAGE IS LIMITED TO 9 DAYS ONCE OVERWRAP IS OPENED
A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY AND EFFICACY OF THIS PRODUCT
ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY
EXP

CONTENTS OF EACH 100 mL OF THE ADMIXED INJECTION
DEXTROSE HYDROUS USP 14 g
ESSENTIAL AMINO ACIDS
LEUCINE 584 mg
ISOLEUCINE 480 mg
VALINE 464 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT) 464 mg
PHENYLALANINE 448 mg
HISTIDINE 384 mg
THREONINE 336 mg
METHIONINE 320 mg
TRYPTOPHAN 144 mg
NONESSENTIAL AMINO ACIDS
ALANINE 1656 mg
ARGININE 920 mg
GLYCINE 824 mg
PROLINE 544 mg
SERINE 400 mg
TYROSINE 32 mg
ELECTROLYTES
DIBASIC POTASSIUM PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 205 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE DIHYDRATE USP 33 mg
mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 83
CHLORIDE 76
PHOSPHATE 30 (15 mmol/L)
BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID AND HYDROCHLORIC ACID
STERILE
SINGLE DOSE CONTAINER
STORE AT ROOM TEMPERATURE (25°C/77°F) IN UNOPENED OVERWRAP
AVOID EXCESSIVE HEAT
PROTECT FROM FREEZING
SEE PRESCRIBING INFORMATION
Baxter
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM
LOT
BE-35-04-044

Container Label

EADB9963

1000 mL

NDC 0338-0202-01

CLINIMIX E
WITH ELECTROLYTES 8/14

SULFITE-FREE
(8% Amino Acids
with Electrolytes in
14% Dextrose with
Calcium) Injection

CENTRAL LINE INFUSION ONLY

360 mL INJECTION PORT CHAMBER
39% Dextrose Injection with Calcium

640 mL OUTLET PORT CHAMBER

12.5% Amino Acid Injection

with Electrolytes

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

2000 mL

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS
ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS
ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT ALTER THE
QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

EXP

CONTENTS OF EACH 100 mL
OF THE ADMIXED INJECTION
DEXTROSE HYDROUS USP 14 g
ESSENTIAL AMINO ACIDS
LEUCINE 584 mg
ISOLEUCINE 480 mg
VALINE 464 mg
LYSINE (ADDED AS THE
HYDROCHLORIDE SALT) 464 mg
PHENYLALANINE 448 mg
HISTIDINE 384 mg
THREONINE 336 mg

METHIONINE 320 mg
TRYPTOPHAN 144 mg
NONESSENTIAL AMINO ACIDS
ALANINE 1656 mg
ARGININE 920 mg
GLYCINE 824 mg
PROLINE 544 mg
SERINE 400 mg
TYROSINE 32 mg
ELECTROLYTES
DIBASIC POTASSIUM
PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 205 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE
DIHYDRATE USP 33 mg
mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 83
CHLORIDE 76
PHOSPHATE 30 (15 mmol/L)

BALANCED BY IONS FROM AMINO ACIDS

pH ADJUSTED WITH GLACIAL ACETIC ACID AND HYDROCHLORIC ACID

STERILE

SINGLE DOSE CONTAINER

STORE AT ROOM TEMPERATURE (25ºC/77ºF)
IN UNOPENED OVERWRAP

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

BAXTER LOGo
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

LOT

BE-35-04-044

EADB9965 2000 mL

NDC 0338-0206-01

CLINIMIX E WITH ELECTROLYTES 8/14

SULFITE-FREE

(8% Amino Acids with Electrolytes in 14% Dextrose with Calcium) Injection

CENTRAL LINE INFUSION ONLY

720 mL INJECTION PORT CHAMBER
39% Dextrose Injection with Calcium

1280 mL OUTLET PORT CHAMBER
12.5% Amino Acid Injection
with Electrolytes

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED INJECTION

DEXTROSE HYDROUS USP	14 g
ESSENTIAL AMINO ACIDS	
LEUCINE	584 mg
ISOLEUCINE	480 mg
VALINE	464 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT)	464 mg
PHENYLALANINE	448 mg
HISTIDINE	384 mg
THREONINE	336 mg
METHIONINE	320 mg
TRYPTOPHAN	144 mg
NONESSENTIAL AMINO ACIDS	
ALANINE	1656 mg
ARGININE	920 mg
GLYCINE	824 mg
PROLINE	544 mg
SERINE	400 mg
TYROSINE	32 mg
ELECTROLYTES	
DIBASIC POTASSIUM PHOSPHATE USP	261 mg
SODIUM CHLORIDE USP	205 mg
MAGNESIUM CHLORIDE USP	51 mg
CALCIUM CHLORIDE DIHYDRATE USP	33 mg
mEq/L	
SODIUM	35
POTASSIUM	30
MAGNESIUM	5
CALCIUM	4.5 (2.2 mmol/L)
ACETATE	83
CHLORIDE	76
PHOSPHATE	30 (15 mmol/L)

BALANCED BY IONS FROM AMINO ACIDS

pH ADJUSTED WITH GLACIAL ACETIC ACID AND HYDROCHLORIC ACID

STERILE
SINGLE DOSE CONTAINER

STORE AT ROOM TEMPERATURE (25°C/77°F) IN UNOPENED OVERWRAP

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

Baxter

BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

BE-35-04-048

Exp Lot

Container Label

EADB9965

2000 mL

NDC 0338-0206-01

CLINIMIX E
WITH ELECTROLYTES 8/14

SULFITE-FREE
(8% Amino Acids
with Electrolytes in
14% Dextrose with
Calcium) Injection

CENTRAL LINE INFUSION ONLY

720 mL INJECTION PORT CHAMBER

39% Dextrose Injection with Calcium

1280 mL OUTLET PORT CHAMBER

12.5% Amino Acid Injection
with Electrolytes

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS
ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY AND
EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

EADB9965

Exp

CONTENTS OF EACH 100 mL OF THE
ADMIXED INJECTION
DEXTROSE HYDROUS USP 14 g

ESSENTIAL AMINO ACIDS
LEUCINE 584 mg
ISOLEUCINE 480 mg
VALINE 464 mg
LYSINE (ADDED AS THE
HYDROCHLORIDE SALT) 464 mg
PHENYLALANINE 448 mg
HISTIDINE 384 mg
THREONINE 336 mg
METHIONINE 320 mg
TRYPTOPHAN 144 mg

NONESSENTIAL AMINO ACIDS
ALANINE 1656 mg
ARGININE 920 mg
GLYCINE 824 mg
PROLINE 544 mg
SERINE 400 mg
TYROSINE 32 mg

ELECTROLYTES
DIBASIC POTASSIUM
PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 205 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE
DIHYDRATE USP 33 mg

mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 83
CHLORIDE 76
PHOSPHATE 30 (15 mmol/L)

BALANCED BY IONS FROM AMINO ACIDS

pH ADJUSTED WITH GLACIAL ACETIC ACID
AND HYDROCHLORIC ACID

STERILE

SINGLE DOSE CONTAINER

STORE AT ROOM TEMPERATURE (25ºC/77ºF)
IN UNOPENED OVERWRAP

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

Lot

BE-35-04-048



Container Label

EADB9943

1000 mL

NDC 0338-0210-01

CLINIMIX E
WITH ELECTROLYTES 8/10

SULFITE-FREE
(8% Amino Acids
with Electrolytes in
10% Dextrose with
Calcium) Injection

CENTRAL LINE INFUSION ONLY

360 mL INJECTION PORT CHAMBER
28% Dextrose Injection with Calcium

640 mL OUTLET PORT CHAMBER
12.5% Amino Acid Injection with Electrolytes

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS
ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS
ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT ALTER THE
QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

EXP

CONTENTS OF EACH 100 mL
OF THE ADMIXED INJECTION
DEXTROSE HYDROUS USP 10 g

ESSENTIAL AMINO ACIDS
LEUCINE 584 mg
ISOLEUCINE 480 mg
VALINE 464 mg
LYSINE (ADDED AS THE
HYDROCHLORIDE SALT) 464 mg
PHENYLALANINE 448 mg
HISTIDINE 384 mg
THREONINE 336 mg
METHIONINE 320 mg
TRYPTOPHAN 144 mg

NONESSENTIAL AMINO ACIDS

ALANINE 1656 mg
ARGININE 920 mg
GLYCINE 824 mg
PROLINE 544 mg
SERINE 400 mg
TYROSINE 32 mg

ELECTROLYTES
DIBASIC POTASSIUM
PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 205 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE
DIHYDRATE USP 33 mg

mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 83
CHLORIDE 76
PHOSPHATE 30 (15 mmol/L)

BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
AND HYDROCHLORIC ACID
STERILE
SINGLE DOSE CONTAINER
STORE AT ROOM TEMPERATURE (25ºC/77ºF)
IN UNOPENED OVERWRAP
AVOID EXCESSIVE HEAT
PROTECT FROM FREEZING
SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

LOT

BE-35-04-042

EADB9945 2000 mL

NDC 0338-0214-01

CLINIMIX E WITH ELECTROLYTES 8/10

SULFITE-FREE

(8% Amino Acids with Electrolytes in 10% Dextrose with Calcium) Injection

CENTRAL LINE INFUSION ONLY

720 mL INJECTION PORT CHAMBER
28% Dextrose Injection with Calcium

1280 mL OUTLET PORT CHAMBER
12.5% Amino Acid Injection
with Electrolytes

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED INJECTION

DEXTROSE HYDROUS USP	10 g
ESSENTIAL AMINO ACIDS	
LEUCINE	584 mg
ISOLEUCINE	480 mg
VALINE	464 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT)	464 mg
PHENYLALANINE	448 mg
HISTIDINE	384 mg
THREONINE	336 mg
METHIONINE	320 mg
TRYPTOPHAN	144 mg
NONESSENTIAL AMINO ACIDS	
ALANINE	1656 mg
ARGININE	920 mg
GLYCINE	824 mg
PROLINE	544 mg
SERINE	400 mg
TYROSINE	32 mg
ELECTROLYTES	
DIBASIC POTASSIUM PHOSPHATE USP	261 mg
SODIUM CHLORIDE USP	205 mg
MAGNESIUM CHLORIDE USP	51 mg
CALCIUM CHLORIDE DIHYDRATE USP	33 mg
mEq/L	
SODIUM	35
POTASSIUM	30
MAGNESIUM	5
CALCIUM	4.5 (2.2 mmol/L)
ACETATE	83
CHLORIDE	76
PHOSPHATE	30 (15 mmol/L)

BALANCED BY IONS FROM AMINO ACIDS

pH ADJUSTED WITH GLACIAL ACETIC ACID AND HYDROCHLORIC ACID

STERILE
SINGLE DOSE CONTAINER

STORE AT ROOM TEMPERATURE (25°C/77°F) IN UNOPENED OVERWRAP

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

Baxter

BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

BE-35-04-046

Exp Lot

Container Label

EADB9945

2000 mL

NDC 0338-0214-01

CLINIMIX E
WITH ELECTROLYTES 8/10

SULFITE-FREE
(8% Amino Acids
with Electrolytes in
10% Dextrose with
Calcium) Injection

CENTRAL LINE INFUSION ONLY

720 mL INJECTION PORT CHAMBER

28% Dextrose Injection with Calcium

1280 mL OUTLET PORT CHAMBER

12.5% Amino Acid Injection
with Electrolytes

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS
ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY AND
EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

Exp

CONTENTS OF EACH 100 mL OF THE
ADMIXED INJECTION
DEXTROSE HYDROUS USP 10 g

ESSENTIAL AMINO ACIDS
LEUCINE 584 mg
ISOLEUCINE 480 mg
VALINE 464 mg
LYSINE (ADDED AS THE
HYDROCHLORIDE SALT) 464 mg
PHENYLALANINE 448 mg
HISTIDINE 384 mg
THREONINE 336 mg
METHIONINE 320 mg
TRYPTOPHAN 144 mg

NONESSENTIAL AMINO ACIDS
ALANINE 1656 mg
ARGININE 920 mg
GLYCINE 824 mg
PROLINE 544 mg
SERINE 400 mg
TYROSINE 32 mg

ELECTROLYTES
DIBASIC POTASSIUM
PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 205 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE
DIHYDRATE USP 33 mg

mEq/L
SODIUM35
POTASSIUM30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 83
CHLORIDE 76
PHOSPHATE 30 (15 mmol/L)

BALANCED BY IONS FROM AMINO ACIDS

pH ADJUSTED WITH GLACIAL ACETIC ACID
AND HYDROCHLORIC ACID

STERILE

SINGLE DOSE CONTAINER

STORE AT ROOM TEMPERATURE (25ºC/77ºF)
IN UNOPENED OVERWRAP

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

Lot

BE35-04-046

2B7735L 1000 mL
NDC 0338-7020-01

(01) 00303387020013

CLINIMIX E
WITH ELECTROLYTES 2.75/5
SULFITE-FREE

(2.75% Amino Acids with Electrolytes in 5% Dextrose with Calcium) Injection

500 mL INJECTION PORT CHAMBER
10% Dextrose Injection with Calcium

500 mL OUTLET PORT CHAMBER
5.5% Amino Acid Injection with Electrolytes

2D Datamatrix (GS1):
NDC, EXP, LOT

ACTIVATE SEAL AND MIX THOROUGHLY BEFORE USE

2D Datamatrix Barcode

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY
Rx Only

EXP

CONTENTS OF EACH 100 mL OF THE ADMIXED INJECTION

DEXTROSE HYDROUS USP	5 g	1600
ESSENTIAL AMINO ACIDS		
LEUCINE	201 mg	
ISOLEUCINE	165 mg	
VALINE	160 mg	
LYSINE (ADDED AS THE HYDROCHLORIDE SALT)	159 mg	
PHENYLALANINE	154 mg	
HISTIDINE	132 mg	
THREONINE	116 mg	
METHIONINE	110 mg	
TRYPTOPHAN	50 mg	
NONESSENTIAL AMINO ACIDS		
ALANINE	570 mg	1000
ARGININE	316 mg	
GLYCINE	283 mg	
PROLINE	187 mg	
SERINE	138 mg	
TYROSINE	11 mg	800
ELECTROLYTES		
SODIUM ACETATE TRIHYDRATE USP	216 mg	
DIBASIC POTASSIUM PHOSPHATE USP	261 mg	
SODIUM CHLORIDE USP	112 mg	
MAGNESIUM CHLORIDE USP	51 mg	
CALCIUM CHLORIDE DIHYDRATE USP	33 mg	600

mEq/L	
SODIUM	35
POTASSIUM	30
MAGNESIUM	5
CALCIUM	4.5 (2.2 mmol/L)
ACETATE	51
CHLORIDE	39
PHOSPHATE	30 (15 mmol/L)

BALANCED BY IONS FROM AMINO ACIDS 400
pH ADJUSTED WITH GLACIAL ACETIC ACID AND HYDROCHLORIC ACID

STERILE SINGLE DOSE CONTAINER

STORE AT ROOM TEMPERATURE (25°C/77°F)
IN UNOPENED OVERWRAP
AVOID EXCESSIVE HEAT
PROTECT FROM FREEZING
SEE PRESCRIBING INFORMATION

BE-35-05-068

Baxter
BAXTER HEALTHCARE CORPORATION 200
DEERFIELD IL 60015 USA
MADE IN BELGIUM

LOT

Container Label

2B7735L
NDC 0338-7020-01
1000 mL

Barcode
(01) 00303387020013

CLINIMIX E
WITH ELECTROLYTES 2.75/5
SULFITE-FREE

(2.75% Amino Acids with Electrolytes in 5% Dextrose with Calcium) Injection

500 mL INJECTION PORT CHAMBER
10% Dextrose Injection with Calcium

500 mL OUTLET PORT CHAMBER
5.5% Amino Acid Injection

Rx Only

2D Datamatrix (GS1):
NDC, LOT, EXP

2D Datamatrix
Barcode

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESRCIBING INFORMATION FOR
INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT
REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO
9 DAYS ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT
ALTER THE QUALITY AND EFFICACY OF
THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE
COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED INJECTION
DEXTROSE HYDROUS USP 5 g
ESSENTIAL AMINO ACIDS
LEUCINE 201 mg
ISOLEUCINE 165 mg
VALINE 160 mg
LYSINE (ADDED AS THE
HYDROCHLORIDE SALT) 159 mg
PHENYLALANINE 154 mg
HISTIDINE 132 mg
THREONINE 116 mg

METHIONINE 110 mg
TRYPTOPHAN 50 mg
NONESSENTIAL AMINO ACIDS
ALANINE 570 mg
ARGININE 316 mg
GLYCINE 283 mg
PROLINE 187 mg
SERINE 138 mg
TYROSINE 11 mg
ELECTROLYTES
SODIUM ACETATE TRIHYDRATE USP 216 mg
DIBASIC POTASSIUM PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 112 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE DIHYDRATE USP 33 mg
mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 51
CHLORIDE 39
PHOSPHATE 30 (15 mmol/L)
BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
STERILE SINGLE DOSE CONTAINER
STORE AT ROOM TEMPERATURE (25°C/77°F)
AVOID EXCESSIVE HEAT
PROTECT FROM FREEZING
SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

EXP LOT

1600

1000

800

600

400

BE-35-05-069

200



Container Label

2B7737L
NDC 0338-7022-03
1000 mL

Barcode
(01) 00303387022017

CLINIMIX E
WITH ELECTROLYTES 4.25/5
SULFITE-FREE
(4.25% Amino Acids with Electrolytes in
5% Dextrose with Calcium) Injection

500 mL INJECTION PORT CHAMBER
10% Dextrose Injection with Calcium

500 mL OUTLET PORT CHAMBER
8.5% Amino Acid Injection with Electrolytes

Rx Only

2D Datamatrix (GS1):
NDC, LOT, EXP

2D Datamatrix
Barcode

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESRCIBING INFORMATION FOR
INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT
REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED
TO 9 DAYS ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT
ALTER THE QUALITY AND EFFICACY OF
THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE
COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED INJECTION
DEXTROSE HYDROUS USP 5 g
ESSENTIAL AMINO ACIDS
LEUCINE 311 mg
ISOLEUCINE 255 mg
VALINE 247 mg
LYSINE (ADDED AS THE
HYDROCHLORIDE SALT) 247 mg
PHENYLALANINE 238 mg
HISTIDINE 204 mg
THREONINE 179 mg
METHIONINE 170 mg
TRYPTOPHAN 77 mg
NONESSENTIAL AMINO ACIDS
ALANINE 880 mg
ARGININE 489 mg
GLYCINE 438 mg
PROLINE 289 mg
SERINE 213 mg
TYROSINE 17 mg
ELECTROLYTES
SODIUM ACETATE TRIHYDRATE USP 297 mg
DIBASIC POTASSIUM PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 77 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE DIHYDRATE USP 33 mg
mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 70
CHLORIDE 39
PHOSPHATE 30 (15 mmol/L)
BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
AND HYDROCHLORIC ACID
STERILE SINGLE DOSE CONTAINER
STORE AT ROOM TEMPERATURE (25°C/77°F)
IN UNOPENED OVERWRAP
AVOID EXCESSIVE HEAT
PROTECT FROM FREEZING
SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

EXP LOT

1600

1000

800

600

400

BE-35-05-069

200

2B7716L 2000 mL
NDC 0338-7024-01

(01)00303387024011

CLINIMIX E
WITH ELECTROLYTES 4.25/5

SULFITE-FREE

(4.25% Amino Acids with Electrolytes in 5% Dextrose with Calcium) Injection

1000 mL INJECTION PORT CHAMBER
10% Dextrose Injection with Calcium

1000 mL OUTLET PORT CHAMBER
8.5% Amino Acid Injection with Electrolytes

ACTIVATE SEAL AND MIX THOROUGHLY BEFORE USE

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

Rx Only

2D Datamatrix (GS1):
NDC, EXP, LOT

2D Datamatrix Barcode

EXP LOT

CONTENTS OF EACH 100 mL OF THE ADMIXED INJECTION

DEXTROSE HYDROUS USP	5 g
ESSENTIAL AMINO ACIDS	
LEUCINE	311 mg
ISOLEUCINE	255 mg
VALINE	247 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT)	247 mg
PHENYLALANINE	238 mg
HISTIDINE	204 mg
THREONINE	179 mg
METHIONINE	170 mg
TRYPTOPHAN	77 mg
NONESSENTIAL AMINO ACIDS	
ALANINE	880 mg
ARGININE	489 mg
GLYCINE	438 mg
PROLINE	289 mg
SERINE	213 mg
TYROSINE	17 mg
ELECTROLYTES	
SODIUM ACETATE TRIHYDRATE USP	297 mg
DIBASIC POTASSIUM PHOSPHATE USP	261 mg
SODIUM CHLORIDE USP	77 mg
MAGNESIUM CHLORIDE USP	51 mg
CALCIUM CHLORIDE DIHYDRATE USP	33 mg
mEq/L	
SODIUM	35
POTASSIUM	30
MAGNESIUM	5
CALCIUM	4.5 (2.2 mmol/L)
ACETATE	70
CHLORIDE	39
PHOSPHATE	30 (15 mmol/L)

2000

1600

1000

800

BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID AND HYDROCHLORIC ACID

STERILE
SINGLE DOSE CONTAINER

600

STORE AT ROOM TEMPERATURE (25°C/77°F) IN UNOPENED OVERWRAP

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

400

Baxter

BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

BE-35-05-064

200

Container Label

2B7716L
NDC 0338-7024-01
2000 mL

Barcode
(01) 00303387024011

CLINIMIX E
WITH ELECTROLYTES 4.25/5

SULFITE-FREE

(4.25% Amino Acids with Electrolytes in 5% Dextrose with Calcium) Injection

1000 mL INJECTION PORT CHAMBER

10% Dextrose Injection with Calcium

1000 mL OUTLET PORT CHAMBER
8.5% Amino Acid Injection with
Electrolytes

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESRCIBING INFORMATION FOR INSTRUCTIONS
ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO
9 DAYS ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT ALTER THE
QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

2D Datamatrix (GS1):
NDC, LOT, EXP

2D Datamatrix
Barcode

CONTENTS OF EACH 100 mL OF THE ADMIXED
INJECTION
DEXTROSE HYDROUS USP 5 g

ESSENTIAL AMINO ACIDS
LEUCINE 311 mg
ISOLEUCINE 255 mg
VALINE 247 mg
LYSINE (ADDED AS
THE HYDROCHLORIDE SALT) 247 mg
PHENYLALANINE 238 mg
HISTIDINE 204 mg
THREONINE 179 mg
METHIONINE 170 mg
TRYPTOPHAN 77 mg

NONESSENTIAL AMINO ACIDS
ALANINE 880 mg
ARGININE 489 mg
GLYCINE 438 mg
PROLINE 289 mg
SERINE 213 mg
TYROSINE 17 mg

ELECTROLYTES
SODIUM ACETATE
TRIHYDRATE USP 297 mg
DIBASIC POTASSIUM
PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 77 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE
DIHYDRATE USP 33 mg

mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 70
CHLORIDE 39
PHOSPHATE 30 (15 mmol/L)

BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
AND HYDROCHLORIC ACID

STERILE
SINGLE DOSE CONTAINER

STORE AT ROOM TEMPERATURE (25°C/77°F)
IN UNOPENED OVERWRAP

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

EXP LOT

2000

1600

1000

800

600

400

BE-35-05-064

200



Container Label

2B7738L
NDC 0338-7026-01
1000 mL

Barcode
(01) 00303387026015

CLINIMIX E
WITH ELECTROLYTES 4.25/10

SULFITE-FREE

(4.25% Amino Acids with Electrolytes in
10% Dextrose with Calcium) Injection

CENTRAL LINE INFUSION ONLY

500 mL INJECTION PORT CHAMBER
20% Dextrose Injection with Calcium

500 mL OUTLET PORT CHAMBER
8.5% Amino Acid Injection with Electrolytes

Rx Only

2D Datamatrix (GS1):
NDC, LOT, EXP

2D Datamatrix
Barcode

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESRCIBING INFORMATION FOR
INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT
REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO
9 DAYS ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT
ALTER THE QUALITY AND EFFICACY
OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE
COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED
INJECTION
DEXTROSE HYDROUS USP 10 g

ESSENTIAL AMINO ACIDS
LEUCINE 311 mg
ISOLEUCINE 255 mg
VALINE 247 mg
LYSINE (ADDED AS THE
HYDROCHLORIDE SALT) 247 mg
PHENYLALANINE 238 mg
HISTIDINE 204 mg
THREONINE 179 mg
METHIONINE 170 mg
TRYPTOPHAN 77 mg

NONESSENTIAL AMINO ACIDS
ALANINE 880 mg
ARGININE 489 mg
GLYCINE 438 mg
PROLINE 289 mg
SERINE 213 mg
TYROSINE 17 mg

ELECTROLYTES
SODIUM ACETATE TRIHYDRATE USP 297 mg
DIBASIC POTASSIUM PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 77 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE DIHYDRATE USP 33 mg

mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 70
CHLORIDE 39
PHOSPHATE 30 (15 mmol/L)
BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
AND HYDROCHLORIC ACID

STERILE SINGLE DOSE CONTAINER

STORE AT ROOM TEMPERATURE (25°C/77°F)
IN UNOPENED OVERWRAP
AVOID EXCESSIVE HEAT
PROTECT FROM FREEZING
SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

EXP LOT

1600

1000

800

600

400

BE-35-05-070

200

2B7717L 2000 mL
NDC 0338-7028-01

(01) 00303387028019

CLINIMIX E
WITH ELECTROLYTES 4.25/10

SULFITE-FREE

(4.25% Amino Acids with Electrolytes in 10% Dextrose with Calcium) Injection

CENTRAL LINE INFUSION ONLY

1000 mL INJECTION PORT CHAMBER
20% Dextrose Injection with Calcium

1000 mL OUTLET PORT CHAMBER
8.5% Amino Acid Injection with Electrolytes

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

Rx Only

2D Datamatrix (GS1):
NDC, EXP, LOT

2D Datamatrix Barcode

CONTENTS OF EACH 100 mL OF THE ADMIXED INJECTION

DEXTROSE HYDROUS USP	10 g
ESSENTIAL AMINO ACIDS	
LEUCINE	311 mg
ISOLEUCINE	255 mg
VALINE	247 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT)	247 mg
PHENYLALANINE	238 mg
HISTIDINE	204 mg
THREONINE	179 mg
METHIONINE	170 mg
TRYPTOPHAN	77 mg
NONESSENTIAL AMINO ACIDS	
ALANINE	880 mg
ARGININE	489 mg
GLYCINE	438 mg
PROLINE	289 mg
SERINE	213 mg
TYROSINE	17 mg
ELECTROLYTES	
SODIUM ACETATE TRIHYDRATE USP	297 mg
DIBASIC POTASSIUM PHOSPHATE USP	261 mg
SODIUM CHLORIDE USP	77 mg
MAGNESIUM CHLORIDE USP	51 mg
CALCIUM CHLORIDE DIHYDRATE USP	33 mg
mEq/L	
SODIUM	35
POTASSIUM	30
MAGNESIUM	5
CALCIUM	4.5 (2.2 mmol/L)
ACETATE	70
CHLORIDE	39
PHOSPHATE	30 (15 mmol/L)

2000

1600

1000

800

BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID AND HYDROCHLORIC ACID

STERILE
SINGLE DOSE CONTAINER

600

STORE AT ROOM TEMPERATURE (25°C/77°F) IN UNOPENED OVERWRAP

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

400

Baxter
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

BE-35-05-065

200

EXP LOT

Container Label

2B7717L
NDC 0338-7026-01
2000 mL

Barcode
(01) 00303387028019

CLINIMIX E
WITH ELECTROLYTES 4.25/5

SULFITE-FREE

(4.25% Amino Acids with Electrolytes in 5% Dextrose with Calcium) Injection

CENTRL LINE INFUSION ONLY

1000 mL INJECTION PORT CHAMBER
10% Dextrose Injection with Calcium

1000 mL OUTLET PORT CHAMBER
8.5% Amino Acid Injection with Electrolytes

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESRCIBING INFORMATION FOR INSTRUCTIONS

ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO
9 DAYS ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT ALTER THE
QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

2D Datamatrix (GS1):
NDC, LOT, EXP

2D Datamatrix
Barcode

CONTENTS OF EACH 100 mL OF THE ADMIXED
INJECTION
DEXTROSE HYDROUS USP 5 g

ESSENTIAL AMINO ACIDS
LEUCINE 311 mg
ISOLEUCINE 255 mg
VALINE 247 mg
LYSINE (ADDED AS
THE HYDROCHLORIDE SALT) 247 mg
PHENYLALANINE 238 mg
HISTIDINE 204 mg
THREONINE 179 mg
METHIONINE 170 mg
TRYPTOPHAN 77 mg

NONESSENTIAL AMINO ACIDS
ALANINE 880 mg
ARGININE 489 mg
GLYCINE 438 mg
PROLINE 289 mg
SERINE 213 mg
TYROSINE 17 mg

ELECTROLYTES
SODIUM ACETATE
TRIHYDRATE USP 297 mg
DIBASIC POTASSIUM
PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 77 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE
DIHYDRATE USP 33 mg

mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 70
CHLORIDE 39
PHOSPHATE 30 (15 mmol/L)

BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
AND HYDROCHLORIC ACID

STERILE
SINGLE DOSE CONTAINER

STORE AT ROOM TEMPERATURE (25°C/77°F)
IN UNOPENED OVERWRAP

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

EXP LOT

2000

1600

1000

800

600

400

BE-35-05-065

2B7740L 1000 mL
NDC 0338-7030-01

(01) 00303387030012

CLINIMIX E
WITH ELECTROLYTES 5/15

SULFITE-FREE
(5% Amino Acids with Electrolytes in
15% Dextrose with Calcium) Injection

CENTRAL LINE INFUSION ONLY

500 mL INJECTION PORT CHAMBER
30% Dextrose Injection with Calcium
500 mL OUTLET PORT CHAMBER
10% Amino Acid Injection with Electrolytes

2D Datamatrix (GS1):
NDC, EXP, LOT

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESCRIBING INFORMATION FOR
INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT
REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO
9 DAYS ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT
ALTER THE QUALITY AND EFFICACY OF
THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE
COMPATIBILITY
Rx Only

2D Datamatrix
Barcode

EXP

CONTENTS OF EACH 100 mL OF THE ADMIXED INJECTION

DEXTROSE HYDROUS USP	15 g	1600
ESSENTIAL AMINO ACIDS		
LEUCINE	365 mg	
ISOLEUCINE	300 mg	
VALINE	290 mg	
LYSINE (ADDED AS THE HYDROCHLORIDE SALT)	290 mg	
PHENYLALANINE	280 mg	
HISTIDINE	240 mg	
THREONINE	210 mg	
METHIONINE	200 mg	
TRYPTOPHAN	90 mg	
NONESSENTIAL AMINO ACIDS		
ALANINE	1035 mg	1000
ARGININE	575 mg	
GLYCINE	515 mg	
PROLINE	340 mg	
SERINE	250 mg	
TYROSINE	20 mg	800
ELECTROLYTES		
SODIUM ACETATE TRIHYDRATE USP	340 mg	
DIBASIC POTASSIUM PHOSPHATE USP	261 mg	
SODIUM CHLORIDE USP	59 mg	
MAGNESIUM CHLORIDE USP	51 mg	
CALCIUM CHLORIDE DIHYDRATE USP	33 mg	600

mEq/L	
SODIUM	35
POTASSIUM	30
MAGNESIUM	5
CALCIUM	4.5 (2.2 mmol/L)
ACETATE	80
CHLORIDE	39
PHOSPHATE	30 (15 mmol/L)

BALANCED BY IONS FROM AMINO ACIDS 400
pH ADJUSTED WITH GLACIAL ACETIC ACID
AND HYDROCHLORIC ACID

STERILE SINGLE DOSE CONTAINER

STORE AT ROOM TEMPERATURE (25°C/77°F)
IN UNOPENED OVERWRAP
AVOID EXCESSIVE HEAT
PROTECT FROM FREEZING
SEE PRESCRIBING INFORMATION

BE-35-05-071

Baxter
BAXTER HEALTHCARE CORPORATION 200
DEERFIELD IL 60015 USA
MADE IN BELGIUM

LOT

Container Label

2B7740L
NDC 0338-7030-01
1000 mL

Bacode
(01) 00303387030012

CLINIMIX E
WITH ELECTROLYTES 5/15

SULFITE-FREE
(5% Amino Acid with Electrolytes in
15% Dextrose with Calcium) Injection

CENTRAL LINE INFUSION ONLY

500 mL INJECTION PORT CHAMBER
30% Dextrose Injection with Calcium

500 mL OUTLET PORT CHAMBER
10% Amino Acid Injection

Rx Only

2D Datamatrix (GS1):
NDC, LOT, EXP

2D Datamatrix
Barcode

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESRCIBING INFORMATION FOR
INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT
REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO
9 DAYS ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT
ALTER THE QUALITY AND EFFICACY OF
THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE
COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED
INJECTION
DEXTROSE HYDROUS USP 15 g

ESSENTIAL AMINO ACIDS
LEUCINE 365 mg
ISOLEUCINE 300 mg

VALINE 290 mg
LYSINE (ADDED AS THE
HYDROCHLORIDE SALT) 290 mg
PHENYLALANINE 280 mg
HISTIDINE 240 mg
THREONINE 210 mg
METHIONINE 200 mg
TRYPTOPHAN 90 mg

NONESSENTIAL AMINO ACIDS
ALANINE 1035 mg
ARGININE 575 mg
GLYCINE 515 mg
PROLINE 340 mg
SERINE 250 mg
TYROSINE 20 mg

ELECTROLYTES
SODIUM ACETATE TRIHYDRATE USP 340 mg
DIBASIC POTASSIUM PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 59 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE DIHYDRATE USP 33 mg

mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 80
CHLORIDE 39
PHOSPHATE 30 (15 mmol/L)

BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
AND HYDROCHLORIC ACID

STERILE SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)
IN UNOPENED OVERWRAP
AVOID EXCESSIVE HEAT
PROTECT FROM FREEZING
SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

EXP LOT

1600

1000

800

600

400

BE-35-05-071

200

2B7721L 2000 mL
NDC 0338-7032-01

(01) 00303387032016

CLINIMIX E
WITH ELECTROLYTES 5/15

SULFITE-FREE

(5% Amino Acids with Electrolytes in 15% Dextrose with Calcium) Injection

CENTRAL LINE INFUSION ONLY

1000 mL INJECTION PORT CHAMBER
30% Dextrose Injection with Calcium

1000 mL OUTLET PORT CHAMBER
10% Amino Acid Injection with Electrolytes

ACTIVATE SEAL AND MIX THOROUGHLY BEFORE USE

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

Rx Only

2D Datamatrix (GS1): NDC, EXP, LOT

2D Datamatrix Barcode

EXP LOT

CONTENTS OF EACH 100 mL OF THE ADMIXED INJECTION

DEXTROSE HYDROUS USP	15 g	
ESSENTIAL AMINO ACIDS		
LEUCINE	365 mg	
ISOLEUCINE	300 mg	
VALINE	290 mg	
LYSINE (ADDED AS THE HYDROCHLORIDE SALT)	290 mg	
PHENYLALANINE	280 mg	2000
HISTIDINE	240 mg	
THREONINE	210 mg	
METHIONINE	200 mg	
TRYPTOPHAN	90 mg	
NONESSENTIAL AMINO ACIDS		
ALANINE	1035 mg	
ARGININE	575 mg	
GLYCINE	515 mg	
PROLINE	340 mg	
SERINE	250 mg	1600
TYROSINE	20 mg	
ELECTROLYTES		
SODIUM ACETATE TRIHYDRATE USP	340 mg	
DIBASIC POTASSIUM PHOSPHATE USP	261 mg	
SODIUM CHLORIDE USP	59 mg	1000
MAGNESIUM CHLORIDE USP	51 mg	
CALCIUM CHLORIDE DIHYDRATE USP	33 mg	
mEq/L		
SODIUM	35	
POTASSIUM	30	
MAGNESIUM	5	
CALCIUM	4.5 (2.2 mmol/L)	
ACETATE	80	800
CHLORIDE	39	
PHOSPHATE	30 (15 mmol/L)	

BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID AND HYDROCHLORIC ACID

STERILE
SINGLE DOSE CONTAINER 600

STORE AT ROOM TEMPERATURE (25°C/77°F) IN UNOPENED OVERWRAP

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION 400

Baxter
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

BE-35-05-066

200

Container Label

2B7721L
NDC 0338-7032-01
2000 mL

Barcode
(01) 00303387032016

CLINIMIX E
WITH ELECTROLYTES 5/15

SULFITE-FREE

(5% Amino Acids with Electrolytes in 15% Dextrose with Calcium) Injection

CENTRAL LINE INFUSION ONLY

1000 mL INJECTION PORT CHAMBER
30% Dextrose Injection with Calcium

1000 mL OUTLET PORT CHAMBER
10% Amino Acid Injection

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESRCIBING INFORMATION FOR INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO
9 DAYS ONCE OVERWARP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT ALTER THE
QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

2D Datamatrix (GS1):
NDC, LOT, EXP

2D Datamatrix
Barcode

CONTENTS OF EACH 100 mL OF THE ADMIXED
INJECTION
DEXTROSE HYDROUS USP 15

ESSENTIAL AMINO ACIDS
LEUCINE 365 mg
ISOLEUCINE 300 mg
VALINE 290 mg
LYSINE (ADDED AS THE
HYDROCHLORIDE SALT) 290 mg
PHENYLALANINE 280 mg
HISTIDINE 240 mg
THREONINE 210 mg
METHIONINE 200 mg
TRYPTOPHAN 90 mg

NONESSENTIAL AMINO ACIDS
ALANINE 1035 mg
ARGININE 575 mg
GLYCINE 515 mg
PROLINE 340 mg
SERINE 250 mg
TYROSINE 20 mg

ELECTROLYTES
SODIUM ACETATE
TRIHYDRATE USP 340 mg
DIBASIC POTASSIUM
PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 59 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE DIHYDRATE USP 33 mg

mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 80
CHLORIDE 39
PHOSPHATE 30 (15 mmol/L)

BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
AND HYDROCHLORIC ACID

STERILE
SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)
IN UNOPENED OVERWRAP

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

EXP LOT

2000

1600

1000

800

600

400

BE-35-05-066

200

2B7741L 1000 mL
NDC 0338-7034-01

(01)00303387034010

CLINIMIX E
WITH ELECTROLYTES 5/20

SULFITE-FREE
(5% Amino Acids with Electrolytes in 20% Dextrose with Calcium) Injection

CENTRAL LINE INFUSION ONLY

500 mL INJECTION PORT CHAMBER
40% Dextrose Injection with Calcium
500 mL OUTLET PORT CHAMBER
10% Amino Acid Injection with Electrolytes

2D Datamatrix (GS1):
NDC, EXP, LOT

ACTIVATE SEAL AND MIX THOROUGHLY BEFORE USE

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

Rx Only

2D Datamatrix Barcode

EXP

CONTENTS OF EACH 100 mL OF THE ADMIXED INJECTION

DEXTROSE HYDROUS USP	20 g	1600
ESSENTIAL AMINO ACIDS		
LEUCINE	365 mg	
ISOLEUCINE	300 mg	
VALINE	290 mg	
LYSINE (ADDED AS THE HYDROCHLORIDE SALT)	290 mg	
PHENYLALANINE	280 mg	
HISTIDINE	240 mg	
THREONINE	210 mg	
METHIONINE	200 mg	
TRYPTOPHAN	90 mg	
NONESSENTIAL AMINO ACIDS		
ALANINE	1035 mg	1000
ARGININE	575 mg	
GLYCINE	515 mg	
PROLINE	340 mg	
SERINE	250 mg	
TYROSINE	20 mg	800
ELECTROLYTES		
SODIUM ACETATE TRIHYDRATE USP	340 mg	
DIBASIC POTASSIUM PHOSPHATE USP	261 mg	
SODIUM CHLORIDE USP	59 mg	
MAGNESIUM CHLORIDE USP	51 mg	
CALCIUM CHLORIDE DIHYDRATE USP	33 mg	600

mEq/L	
SODIUM	35
POTASSIUM	30
MAGNESIUM	5
CALCIUM	4.5 (2.2 mmol/L)
ACETATE	80
CHLORIDE	39
PHOSPHATE	30 (15 mmol/L)

BALANCED BY IONS FROM AMINO ACIDS 400
pH ADJUSTED WITH GLACIAL ACETIC ACID
AND HYDROCHLORIC ACID

STERILE SINGLE DOSE CONTAINER

STORE AT ROOM TEMPERATURE (25°C/77°F)
IN UNOPENED OVERWRAP
AVOID EXCESSIVE HEAT
PROTECT FROM FREEZING
SEE PRESCRIBING INFORMATION

BE-35-05-072

Baxter
BAXTER HEALTHCARE CORPORATION 200
DEERFIELD IL 60015 USA
MADE IN BELGIUM
LOT

Container Label

2B7741L
NDC 0338-7034-01
1000 mL

Barcode
(01) 00303387034010

CLINIMIX E
WITH ELECTROLYTES 5/20

SULFITE-FREE

(5% Amino Acids with Electrolytes in
20% Dextrose with Calcium) Injection

CENTRAL LINE INFUSION ONLY

500 mL INJECTION PORT CHAMBER
40% Dextrose Injection with Calcium

500 mL OUTLET PORT CHAMBER
10% Amino Acid Injection

Rx Only

2D Datamatrix (GS1):
NDC, LOT, EXP

2D Datamatrix
Barcode

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESRCIBING INFORMATION FOR
INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT
REPRESENTS 1000 mL

REFRIGERATED STORAGE IS LIMITED TO
9 DAYS ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT
ALTER THE QUALITY AND EFFICACY OF
THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE
COMPATIBILITY

CONTENTS OF EACH 100 mL OF THE ADMIXED INJECTION
DEXTROSE HYDROUS USP 20 g

ESSENTIAL AMINO ACIDS
LEUCINE 365 mg
ISOLEUCINE 300 mg
VALINE 290 mg
LYSINE (ADDED AS THE
HYDROCHLORIDE SALT) 290 mg
PHENYLALANINE 280 mg

HISTIDINE 240 mg
THREONINE 210 mg
METHIONINE 200 mg
TRYPTOPHAN 90 mg

NONESSENTIAL AMINO ACIDS
ALANINE 1035 mg
ARGININE 575 mg
GLYCINE 515 mg
PROLINE 340 mg
SERINE 250 mg
TYROSINE 20 mg

ELECTROLYTES
SODIUM ACETATE TRIHYDRATE USP 340 mg
DIBASIC POTASSIUM PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 59 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE DIHYDRATE USP 33 mg

mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 80
CHLORIDE 39
PHOSPHATE 30 (15 mmol/L)

BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
AND HYDROCHLORIC ACID

STERILE SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)
IN UNOPENED OVERWRAP
AVOID EXCESSIVE HEAT
PROTECT FROM FREEZING
SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

EXP LOT

1600

1000

800

600

400

BE-35-05-072

200

2B7721L 2000 mL
NDC 0338-7032-01

(01) 00303387032016

CLINIMIX E
WITH ELECTROLYTES 5/15

SULFITE-FREE

(5% Amino Acids with Electrolytes in 15% Dextrose with Calcium) Injection

CENTRAL LINE INFUSION ONLY

1000 mL INJECTION PORT CHAMBER
30% Dextrose Injection with Calcium

1000 mL OUTLET PORT CHAMBER
10% Amino Acid Injection with Electrolytes

ACTIVATE SEAL AND MIX THOROUGHLY BEFORE USE

SEE PRESCRIBING INFORMATION FOR INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO 9 DAYS ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT ALTER THE QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

Rx Only

2D Datamatrix (GS1):
NDC, EXP, LOT

2D Datamatrix Barcode

EXP LOT

CONTENTS OF EACH 100 mL OF THE ADMIXED INJECTION

DEXTROSE HYDROUS USP	15 g
ESSENTIAL AMINO ACIDS	
LEUCINE	365 mg
ISOLEUCINE	300 mg
VALINE	290 mg
LYSINE (ADDED AS THE HYDROCHLORIDE SALT)	290 mg
PHENYLALANINE	280 mg
HISTIDINE	240 mg
THREONINE	210 mg
METHIONINE	200 mg
TRYPTOPHAN	90 mg
NONESSENTIAL AMINO ACIDS	
ALANINE	1035 mg
ARGININE	575 mg
GLYCINE	515 mg
PROLINE	340 mg
SERINE	250 mg
TYROSINE	20 mg
ELECTROLYTES	
SODIUM ACETATE TRIHYDRATE USP	340 mg
DIBASIC POTASSIUM PHOSPHATE USP	261 mg
SODIUM CHLORIDE USP	59 mg
MAGNESIUM CHLORIDE USP	51 mg
CALCIUM CHLORIDE DIHYDRATE USP	33 mg
mEq/L	
SODIUM	35
POTASSIUM	30
MAGNESIUM	5
CALCIUM	4.5 (2.2 mmol/L)
ACETATE	80
CHLORIDE	39
PHOSPHATE	30 (15 mmol/L)

BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID AND HYDROCHLORIC ACID

STERILE
SINGLE DOSE CONTAINER

STORE AT ROOM TEMPERATURE (25°C/77°F) IN UNOPENED OVERWRAP

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

Baxter
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

BE-35-05-066

2000 1600 1000 800 600 400 200

Container Label

2B7722L
NDC 0338-7036-01
2000 mL

Barcode
(01) 00303387032016

CLINIMIX E
WITH ELECTROLYTES 5/20

SULFITE-FREE

(5% Amino Acids with Electrolytes in 20% Dextrose with Calcium) Injection

CENTRAL LINE INFUSION ONLY

1000 mL INJECTION PORT CHAMBER
40% Dextrose Injection with Calcium

1000 mL OUTLET PORT CHAMBER
10% Amino Acid Injection

Rx Only

ACTIVATE SEAL AND
MIX THOROUGHLY BEFORE USE

SEE PRESRCIBING INFORMATION FOR INSTRUCTIONS ON ACTIVATION

AFTER MIXING THE PRODUCT REPRESENTS 2000 mL

REFRIGERATED STORAGE IS LIMITED TO
9 DAYS ONCE OVERWRAP IS OPENED

A SLIGHT YELLOW COLOR DOES NOT ALTER THE
QUALITY AND EFFICACY OF THIS PRODUCT

ASK PHARMACIST ABOUT ADDITIVE COMPATIBILITY

2D Datamatrix (GS1):
NDC, LOT, EXP

2D Datamatrix
Barcode

CONTENTS OF EACH 100 mL OF THE ADMIXED
INJECTION
DEXTROSE HYDROUS USP 20 g

ESSENTIAL AMINO ACIDS
LEUCINE 365 mg
ISOLEUCINE 300 mg
VALINE 290 mg
LYSINE (ADDED AS THE
HYDROCHLORIDE SALT) 290 mg
PHENYLALANINE 280 mg
HISTIDINE 240 mg
THREONINE 210 mg
METHIONINE 200 mg
TRYPTOPHAN 90 mg

NONESSENTIAL AMINO ACIDS
ALANINE 1035 mg
ARGININE 575 mg
GLYCINE 515 mg
PROLINE 340 mg
SERINE 250 mg
TYROSINE 20 mg

ELECTROLYTES
SODIUM ACETATE
TRIHYDRATE USP 340 mg
DIBASIC POTASSIUM
PHOSPHATE USP 261 mg
SODIUM CHLORIDE USP 59 mg
MAGNESIUM CHLORIDE USP 51 mg
CALCIUM CHLORIDE
DIHYDRATE USP 33 mg

mEq/L
SODIUM 35
POTASSIUM 30
MAGNESIUM 5
CALCIUM 4.5 (2.2 mmol/L)
ACETATE 80
CHLORIDE 39
PHOSPHATE 30 (15 mmol/L)

BALANCED BY IONS FROM AMINO ACIDS
pH ADJUSTED WITH GLACIAL ACETIC ACID
AND HYDROCHLORIC ACID

STERILE
SINGLE DOSE CONTAINER

ROOM TEMPERATURE (25°C/77°F)
IN UNOPENED OVERWARP

AVOID EXCESSIVE HEAT

PROTECT FROM FREEZING

SEE PRESCRIBING INFORMATION

BAXTER LOGO
BAXTER HEALTHCARE CORPORATION
DEERFIELD IL 60015 USA
MADE IN BELGIUM

EXP LOT

2000

1600

1000

800

600

400

BE-35-05-066

200

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-1142
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	201 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	154 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	159 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	110 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	165 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	160 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	132 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	116 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	50 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	570 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	283 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	316 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	187 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	138 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	11 mg in 100 mL
SODIUM ACETATE (UNII: 4550K0SC9B) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM ACETATE	217 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	112 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	5 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1142-03	1000 mL in 1 BAG; Type 0: Not a Combination Product	03/26/1997	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	03/26/1997	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-1144
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	311 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	238 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	247 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	170 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	255 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	247 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	204 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	179 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	77 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	880 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	438 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	489 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	289 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	213 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	17 mg in 100 mL
SODIUM ACETATE (UNII: 4550K0SC9B) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM ACETATE	297 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	77 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	5 g in 100 mL

Inactive Ingredients	
Ingredient Name	**Strength**
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	

Packaging				
#	**Item Code**	**Package Description**	**Marketing Start Date**	**Marketing End Date**
1	NDC:0338-1144-03	1000 mL in 1 BAG; Type 0: Not a Combination Product	03/26/1997	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	03/26/1997	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-1113
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	311 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	238 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	247 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	170 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	255 mg in 100 mL

VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	247 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	204 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	179 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	77 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	880 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	438 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	489 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	289 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	213 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	17 mg in 100 mL
SODIUM ACETATE (UNII: 4550K0SC9B) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM ACETATE	297 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	77 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	5 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1113-04	2000 mL in 1 BAG; Type 0: Not a Combination Product	03/26/1997	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	03/26/1997	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-1145
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	311 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	238 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	247 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	170 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	255 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	247 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	204 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	179 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	77 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	880 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	438 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	489 mg in 100 mL

PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	289 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	213 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	17 mg in 100 mL
SODIUM ACETATE (UNII: 4550K0SC9B) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM ACETATE	297 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	77 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	10 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1145-03	1000 mL in 1 BAG; Type 0: Not a Combination Product	03/26/1997	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	03/26/1997	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-1115
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	311 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	238 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	247 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	170 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	255 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	247 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	204 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	179 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	77 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	880 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	438 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	489 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	289 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	213 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	17 mg in 100 mL
SODIUM ACETATE (UNII: 4550K0SC9B) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM ACETATE	297 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37,	SODIUM CHLORIDE	77 mg

CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	10 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1115-04	2000 mL in 1 BAG; Type 0: Not a Combination Product	03/26/1997	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	03/26/1997	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-1147
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	365 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	280 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	290 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	200 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	300 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	290 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	240 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	210 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	90 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	1035 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	515 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	575 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	340 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	250 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	20 mg in 100 mL
SODIUM ACETATE (UNII: 4550K0SC9B) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM ACETATE	340 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	59 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	15 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1147-03	1000 mL in 1 BAG; Type 0: Not a Combination Product	03/26/1997	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	03/26/1997	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-1123
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	365 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	280 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	290 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	200 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	300 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	290 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	240 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	210 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	90 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	1035 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	515 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	575 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	340 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	250 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	20 mg in 100 mL
SODIUM ACETATE (UNII: 4550K0SC9B) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM ACETATE	340 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	59 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	15 g in 100 mL

Inactive Ingredients	
Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1123-04	2000 mL in 1 BAG; Type 0: Not a Combination Product	03/26/1997	

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End

Category	Citation	Date	Date
NDA	NDA020678	03/26/1997	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-1125
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	365 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	280 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	290 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	200 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	300 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	290 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	240 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	210 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	90 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	1035 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	515 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	575 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	340 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	250 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	20 mg in 100 mL
SODIUM ACETATE (UNII: 4550K0SC9B) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM ACETATE	340 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	59 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	20 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1125-04	2000 mL in 1 BAG; Type 0: Not a Combination Product	03/26/1997	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	03/26/1997	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-1148
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	365 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	280 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	290 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	200 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	300 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	290 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	240 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	210 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	90 mg in 100 mL
Alanine (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	Alanine	1035 mg in 100 mL
Glycine (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	Glycine	515 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	575 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	340 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	250 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	20 mg in 100 mL
SODIUM ACETATE (UNII: 4550K0SC9B) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM ACETATE	340 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	59 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	20 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-1148-03	1000 mL in 1 BAG; Type 0: Not a Combination Product	03/26/1997	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	03/26/1997	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-0202
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	584 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	448 mg in 100 mL
		464 mg

LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	464 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	320 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	480 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	464 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	384 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	336 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	144 mg in 100 mL
Alanine (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	Alanine	1656 mg in 100 mL
Glycine (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	Glycine	824 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	920 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	544 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	400 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	32 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	205 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	14 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0202-06	6 in 1 CARTON	09/21/2020	
1	NDC:0338-0202-01	1000 mL in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	09/21/2020	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-0206
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	584 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	448 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	464 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	320 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	480 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	464 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	384 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	336 mg in 100 mL

TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	144 mg in 100 mL
Alanine (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	Alanine	1656 mg in 100 mL
Glycine (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	Glycine	824 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	920 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	544 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	400 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	32 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	205 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	14 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0206-04	4 in 1 CARTON	09/21/2020	
1	NDC:0338-0206-01	2000 mL in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	09/21/2020	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-0210
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	584 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	448 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	464 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	320 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	480 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	464 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	384 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	336 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	144 mg in 100 mL
Alanine (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	Alanine	1656 mg in 100 mL
Glycine (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	Glycine	824 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	920 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	544 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	400 mg in 100 mL

TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	32 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	205 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	10 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0210-06	6 in 1 CARTON	09/21/2020	
1	NDC:0338-0210-01	1000 mL in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	09/21/2020	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-0214
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	584 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	448 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	464 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	320 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	480 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	464 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	384 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	336 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	144 mg in 100 mL
Alanine (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	Alanine	1656 mg in 100 mL
Glycine (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	Glycine	824 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	920 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	544 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	400 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	32 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	205 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	10 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-0214-04	4 in 1 CARTON	09/21/2020	
1	NDC:0338-0214-01	2000 mL in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	09/21/2020	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-7020
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	201 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	154 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	159 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	110 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	165 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	160 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	132 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	116 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	50 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	570 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	283 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	316 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	187 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	138 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	11 mg in 100 mL
SODIUM ACETATE (UNII: 4550K0SC9B) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM ACETATE	216 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	112 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	5 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-7020-06	6 in 1 CARTON	07/15/2024	
1	NDC:0338-7020-01	1000 mL in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	07/15/2024	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-7022
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	311 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	238 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	247 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	170 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	255 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	247 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	204 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	179 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	77 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	880 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	438 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	489 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	289 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	213 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	17 mg in 100 mL
SODIUM ACETATE (UNII: 4550K0SC9B) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM ACETATE	297 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	77 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	5 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-7022-06	6 in 1 CARTON	07/15/2024	
1	NDC:0338-7022-01	1000 mL in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	07/15/2024	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-7024
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	311 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	238 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	247 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	170 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	255 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	247 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	204 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	179 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	77 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	880 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	438 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	489 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	289 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	213 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	17 mg in 100 mL
SODIUM ACETATE (UNII: 4550K0SC9B) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM ACETATE	297 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	77 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	5 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-7024-04	4 in 1 CARTON	07/15/2024	
1	NDC:0338-7024-01	2000 mL in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	07/15/2024	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-7026
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	311 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	238 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	247 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	170 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	255 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	247 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	204 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	179 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	77 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	880 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	438 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	489 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	289 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	213 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	17 mg in 100 mL
SODIUM ACETATE (UNII: 4550K0SC9B) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM ACETATE	297 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	77 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	10 g in 100 mL

Inactive Ingredients	
Ingredient Name	**Strength**
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging				
#	**Item Code**	**Package Description**	**Marketing Start Date**	**Marketing End Date**
1	NDC:0338-7026-06	6 in 1 CARTON	07/15/2024	
1	NDC:0338-7026-01	1000 mL in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	07/15/2024	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-7028
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	311 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	238 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	247 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	170 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	255 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	247 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	204 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	179 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	77 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	880 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	438 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	489 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	289 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	213 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	17 mg in 100 mL
SODIUM ACETATE (UNII: 4550K0SC9B) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM ACETATE	297 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	77 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	10 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-7028-04	4 in 1 CARTON	07/15/2024	
1	NDC:0338-7028-01	2000 mL in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	07/15/2024	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-7030
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	365 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	280 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	290 mg

LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	200 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	300 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	290 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	240 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	210 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	90 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	1035 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	515 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	575 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	340 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	250 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	20 mg in 100 mL
SODIUM ACETATE (UNII: 4550K0SC9B) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM ACETATE	340 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	59 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	15 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-7030-06	6 in 1 CARTON	07/15/2024	
1	NDC:0338-7030-01	1000 mL in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	07/15/2024	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-7032
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	365 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	280 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	290 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	200 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	300 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	290 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	240 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	210 mg

THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	90 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	1035 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	515 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	575 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	340 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	250 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	20 mg in 100 mL
SODIUM ACETATE (UNII: 4550K0SC9B) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM ACETATE	340 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	59 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	15 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-7032-04	4 in 1 CARTON	07/15/2024	
1	NDC:0338-7032-01	2000 mL in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	07/15/2024	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-7034
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	365 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	280 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	290 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	200 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	300 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	290 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	240 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	210 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	90 mg in 100 mL
Alanine (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	Alanine	1035 mg in 100 mL
Glycine (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	Glycine	515 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	575 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	340 mg

PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	250 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	20 mg in 100 mL
SODIUM ACETATE (UNII: 4550K0SC9B) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM ACETATE	340 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL
MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	59 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	20 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-7034-06	6 in 1 CARTON	07/15/2024	
1	NDC:0338-7034-01	1000 mL in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	07/15/2024	

CLINIMIX E

leucine, phenylalanine, lysine, methionine, isoleucine, valine, histidine, threonine, tryptophan, alanine, glycine, arginine, proline, serine, tyrosine, sodium acetate, dibasic potassium phosphate, magnesium chloride, sodium chloride, calcium chloride, dextrose injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	**Item Code (Source)**	NDC:0338-7036
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LEUCINE (UNII: GMW67QNF9C) (LEUCINE - UNII:GMW67QNF9C)	LEUCINE	365 mg in 100 mL
PHENYLALANINE (UNII: 47E5O17Y3R) (PHENYLALANINE - UNII:47E5O17Y3R)	PHENYLALANINE	280 mg in 100 mL
LYSINE (UNII: K3Z4F929H6) (LYSINE - UNII:K3Z4F929H6)	LYSINE	290 mg in 100 mL
METHIONINE (UNII: AE28F7PNPL) (METHIONINE - UNII:AE28F7PNPL)	METHIONINE	200 mg in 100 mL
ISOLEUCINE (UNII: 04Y7590D77) (ISOLEUCINE - UNII:04Y7590D77)	ISOLEUCINE	300 mg in 100 mL
VALINE (UNII: HG18B9YRS7) (VALINE - UNII:HG18B9YRS7)	VALINE	290 mg in 100 mL
HISTIDINE (UNII: 4QD397987E) (HISTIDINE - UNII:4QD397987E)	HISTIDINE	240 mg in 100 mL
THREONINE (UNII: 2ZD004190S) (THREONINE - UNII:2ZD004190S)	THREONINE	210 mg in 100 mL
TRYPTOPHAN (UNII: 8DUH1N11BX) (TRYPTOPHAN - UNII:8DUH1N11BX)	TRYPTOPHAN	90 mg in 100 mL
ALANINE (UNII: OF5P57N2ZX) (ALANINE - UNII:OF5P57N2ZX)	ALANINE	1035 mg in 100 mL
GLYCINE (UNII: TE7660XO1C) (GLYCINE - UNII:TE7660XO1C)	GLYCINE	515 mg in 100 mL
ARGININE (UNII: 94ZLA3W45F) (ARGININE - UNII:94ZLA3W45F)	ARGININE	575 mg in 100 mL
PROLINE (UNII: 9DLQ4CIU6V) (PROLINE - UNII:9DLQ4CIU6V)	PROLINE	340 mg in 100 mL
SERINE (UNII: 452VLY9402) (SERINE - UNII:452VLY9402)	SERINE	250 mg in 100 mL
TYROSINE (UNII: 42HK56048U) (TYROSINE - UNII:42HK56048U)	TYROSINE	20 mg in 100 mL
SODIUM ACETATE (UNII: 4550K0SC9B) (ACETATE ION - UNII:569DQM74SC, SODIUM CATION - UNII:LYR4M0NH37)	SODIUM ACETATE	340 mg in 100 mL
POTASSIUM PHOSPHATE, DIBASIC (UNII: CI71S98N1Z) (PHOSPHATE ION - UNII:NK08V8K8HR, POTASSIUM CATION - UNII:295O53K152)	POTASSIUM PHOSPHATE, DIBASIC	261 mg in 100 mL

MAGNESIUM CHLORIDE (UNII: 02F3473H9O) (MAGNESIUM CATION - UNII:T6V3LHY838, CHLORIDE ION - UNII:Q32ZN48698)	MAGNESIUM CHLORIDE	51 mg in 100 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X) (SODIUM CATION - UNII:LYR4M0NH37, CHLORIDE ION - UNII:Q32ZN48698)	SODIUM CHLORIDE	59 mg in 100 mL
CALCIUM CHLORIDE (UNII: M4I0D6VV5M) (CALCIUM CATION - UNII:2M83C4R6ZB, CHLORIDE ION - UNII:Q32ZN48698)	CALCIUM CHLORIDE	33 mg in 100 mL
DEXTROSE (UNII: IY9XDZ35W2) (ANHYDROUS DEXTROSE - UNII:5SL0G7R0OK)	DEXTROSE	20 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
WATER (UNII: 059QF0KO0R)	
NITROGEN (UNII: N762921K75)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0338-7036-04	4 in 1 CARTON	07/15/2024	
1	NDC:0338-7036-01	2000 mL in 1 BAG; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020678	07/15/2024	

Labeler - Baxter Healthcare Company (005083209)

Establishment

Name	Address	ID/FEI	Business Operations
Baxter Healthcare Corporation		189326168	ANALYSIS(0338-1142, 0338-1144, 0338-1113, 0338-1145, 0338-1115, 0338-1147, 0338-1123, 0338-1125, 0338-1148) , LABEL(0338-1142, 0338-1144, 0338-1113, 0338-1145, 0338-1115, 0338-1147, 0338-1123, 0338-1125, 0338-1148) , MANUFACTURE(0338-1142, 0338-1144, 0338-1113, 0338-1145, 0338-1115, 0338-1147, 0338-1123, 0338-1125, 0338-1148) , PACK(0338-1142, 0338-1144, 0338-1113, 0338-1145, 0338-1115, 0338-1147, 0338-1123, 0338-1125, 0338-1148) , STERILIZE(0338-1142, 0338-1144, 0338-1113, 0338-1145, 0338-1115, 0338-1147, 0338-1123, 0338-1125, 0338-1148)

Establishment

Name	Address	ID/FEI	Business Operations
Baxter Healthcare Corporation		194684502	ANALYSIS(0338-1142, 0338-1144, 0338-1113, 0338-1145, 0338-1115, 0338-1147, 0338-1123, 0338-1125, 0338-1148)

Establishment

Name	Address	ID/FEI	Business Operations
Baxter Healthcare Corporation		059140764	ANALYSIS(0338-1142, 0338-1144, 0338-1113, 0338-1145, 0338-1115, 0338-1147, 0338-1123, 0338-1125, 0338-1148) , MANUFACTURE(0338-1142, 0338-1144, 0338-1113, 0338-1145, 0338-1115, 0338-1147, 0338-1123, 0338-1125, 0338-1148) , LABEL(0338-1142, 0338-1144, 0338-1113, 0338-1145, 0338-1115, 0338-1147, 0338-1123, 0338-1125, 0338-1148) , PACK(0338-1142, 0338-1144, 0338-1113, 0338-1145, 0338-1115, 0338-1147, 0338-1123, 0338-1125, 0338-1148) , STERILIZE(0338-1142, 0338-1144, 0338-1113, 0338-1145, 0338-1115, 0338-1147, 0338-1123, 0338-1125, 0338-1148)

Establishment

Name	Address	ID/FEI	Business Operations
Baxter SA		370353835	ANALYSIS(0338-0202, 0338-0206, 0338-0210, 0338-0214, 0338-7020, 0338-7022, 0338-7024, 0338-7026, 0338-7028, 0338-7030, 0338-7032, 0338-7034, 0338-7036) , MANUFACTURE(0338-0202, 0338-0206, 0338-0210, 0338-0214, 0338-7020, 0338-7022, 0338-7024, 0338-7026, 0338-7028, 0338-7030, 0338-7032, 0338-7034, 0338-7036) , PACK(0338-0202, 0338-0206, 0338-0210, 0338-0214, 0338-7020, 0338-7022, 0338-7024, 0338-7026, 0338-7028, 0338-7030, 0338-7032, 0338-7034, 0338-7036) , STERILIZE(0338-0202, 0338-0206, 0338-0210, 0338-0214, 0338-7020, 0338-7022, 0338-7024, 0338-7026, 0338-7028, 0338-7030, 0338-7032, 0338-7034, 0338-7036) , LABEL(0338-0202, 0338-0206, 0338-0210, 0338-0214, 0338-7020, 0338-7022, 0338-7024, 0338-7026, 0338-7028, 0338-7030, 0338-7032, 0338-7034, 0338-7036)