BENZOICUM ACIDUM- benzoic acid pellet Boiron

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

Benzoicum acidum 30C

Benzoicum acidum 30C

Active ingredient**: See product name on front panel (**contains 0.443 mg of the active ingredient per pellet).

Relieves minor arthritis pain in the wrists *

Uses: See symptoms on front panel.

Stop use and ask a doctor if symptoms persist for more than 3 days or worsen.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Do not use if pellet dispenser seal is broken.

Contains approx 80 pellets.

How to dispense pellets? Turn tube upside down. Twist until 5 pellets are dispensed into cap. Carefully remove the cap and use it to pour pellets under the tongue.

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

*C,K,CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

lactose, sucrose

Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

1-800-BOIRON-1 (1-800-264-7661), BoironUSA.com Info@boiron.com Distributed by Boiron, Inc. Newtown Square, PA 19073



Drug Facts

Active ingredient**: See product name on front panel (contains 0.443 mg of the active ingredient per pellet).

Uses: See symptoms on front panel.

Warnings: Stop use and ask a doctor if symptoms persist for more than 3 days or worsen. If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Directions: ■ Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.





Lot:

Exp:

06960

11113

Contains approx. 80 pellets. US

Peel for Drugs Facts and instructions for use.



Active ingredient**: See product name on front panel (contains 0.443 mg of the active ingredient per pellet).

Uses: See symptoms on front panel.

Warnings: Stop use and ask a doctor if symptoms persist for more than 3 days or worsen. If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children.

Directions: • Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

Other information: • Do not use if pellet dispenser seal is broken.

Drug Facts (continued) **Inactive ingredients:** lactose, sucrose



BENZOICUM ACIDUM

benzoic acid pellet

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0220-0824	
Route of Administration	ORAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
BENZOIC ACID (UNII: 85KN0B0MIM) (BENZOIC ACID - UNII:85KN0B0MIM)	BENZOIC ACID	30 [hp_C] in 30 [hp_C]	

Inactive Ingredients			
Ingredient Name	Strength		
SUCROSE (UNII: C151H8M554)			
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)			

Product Characteristics			
Color	white	Score	

Shape	ROUND	Size	4mm
Flavor		Imprint Code	
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0220-0824- 41	30 [hp_C] in 1 TUBE; Type 0: Not a Combination Product	03/03/1983	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
unapproved homeopathic		03/03/1983		

Labeler - Boiron (282560473)

Registrant - Boiron, Inc. (014892269)

Establishment				
Name	Address	ID/FEI	Business Operations	
Boiron		282560473	manufacture(0220-0824)	

Revised: 2/2024 Boiron