

**SEVERE SINUS RELIEF MIST- oxymetazoline spray liquid**  
**Walgreens Company**

-----  
**Severe Nasal Spray**

**Drug Facts**

**Active Ingredient**

Oxymetazoline Hydrochloride 0.05%

**Purpose**

Nasal Decongestant

**Uses**

Temporarily relieves nasal congestion due to:

- common cold
- hay fever
- upper respiratory allergies
- temporarily relieves sinus congestion and pressure
- Shrinks swollen membranes so you can breathe more freely

**Warnings**

**Ask a Doctor before use if you have**

- heart disease
- high blood pressure
- diabetes
- thyroid disease
- trouble urinating due to enlarged prostate gland.

**When using this product**

- **Do not use more than directed**
- do not use for more than three days, Use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.
- temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur.
- use of this container by more than one person may spread infection.

**stop use and ask a doctor if**

symptoms persist.

**if pregnant or breast-feeding**

ask a health professional before use.

**Keep out of the reach of children**

if swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

- Before using for the first time, remove the protective cap from the tip and prime the pump by depressing pump firmly several times. To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress rim with a firm and even stroke, and sniff deeply.
- Wipe nozzle clean after use.
- Adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.
- Children under 6 years of age consult a doctor.

**Other information**

- store between 20° and 25°C (68° to 77°F)
- retain carton for future reference on full labeling

**Questions or Comments ?**

**1-800-222-1087**

**Inactive Ingredients**

Benzalkonium Chloride, Benzyl Alcohol, Citric Acid Anhydrous, Edetate Sodium, Fragrance, Polysorbate 80, Purified Water, Sodium Citrate, Sorbitol

**Principal Panel 0.5 FLOZ Bottle**

Walgreens

NDC 0363-2305-52

**Severe Sinus Relief Mist**  
OXYMETAZOLINE HCl 0.05% /  
NASAL DECONGESTANT

• 12 hour

0.5 FL OZ (15 mL)

SEE CARTON FOR FULL LABELING

DO NOT USE IF PRINTED SEAL OVER  
CAP IS TORN OR MISSING

**Active ingredient** Oxymetazoline hydrochloride 0.05% **Uses** • See Carton.

**Warnings Ask a doctor before use if you have**  
• heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to an enlarged prostate gland. **When using this product**  
• do not use more than directed • do not use for more than 3 days. Frequent or prolonged use may cause nasal congestion to recur or worsen. • temporary nasal congestion to recur or worsen. • increase in nasal discharge may occur • use of this container by more than one person may spread infection. **Stop use and ask a doctor if symptoms persist. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions** To use: squeeze cap at base on opposite sides and twist cap counterclockwise to remove. Secure cap after use. • adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period. • children under 6 years of age: ask a doctor.

**Questions or comments?**

1-800-222-1087

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com ©2018 Walgreen Co.  
ITEM 821792 W000000-0000-0

REV0319  
ORG0518-F

Walgreens Severe Sinus Relief Mist  
Nasal Spray  
Oxymetazoline HCL 0.05%  
12 hour Nasal Spray  
Net Wt 0.5 FLOZ (15ml)

Walgreens

NDC 0363-2305-52

**Severe Sinus Relief Mist**  
OXYMETAZOLINE HCl 0.05% /  
NASAL DECONGESTANT

• 12 hour

0.5 FL OZ (15 mL)

SEE CARTON FOR FULL LABELING

DO NOT USE IF PRINTED SEAL OVER  
CAP IS TORN OR MISSING

**Active ingredient** Oxymetazoline hydrochloride 0.05% **Uses** • See Carton.

**Warnings Ask a doctor before use if you have**  
• heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to an enlarged prostate gland. **When using this product**  
• do not use more than directed • do not use for more than 3 days. Frequent or prolonged use may cause nasal congestion to recur or worsen. • temporary nasal congestion to recur or worsen. • increase in nasal discharge may occur • use of this container by more than one person may spread infection. **Stop use and ask a doctor if symptoms persist. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions** To use: squeeze cap at base on opposite sides and twist cap counterclockwise to remove. Secure cap after use. • adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period. • children under 6 years of age: ask a doctor.

**Questions or comments?**

1-800-222-1087

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com ©2018 Walgreen Co.  
ITEM 821792 W000000-0000-0

REV0319  
ORG0518-F

Walgreens

NDC 0363-2305-52

**Severe Sinus Relief Mist**  
OXYMETAZOLINE HCl 0.05% /  
NASAL DECONGESTANT

• 12 hour

0.5 FL OZ (15 mL)

SEE CARTON FOR FULL LABELING

DO NOT USE IF PRINTED SEAL OVER  
CAP IS TORN OR MISSING

**Active ingredient** Oxymetazoline hydrochloride 0.05% **Uses** • See Carton.

**Warnings Ask a doctor before use if you have**  
• heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to an enlarged prostate gland. **When using this product**  
• do not use more than directed • do not use for more than 3 days. Frequent or prolonged use may cause nasal congestion to recur or worsen. • temporary nasal congestion to recur or worsen. • increase in nasal discharge may occur • use of this container by more than one person may spread infection. **Stop use and ask a doctor if symptoms persist. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions** To use: squeeze cap at base on opposite sides and twist cap counterclockwise to remove. Secure cap after use. • adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period. • children under 6 years of age: ask a doctor.

**Questions or comments?**

1-800-222-1087

DISTRIBUTED BY: WALGREEN CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com ©2018 Walgreen Co.  
ITEM 821792 W000000-0000-0

REV0319  
ORG0518-F

*Walgreens*

NDC 0363-2305-52  
**Severe Sinus Relief Mist**  
 OXYMETAZOLINE HCl 0.05% /  
 NASAL DECONGESTANT

• 12 hour  
 0.5 FL OZ (15 mL)

**SEE CARTON FOR FULL LABELING**

**DO NOT USE IF PRINTED SEAL OVER CAP IS TORN OR MISSING**

**Active ingredient:** Oxymetazoline hydrochloride 0.05% **Uses:** See Carton.

**Warnings: Ask a doctor before use if you have**

- heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to an enlarged prostate gland. **When using this product**
- do not use more than directed • do not use for more than 3 days. Frequent or prolonged use may cause nasal congestion to recur or worsen. • temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur • use of this container by more than one person may spread infection. **Stop use and ask a doctor if symptoms persist. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions:** To use: squeeze cap at base on opposite sides and twist cap counterclockwise to remove. Secure cap after use. • adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period. • children under 6 years of age: ask a doctor.

**Questions or comments?**  
**1-800-222-1087**  
 DISTRIBUTED BY: WALGREEN CO.  
 200 WILMOT RD., DEERFIELD, IL 60015  
**100% SATISFACTION GUARANTEED**  
 walgreens.com ©2018 Walgreen Co.  
 ITEM 821792 W00000-0000-0

**REV0319**  
**ORG0518-F**

*Walgreens*

NDC 0363-2305-52  
**Severe Sinus Relief Mist**  
 OXYMETAZOLINE HCl 0.05% /  
 NASAL DECONGESTANT

• 12 hour  
 0.5 FL OZ (15 mL)

**SEE CARTON FOR FULL LABELING**

**DO NOT USE IF PRINTED SEAL OVER CAP IS TORN OR MISSING**

**Active ingredient:** Oxymetazoline hydrochloride 0.05% **Uses:** See Carton.

**Warnings: Ask a doctor before use if you have**

- heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to an enlarged prostate gland. **When using this product**
- do not use more than directed • do not use for more than 3 days. Frequent or prolonged use may cause nasal congestion to recur or worsen. • temporary discomfort such as burning, stinging, sneezing or an increase in nasal discharge may occur • use of this container by more than one person may spread infection. **Stop use and ask a doctor if symptoms persist. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions:** To use: squeeze cap at base on opposite sides and twist cap counterclockwise to remove. Secure cap after use. • adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period. • children under 6 years of age: ask a doctor.

**Questions or comments?**  
**1-800-222-1087**  
 DISTRIBUTED BY: WALGREEN CO.  
 200 WILMOT RD., DEERFIELD, IL 60015  
**100% SATISFACTION GUARANTEED**  
 walgreens.com ©2018 Walgreen Co.  
 ITEM 821792 W00000-0000-0

**REV0319**  
**ORG0518-F**

## Principal display panel -Carton

Walgreens Severe Sinus Relief Mist

Nasal Spray

Oxymetazoline HCL 0.05%

12 hour Nasal Spray

Net Wt 0.5 FLOZ (15ml)

INTEGRATING 0.05% / NASAL DECONGESTANT

# Severe Sinus Relief Mist

Walgreens

DO NOT USE IF PRINTED SEAL OVER CAP IS TORN OR MISSING

### Drug Facts

**Active ingredient** Oxymetazoline hydrochloride 0.05%, Nasal decongestant

**Purpose** temporarily relieves nasal congestion due to:

• common cold • hay fever • upper respiratory allergies • temporarily relieves sinus congestion and pressure • shrinks swollen nasal membranes so you can breathe more freely

### Warnings

**Ask a doctor before use if you have** • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to an enlarged prostate gland

**When using this product**

• do not use more than directed  
• do not use for more than 3 days (use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.)

• temporary discomfort such as burning, stinging, itching or an increase in nasal discharge may occur  
• use of this container by more than one person may spread infection

**Stop use and ask a doctor if symptoms persist**

**if pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.**

### Directions

Before using the first time, remove the protective cap from the tip and prime pump by depressing pump firmly several times. To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress rim with a firm, even stroke and sniff deeply. Wipe nozzle clean after use.

• adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.  
• children under 6 years of age: ask a doctor.

### Drug Facts (continued)

**Other information**

• store between 20° to 25°C (68° to 77°F)

• refer to carton for future reference on full labeling

### Inactive ingredients

Benzalkonium Chloride, Benzyl Alcohol, Citric Acid Anhydrous, Disodium Phosphate, Fragrance, Polyethylene Glycol, Purified Water, Sodium Citrate, Sorbitol

### IMPORTANT:

**KEEP OUTER CARTON FOR FUTURE REFERENCE ON FULL LABELING**

### Cap Removal Instructions:

To remove cap, squeeze cap at base on opposite sides on the two firm impressions with fingers. While squeezing, twist cap counterclockwise to remove. Press down to lock. Refer to arrow on cap. To reapply cap, turn cap on pump clockwise until it locks.

### Walgreens Pharmacy Recommendation

Walgreens Pharmacy Survey

This product is not manufactured or distributed by Procter & Gamble, the distributor of Vicks® Sinex™ Severe.

### Questions or comments?

1-800-222-1037

DISTRIBUTED BY: WALGREENS CO.

200 WILMOT RD. DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED

walgreens.com ©2010 Walgreen Co.

Walgreens

Compare to Vicks® Sinex™ Severe active ingredient!

NDC 0583-2205-02

# Severe Sinus Relief Mist

OXYMETAZOLINE HCl 0.05% / NASAL DECONGESTANT

SEVERE

12 HOUR

ULTRA-FINE SPRAY

• Helps relieve congestion & sinus pressure

0.5 FL OZ (15 mL)

NET WT

LOT

ORG0518-F

FSP



INTEGRATING 0.05% / NASAL DECONGESTANT

# Severe Sinus Relief Mist

Walgreens

DO NOT USE IF PRINTED SEAL OVER CAP IS TORN OR MISSING

### Drug Facts

**Active ingredient** Oxymetazoline hydrochloride 0.05%, Nasal decongestant

**Purpose** temporarily relieves nasal congestion due to:

• common cold • hay fever • upper respiratory allergies • temporarily relieves sinus congestion and pressure • shrinks swollen nasal membranes so you can breathe more freely

### Warnings

**Ask a doctor before use if you have** • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to an enlarged prostate gland

**When using this product**

• do not use more than directed  
• do not use for more than 3 days (use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.)

• temporary discomfort such as burning, stinging, itching or an increase in nasal discharge may occur  
• use of this container by more than one person may spread infection

**Stop use and ask a doctor if symptoms persist**

**if pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.**

### Directions

Before using the first time, remove the protective cap from the tip and prime pump by depressing pump firmly several times. To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress rim with a firm, even stroke and sniff deeply. Wipe nozzle clean after use.

• adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.  
• children under 6 years of age: ask a doctor.

### Drug Facts (continued)

**Other information**

• store between 20° to 25°C (68° to 77°F)  
• refer to carton for future reference on full labeling

### Inactive ingredients

Benzalkonium Chloride, Benzyl Alcohol, Citric Acid Anhydrous, Disodium Phosphate, Fragrance, Polyethylene Glycol, Purified Water, Sodium Citrate, Sorbitol

**IMPORTANT: KEEP OUTER CARTON FOR FUTURE REFERENCE ON FULL LABELING**

### Cap Removal Instructions

To remove cap, squeeze cap at base on opposite sides on the two firm impressions with fingers. While squeezing, twist cap counterclockwise to remove. Press down to lock. Repeat to remove on cap. To reapply cap, turn cap on pump clockwise until it locks.

Walgreens Pharmacy Recommended  
Walgreens Pharmacy Survey

This product is not manufactured or distributed by Procter & Gamble, the distributor of Vicks® Sinex™ Severe.

### Questions or comments?

1-800-222-1037

DISTRIBUTED BY: WALGREENS CO.  
200 WILMOT RD., DEERFIELD, IL 60015  
100% SATISFACTION GUARANTEED  
walgreens.com ©2010 Walgreen Co.

Walgreens

Compare to Vicks® Sinex™ Severe active ingredient!

NDC 0583-2205-02

# Severe Sinus Relief Mist

OXYMETAZOLINE HCl 0.05% / NASAL DECONGESTANT

SEVERE

12 HOUR

ULTRA-FINE SPRAY

• Helps relieve congestion & sinus pressure

0.5 FL OZ (15 mL)

NOT FOR SALE

LOT

ORG0518-F

FSP



INTEGRATING 0.05% / NASAL DECONGESTANT

# Severe Sinus Relief Mist

Walgreens

DO NOT USE IF PRINTED SEAL OVER CAP IS TORN OR MISSING

### Drug Facts

**Active ingredient** Oxymetazoline hydrochloride 0.05%, Nasal decongestant

**Purpose** temporarily relieves nasal congestion due to:

• common cold • hay fever • upper respiratory allergies • temporarily relieves sinus congestion and pressure • shrinks swollen nasal membranes so you can breathe more freely

### Warnings

**Ask a doctor before use if you have** • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to an enlarged prostate gland

**When using this product**

• do not use more than directed  
• do not use for more than 3 days (use only as directed. Frequent or prolonged use may cause nasal congestion to recur or worsen.)

• temporary discomfort such as burning, stinging, itching or an increase in nasal discharge may occur  
• use of this container by more than one person may spread infection

**Stop use and ask a doctor if symptoms persist**

**if pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.**

### Directions

Before using the first time, remove the protective cap from the tip and prime pump by depressing pump firmly several times. To spray, hold bottle with thumb at base and nozzle between first and second fingers. Without tilting head, insert nozzle into nostril. Fully depress rim with a firm, even stroke and sniff deeply. Wipe nozzle clean after use.

• adults and children 6 to under 12 years of age (with adult supervision): 2 or 3 sprays in each nostril not more often than every 10 to 12 hours. Do not exceed 2 doses in any 24-hour period.  
• children under 6 years of age: ask a doctor.

### Drug Facts (continued)

**Other information**

• store between 20° to 25°C (68° to 77°F)

• refer to carton for future reference on full labeling

### Inactive ingredients

Benzalkonium Chloride, Benzyl Alcohol, Citric Acid Anhydrous, Disodium Phosphate, Fragrance, Polyethylene Glycol, Purified Water, Sodium Citrate, Sorbitol

### IMPORTANT:

**KEEP OUTER CARTON FOR FUTURE REFERENCE ON FULL LABELING**

### Cap Removal Instructions:

To remove cap, squeeze cap at base on opposite sides on the two firm impressions with fingers. While squeezing, twist cap counterclockwise to remove. Press down to lock. Refer to arrow on cap. To reapply cap, turn cap on pump clockwise until it locks.

### Walgreens Pharmacy Recommendation

Walgreens Pharmacy Survey

This product is not manufactured or distributed by Procter & Gamble, the distributor of Vicks® Sinex™ Severe.

### Questions or comments?

1-800-222-1007

DISTRIBUTED BY: WALGREENS CO.

200 WILMOT RD. DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED

walgreens.com ©2010 Walgreen Co.

Walgreens

Compare to Vicks® Sinex™ Severe active ingredient!

NDC 0583-2205-02

# Severe Sinus Relief Mist

OXYMETAZOLINE HCl 0.05% / NASAL DECONGESTANT

SEVERE

12 HOUR

ULTRA-FINE SPRAY

• Helps relieve congestion & sinus pressure

0.5 FL OZ (15 mL)

NET WT

LOT

ORG0518-F

FSP





## SEVERE SINUS RELIEF MIST

oxymetazoline spray liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0363-2305
<b>Route of Administration</b>	NASAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>OXYMETAZOLINE HYDROCHLORIDE</b> (UNII: K89MJ0S5VY) (OXYMETAZOLINE - UNII:8VLN5B44ZY)	OXYMETAZOLINE HYDROCHLORIDE	50 mg in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>SODIUM PHOSPHATE, MONOBASIC</b> (UNII: 3980JIH2SW)	
<b>SODIUM PHOSPHATE, DIBASIC</b> (UNII: GR686LBA74)	
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>BENZALKONIUM CHLORIDE</b> (UNII: F5UM2KM3W7)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>MENTHOL</b> (UNII: L7T10EIP3A)	
<b>CAMPHOR (SYNTHETIC)</b> (UNII: 5TJD82A1ET)	
<b>EUCALYPTOL</b> (UNII: RV6J6604TK)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-2305-52	1 in 1 CARTON	08/30/2018	
1		15 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/30/2018	

**Labeler** - Walgreens Company (008965063)