

CYSTADANE - betaine powder, for solution
Rare Disease Therapeutics, Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Cystadane safely and effectively. See full prescribing information for Cystadane.

Cystadane (R) (betaine anhydrous for oral solution) powder
Initial U.S. Approval: 1996

----- **RECENT MAJOR CHANGES** -----

Changes of Address 08/2008
Address of adverse reactions 07/2010
Change of distributor 07/2014
Manufactured For / Under License From 07/2014

----- **INDICATIONS AND USAGE** -----

Cystadane is a methylating agent indicated for the treatment of homocystinuria to decrease elevated homocysteine blood levels. Included within the category of homocystinuria are (1):

- Cystathionine beta-synthase (CBS) deficiency
- 5,10-methylenetetrahydrofolate reductase (MTHFR) deficiency
- Cobalamin cofactor metabolism (cbl) defect

----- **DOSAGE AND ADMINISTRATION** -----

- Usual dose in adult and pediatric patients is 6 grams per day, administered orally in divided doses of 3 grams two times a day.(2)
- In children less than 3 years of age, may initiate dosing at 100 mg/kg/day, divided in twice daily doses, and then increased weekly by 50 mg/kg increments.(2)
- Dose can be gradually increased until plasma total homocysteine is undetectable or present only in small amounts.(2)
- Monitor patient response by plasma homocysteine levels.(2)
- Prescribed amount of Cystadane should be measured with the measuring scoop provided and then dissolved in 4 to 6 ounces of water, juice, milk, or formula, or mixed with food for immediate ingestion.(2)

----- **DOSAGE FORMS AND STRENGTHS** -----

- Powder for oral solution available in bottles containing 180 grams of betaine anhydrous.(3)

----- **CONTRAINDICATIONS** -----

- None (4)

----- **WARNINGS AND PRECAUTIONS** -----

- Hypermethioninemia: Cystadane may worsen elevated plasma methionine concentrations in patients with CBS deficiency. Cerebral edema has been reported in patients receiving Cystadane.(5.1)
- Monitoring: Monitor plasma methionine concentrations in patients with CBS deficiency. Keep plasma methionine concentrations below 1,000 µmol/L through dietary medication and, if necessary, a reduction of Cystadane dose. (5.1)

----- **ADVERSE REACTIONS** -----

- Most common adverse reactions (incidence > 2%) were nausea and gastrointestinal distress, based on physician survey.

To report SUSPECTED ADVERSE REACTIONS, contact 877-828-8874, or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

----- **USE IN SPECIFIC POPULATIONS** -----

- Pregnancy: Animal reproduction studies have not been conducted with Cystadane. Use only if clearly needed.(8.1)
- Nursing women: It is not known whether Cystadane is excreted in human milk. Use only if clearly needed.(8.3)
- Pediatrics: Pediatric patients ranging in age from 24 days to 17 years have been treated with Cystadane. Children younger than 3 years of age may benefit from dose titration.

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 7/2014

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

2.1 Dosage

2.2 Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

5.1 Hypermethioninemia

6 ADVERSE REACTIONS

6.1 Adverse Reactions in Clinical Studies

6.2 Postmarketing Experience

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

8.3 Nursing Mothers

8.4 Pediatric Use

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 Storage

17 PATIENT COUNSELING INFORMATION

17.1 Dosing and Administration

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Cystadane® (betaine anhydrous for oral solution) is indicated for the treatment of homocystinuria to decrease elevated homocysteine blood levels. Included within the category of homocystinuria are:

- Cystathionine beta-synthase (CBS) deficiency
- 5,10-methylenetetrahydrofolate reductase (MTHFR) deficiency
- Cobalamin cofactor metabolism (cbl) defect

2 DOSAGE AND ADMINISTRATION

2.1 Dosage

The usual dosage in adult and pediatric patients is 6 grams per day administered orally in divided doses of 3 grams twice daily. In pediatric patients less than 3 years of age, dosage may be started at 100 mg/kg/day divided in twice daily doses, and then increased weekly by 50 mg/kg increments.

Therapy with Cystadane should be directed by physicians knowledgeable in the management of patients with homocystinuria. Patient response to Cystadane can be monitored by homocysteine plasma levels. Dosage in all patients can be gradually increased until plasma total homocysteine is undetectable or present only in small amounts. Response (by homocysteine plasma levels) usually occurs within several days and steady state within a month. Plasma methionine concentrations should be monitored in patients with CBS deficiency [*See Warnings and Precautions (5.1)*].

Dosages of up to 20 grams per day have been necessary to control homocysteine levels in some patients. However, one pharmacokinetic and pharmacodynamic *in vitro* simulation study indicated minimal benefit from exceeding a twice-daily dosing schedule and a 150 mg/kg/day dosage for Cystadane.

2.2 Administration

The prescribed amount of Cystadane should be measured with the measuring scoop provided (one level 1.7 mL scoop is equal to 1 gram of betaine anhydrous powder) and then dissolved in 4 to 6 ounces (120 to 180 mL) of water, juice, milk, or formula, or mixed with food for immediate ingestion.

3 DOSAGE FORMS AND STRENGTHS

Cystadane is a white, granular, hygroscopic powder for oral solution available in bottles containing 180 grams of betaine anhydrous.

4 CONTRAINDICATIONS

None.

5 WARNINGS AND PRECAUTIONS

5.1 Hypermethioninemia

Risk of Hypermethioninemia in Patients with CBS Deficiency

Patients with homocystinuria due to cystathionine beta-synthase (CBS) deficiency may also have elevated plasma methionine concentrations. Treatment with Cystadane may further increase methionine concentrations due to the remethylation of homocysteine to methionine. Cerebral edema has been reported in patients with hypermethioninemia, including patients treated with Cystadane. Plasma methionine concentrations should be monitored in patients with CBS deficiency. Plasma methionine concentrations should be kept below 1,000 $\mu\text{mol/L}$ through dietary modification and, if necessary, a reduction of Cystadane dose.

6 ADVERSE REACTIONS

6.1 Adverse Reactions in Clinical Studies

The most serious adverse reaction reported with Cystadane treatment is the development of hypermethioninemia and cerebral edema in patients with CBS Deficiency [*see Warnings and Precautions (5.1)*].

The assessment of clinical adverse reactions is based on a survey study of 41 physicians, who treated a total of 111 homocystinuria patients with Cystadane. Adverse reactions were retrospectively recalled

and were not collected systematically in this open-label, uncontrolled, physician survey. Thus, this list may not encompass all types of potential adverse reactions, reliably estimate their frequency, or establish a causal relationship to drug exposure. The following adverse reactions were reported (Table 1):

Table 1: Number of Patients with Adverse Reactions to Cystadane by Physician Survey

Adverse Reactions	Number of Patients
Nausea	2
Gastrointestinal distress	2
Diarrhea	1
"Bad Taste"	1
"Caused Odor"	1
Questionable psychological changes	1
"Aspirated the powder"	1

6.2 Postmarketing Experience

The following adverse reactions have been identified during post approval use of Cystadane. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

In postmarketing experience with Cystadane, severe cerebral edema and hypermethioninemia have been reported within 2 weeks to 6 months of starting betaine therapy, with complete recovery after discontinuation of Cystadane. All patients who developed cerebral edema had homocystinuria due to CBS deficiency and had severe elevation in plasma methionine levels (range 1,000 to 3,000 μM). As cerebral edema has also been reported in patients with hypermethioninemia, secondary hypermethioninemia due to betaine therapy has been postulated as a possible mechanism of action.

The following adverse reactions have been reported in patients during postmarketing use of Cystadane: anorexia, agitation, depression, irritability, personality disorder, sleep disturbed, dental disorders, diarrhea, glossitis, nausea, stomach discomfort, vomiting, hair loss, hives, skin odor abnormalities, and urinary incontinence.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C: Animal reproduction studies have not been conducted with Cystadane. It is also not known whether Cystadane can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Cystadane should be given to a pregnant woman only if clearly needed.

8.3 Nursing Mothers

It is not known whether Cystadane is excreted in human milk. Use only if clearly needed.

8.4 Pediatric Use


The majority of case studies of homocystinuria patients treated with Cystadane have been pediatric patients, including patients ranging in age from 24 days to 17 years [see *Clinical Studies (14)*]. Children younger than 3 years of age may benefit from dose titration [see *Dosage and Administration (2)*].

10 OVERDOSAGE

In an acute toxicology study in rats, death occurred frequently at doses equal to or greater than 10 g/kg.

11 DESCRIPTION

Cystadane (betaine anhydrous for oral solution) is an agent for the treatment of homocystinuria. It contains no ingredients other than anhydrous betaine. Cystadane is a white, granular, hygroscopic powder, which is diluted in water and administered orally. The chemical name of betaine anhydrous powder is trimethylglycine. It has a molecular weight of 117.15. The structural formula is:



12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Cystadane acts as a methyl group donor in the remethylation of homocysteine to methionine in patients with homocystinuria. Cystadane occurs naturally in the body. It is a metabolite of choline and is present in small amounts in foods such as beets, spinach, cereals, and seafood.

12.2 Pharmacodynamics

Cystadane was observed to lower plasma homocysteine levels in three types of homocystinuria, including CBS deficiency; MTHFR deficiency; and cbl defect. Patients have taken Cystadane for many years without evidence of tolerance. There has been no demonstrated correlation between Cystadane levels and homocysteine levels.

In CBS-deficient patients, large increases in methionine levels over baseline have been observed. Cystadane has also been demonstrated to increase low plasma methionine and S-adenosylmethionine (SAM) levels in patients with MTHFR deficiency and cbl defect.

12.3 Pharmacokinetics

Pharmacokinetic studies of Cystadane are not available. Plasma levels of Cystadane have not been measured in patients and have not been correlated to homocysteine levels.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term carcinogenicity and fertility studies have not been conducted with Cystadane. No evidence of genotoxicity was demonstrated in the following tests: metaphase analysis of human lymphocytes; bacterial reverse mutation assay; and mouse micronucleus test.

14 CLINICAL STUDIES

Cystadane was studied in a double-blind, placebo-controlled, crossover study in 6 patients with CBS deficiency, ages 7 to 32 years at enrollment. Cystadane was administered at a dosage of 3 grams twice daily, for 12 months. Plasma homocysteine levels were significantly reduced ($p < 0.01$) compared to placebo. Plasma methionine levels were variable and not significantly different compared to placebo.

No adverse events were reported in any patient.

Cystadane has also been evaluated in observational studies without concurrent controls in patients with homocystinuria due to CBS deficiency, MTHFR deficiency, or cbl defect. A review of 16 case studies and the randomized controlled trial previously described was also conducted, and the data available for each study were summarized; however, no formal statistical analyses were performed. The studies included a total of 78 male and female patients with homocystinuria who were treated with Cystadane. This included 48 patients with CBS deficiency, 13 with MTHFR deficiency, and 11 with cbl defect, ranging in age from 24 days to 53 years. The majority of patients (n=48) received 6 gm/day, 3 patients received less than 6 gm/day, 12 patients received doses from 6 to 15 gm/day, and 5 patients received doses over 15 gm/day. Most patients were treated for more than 3 months (n=57) and 30 patients were treated for 1 year or longer (range 1 month to 11 years). Homocystine is formed nonenzymatically from two molecules of homocysteine, and both have been used to evaluate the effect of Cystadane in patients with homocystinuria. Plasma homocystine or homocysteine levels were reported numerically for 62 patients, and 61 of these patients showed decreases with Cystadane treatment. Homocystine decreased by 83-88% regardless of pre-treatment level, and homocysteine decreased by 71-83%, regardless of the pre-treatment level. Clinical improvement, such as improvement in seizures, or behavioral and cognitive functioning, was reported by the treating physicians in about three-fourths of patients. Many of these patients were also taking other therapies such as vitamin B6 (pyridoxine), vitamin B12 (cobalamin), and folate with variable biochemical responses. In most cases, adding Cystadane resulted in a further reduction of either homocystine or homocysteine.

16 HOW SUPPLIED/STORAGE AND HANDLING

Cystadane is available in plastic bottles containing 180 grams of betaine anhydrous. Each bottle is equipped with a plastic child-resistant cap and is supplied with a polystyrene measuring scoop. One level scoop (1.7 mL) is equal to 1 gram of betaine anhydrous powder.

NDC 66621-4000-1 180 g/bottle

Cystadane can be ordered by calling AnovoRx Group, LLC, Customer service at 1-888-487-4703

16.1 Storage

Store at room temperature, 15 – 30 °C (59 – 86 °F). Protect from moisture.

17 PATIENT COUNSELING INFORMATION

Patients should be advised of the following information before beginning treatment with Cystadane:

17.1 Dosing and Administration

- Instruct patients and caregivers that Cystadane should only be taken as directed by their healthcare professional.
 - Instruct patients and caregivers to administer Cystadane as follows:
 - Shake bottle lightly before removing cap.
- Measure with the scoop provided.
- Measure the number of scoops as prescribed by their healthcare professional. One level scoop (1.7 mL) is equivalent to 1 gram of betaine anhydrous powder.
- Mix powder with 4 to 6 ounces (120 to 180 mL) of water, juice, milk, or formula until completely dissolved, or mix with food, then ingest mixture immediately.
- Always replace the cap tightly after using, and protect powder from moisture.

Manufactured For:

Rare Disease Therapeutics, Inc.

Franklin, TN 37067

Under License From:
Orphan Europe, s.a.r.l. Puteaux France

Distributed By:
AnovoRx Distribution, LLC
Memphis, TN 38134

Part No.: RDT C PI007
Part No.: Orphan Europe OEP 829

3" X 7"

Die line

Manufactured for:
Rare Disease Therapeutics, Inc
Franklin, Tennessee 37067
Under License from:
Orphan Europe, s.a.r.l.
Puteaux France
Distributed by:
AnovoRx Distribution, LLC
Memphis, Tennessee 38134

RDT

NDC 66621-4000-1

Cystadane®
*(betaine anhydrous
for oral solution)*

1 gram of betaine anhydrous
powder per scoopful

180 grams

Dosage and Administration:
See package insert for dosage information.

Preparation:

1. Shake bottle lightly before removing cap.
2. Measure with the scoop provided.
3. One level scoop (1.7 cc) is equivalent to 1 gram of betaine anhydrous powder. Measure the number of scoops your physician has prescribed.
4. Mix with 4 to 6 ounces (120 to 180 mL) of water, juice, milk, or formula, until completely dissolved, or mix with food, then ingest immediately.

Keep this and all medication out of the reach of children.
Always replace the cap tightly after using.
Store at room temperature, 15°-30°C (59°-86°F). Protect from moisture.

Lot #:
Exp:
Rev Date: July 2014
Part No.: RDT C L003
OEP824

Made in Germany

Rx only



Black PMS Match Spot non vernis

CYSTADANE

betaine powder, for solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:66621-4000
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BETAINE (UNII: 3SCV180C9W) (BETAINE - UNII:3SCV180C9W)	BETAINE	6 g

Product Characteristics

Color	white (none)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66621-4000-1	180 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020576	10/25/1996	

Labeler - Rare Disease Therapeutics, Inc. (966133100)**Registrant** - Rare Disease Therapeutics, Inc. (966133100)**Establishment**

Name	Address	ID/FEI	Business Operations
Rare Disease Therapeutics, Inc.		966133100	relabel(66621-4000)

Establishment

Name	Address	ID/FEI	Business Operations
Ropack Pharmaceutical		209989631	manufacture(66621-4000)

Revised: 1/2015

Rare Disease Therapeutics, Inc.