

PEPTO BISMOL DIARRHEA LIQUICAPS- bismuth subsalicylate capsule, liquid filled

The Procter & Gamble Manufacturing Company

Pepto Bismol[®] Diarrhea LiquiCaps

Drug Facts

Active ingredient (in each LiquiCap)

Bismuth subsalicylate 262 mg

Purpose

Antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea

Warnings

Reye's syndrome:

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert:

Contains salicylate.

Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use

if you are taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- swallow with water, do not chew
- adults and children 12 years and over:
 - 2 LiquiCaps every ½ hour or 4 LiquiCaps every 1 hour as needed
 - do not exceed 16 LiquiCaps in 24 hours
 - use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each LiquiCap contains:**
 - salicylate 199 mg
 - low sodium
 - do not exceed 25°C

Inactive ingredients

D&C red No. 33, gelatin, glycerin, polyethylene glycol, polysorbate 20, povidone, sorbitol sorbitan solution, titanium dioxide

Questions?

1-800-717-3786

TAMPER EVIDENT: THIS PACKAGE IS SAFETY SEALED & CHILD RESISTANT. USE ONLY IF BLISTERS ARE INTACT. IF DIFFICULT TO OPEN, USE SCISSORS.

MADE IN CANADA

DIST. BY PROCTER & GAMBLE,
CINCINNATI OH 45202

PRINCIPAL DISPLAY PANEL - 48 LiquiCap Carton

FAST & SOOTHING RELIEF

Pepto®

Diarrhea

Bismuth Subsalicylate

Antidiarrheal

LIQUICAPS

COATS & KILLS BACTERIA THAT CAUSE DIARRHEA

48 LIQUICAPS™

**Pepto
Bismol®
Diarrhea**
LIQUICAPS



FAST & SOOTHING RELIEF

Pepto® Bismuth Subsalicylate
Antidiarrheal
Diarrhea
LIQUICAPS



**COATS & KILLS
BACTERIA THAT
CAUSE DIARRHEA**



Actual Size

48 LIQUICAPS™

TAMPER EVIDENT:
THIS PACKAGE IS SAFETY SEALED & CHILD
RESISTANT. USE ONLY IF BLISTERS ARE
INTACT. IF DIFFICULT TO OPEN, USE SCISSORS.

MADE IN CANADA
DIST. BY: PROCTER & GAMBLE,
CINCINNATI, OH 45202

Also sold as Pepto Bismol® LiquiCaps

P&G www.pg.com pepto-bismol.com
Patents: www.pg.com/patents



Drug Facts	
Active ingredient (in each LiquiCap) Bismuth subsalicylate 262 mg	Purpose Antidiarrheal
Uses relieves • travelers' diarrhea • diarrhea	
Warnings Riley's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Riley's syndrome, a rare but serious illness.	
Allergy alert: Contains salicylate. Do not take if you are • allergic to salicylates (including aspirin) • taking other salicylate products	
Do not use if you have • an ulcer • a bleeding problem • bloody or black stool	
Ask a doctor before use if you have • fever • mucus in the stool	
Ask a doctor or pharmacist before use if you are taking any drug for • anticoagulation (thinning the blood) • diabetes • gout • arthritis	
When using this product: a temporary, but harmless, darkening of the stool and/or tongue may occur	
Stop use and ask a doctor if • symptoms get worse • ringing in the ears or loss of hearing occurs • diarrhea lasts more than 2 days	
If pregnant or breast feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions • swallow with water; do not chew • adults and children 12 years and over: • 2 LiquiCaps every 1/2 hour or 4 LiquiCaps every 1 hour as needed • do not exceed 16 LiquiCaps in 24 hours • use until diarrhea stops but not more than 2 days • children under 12 years: ask a doctor • drink plenty of clear fluids to help prevent dehydration caused by diarrhea	
Other information • each LiquiCap contains: • salicylate 199 mg • low sodium • do not exceed 25°C	
Inactive ingredients D&C Red No. 33, gelatin, glycerin, polyethylene glycol, polysorbate 20, povidone, sorbitol sorbitan solution, titanium dioxide	
Questions? 1-800-717-3786	



90240138



PEPTO BISMOL DIARRHEA LIQUICAPS

bismuth subsalicylate capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37000-537
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	262 mg

Inactive Ingredients

Ingredient Name	Strength
POLYSORBATE 20 (UNII: 7T1F30V5YH)	
GLYCERIN (UNII: PDC6A3C0OX)	
SORBITOL (UNII: 506T60A25R)	

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
GELATIN (UNII: 2G86QN327L)	
POVIDONE (UNII: FZ989GH94E)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	

Product Characteristics

Color	pink	Score	no score
Shape	CAPSULE	Size	14mm
Flavor		Imprint Code	PB
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37000-537-12	1 in 1 CARTON	03/01/2019	06/30/2026
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:37000-537-24	2 in 1 CARTON	03/01/2019	08/20/2024
2		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:37000-537-48	4 in 1 CARTON	03/01/2019	06/30/2026
3		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	03/01/2018	06/30/2026

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 11/2023

The Procter & Gamble Manufacturing Company