

CHLORAPREP ONE-STEP- chlorhexidine gluconate and isopropyl alcohol solution
CareFusion 213, LLC

ChloraPrep 3mL Sterile Solution Applicators: Clear, Hi-Lite Orange

WARNING FLAMMABLE.

Keep away from fire or flame.

Active ingredients

Chlorhexidine gluconate 2% w/v

Isopropyl alcohol 70% v/v

Purposes

Antiseptic

Antiseptic

Use

for the preparation of the patient's skin prior to surgery. Helps to reduce bacteria that potentially can cause skin infection.

Warnings

For external use only. Flammable, keep away from fire or flame. To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.** •
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (**minimum of 3 minutes on hairless skin; up to 1 hour in hair**)
- do not allow solution to pool
- remove wet materials from prep area

Allergy alert:

This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing • shock • facial swelling • hives • rash

If an allergic reaction occurs, stop use and seek medical help right away.

Do not use

- on patients allergic to chlorhexidine gluconate or any other ingredient in this product
- for lumbar puncture or in contact with the meninges
- on open skin wounds or as a general skin cleanser

When using this product

keep out of eyes, ears, and mouth. May cause serious or permanent injury if permitted to enter and remain. If contact occurs, rinse with cold water right away and contact a doctor.

Stop use and ask a doctor if

irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- use with care in premature infants or infants under 2 months of age.

These products may cause irritation or chemical burns. • use in a well ventilated area

- maximal treatment area for one applicator is approximately **4 in. x 5 in. (130 cm²)**
- remove applicator from package; do not touch sponge • hold the applicator with the sponge down. Pinch wings **only once** to activate the ampule and release the antiseptic.
- wet the sponge by pressing and releasing the sponge against the treatment area until liquid

is visible on the skin • completely wet the treatment area with antiseptic • **dry surgical sites** (e.g., abdomen or arm): use gentle repeated back-and-forth strokes of the sponge for

approximately 30 seconds Allow solution to air dry for approximately 30 seconds. Do not blot

or wipe away. • **moist surgical sites** (e.g., inguinal fold): use gentle repeated back-and-forth

strokes for approximately 2 minutes. • Allow solution to air dry for approximately 1 minute. Do not blot or wipe away. • **if using an ignition source, allow the**

solution to

completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in hair).

Do not blot

or wipe away. • discard the applicator after a single use along with any portion of the solution

not required to cover the prep area. It is not necessary to use the entire amount available.

Other Information

• store between 15-30 °C (59-86 °F) • avoid freezing and excessive heat above 40 °C (104 °F)


Inactive Ingredient

USP purified water

Questions?

www.chloraprep.com • call **1-800-523-0502** (M-F 8 a.m.-5 p.m. CST)

Principal Display Panel-Carton 3mL Sterile Solution Applicators: Clear, Hi-Lite Orange



Principal Display Panel-Carton

BD Chloraprep Clear

3 mL Applicators

2% w/v chlorhexidine gluconate (CHG)

and 70% v/v Isopropyl alcohol (IPA)

Patient Preoperative Skin Preparation

Sterile Solution

Applicator is sterile if package is intact

Store between 15-30 °C (59-86 °F)

Avoid freezing and excessive heat above 40 °C (104 °F)

0.10 fl. oz. (3 mL) each

25 Applicators

NDC 54365-400-32

REF 930400

WARNING

FLAMMABLE

Keep away from fire or flame

To reduce risk of fire, PREP CAREFULLY:

- solution contains alcohol and gives off **flammable vapors**
- avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. **Wet hair is flammable.**
- do not drape or use ignition source (e.g., cautery, laser) until solution is completely

dry (**minimum of 3 minutes on hairless skin; up to 1 hour in**

hair) • do not allow solution to pool • remove wet materials from prep area

Not made with natural rubber latex

Do not reuse

STERILE EO


STERILE SOLUTION

External Use Only

Professional Use Only

CareFusion 213, LLC, El Paso, TX 79912, subsidiary of Becton, Dickinson and Co.

Made in the USA of US and imported parts. www.bd.com



Principal Display Panel-Carton

BD ChloraPrep Hi-Lite Orange

3 mL Applicators

2% w/v chlorhexidine gluconate (CHG)

and 70% v/v Isopropyl alcohol (IPA)

Patient Preoperative Skin Preparation

Sterile Solution

Applicator is sterile if package is intact

Store between 15-30 C (59-86 F)

Avoid freezing and excessive heat above 40 C (104 F)

0.10 fl. oz. (3 mL) each

25 Applicators

NDC 54365-400-33

REF 930415

WARNING

FLAMMABLE

Keep away from fire or flame

To reduce risk of fire, PREP CAREFULLY:

solution contains alcohol and gives off flammable vapors

avoid getting solution into hairy areas. Hair may take up to 1 hour to dry. Wet hair is flammable.

do not drape or use ignition source (e.g., cautery, laser) until solution is completely dry (minimum of 3 minutes on hairless skin; up to 1 hour in

hair) • do not allow solution to pool • remove wet materials from prep area

Not made with natural rubber latex

Do not reuse

STERILE EO

STERILE SOLUTION

External Use Only

Professional Use Only

CareFusion 213, LLC, El Paso, TX 79912, subsidiary of Becton, Dickinson and Co.

Made in the USA of US and imported parts. www.bd.com

CHLORAPREP ONE-STEP

chlorhexidine gluconate and isopropyl alcohol solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54365-400
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)		CHLORHEXIDINE GLUCONATE	20 mg in 1 mL	
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)		ISOPROPYL ALCOHOL	0.7 mL in 1 mL	
Inactive Ingredients				
Ingredient Name		Strength		
WATER (UNII: 059QF0KO0R)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54365-400-32	25 in 1 CARTON	05/24/2019	
1		1 in 1 POUCH		
1		3 mL in 1 APPLICATOR; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		
2	NDC:54365-400-33	25 in 1 CARTON	05/24/2019	
2		1 in 1 POUCH		
2		3 mL in 1 APPLICATOR; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
NDA	NDA020832	05/24/2019		

Labeler - CareFusion 213, LLC (826496312)

Registrant - Becton, Dickinson and Company (832696038)

Establishment

Name	Address	ID/FEI	Business Operations
CareFusion 213, LLC		826496312	analysis(54365-400) , label(54365-400) , manufacture(54365-400) , pack(54365-400) , sterilize(54365-400)

Revised: 11/2023

CareFusion 213, LLC