

TIGER SUPPLY INC APF- sodium fluoride gel
Tiger Supply Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Active Ingredient:

Fluoride Ion 1.23%

Available from 2.09% Sodium Fluoride and Hydrofluoric Acid

Purpose:

Fluoride Treatment Gel

Indications and Usage:

- A stable thixotropic fluoride treatment gel used to help prevent dental decay.
- For Professional Use Only. This product is not intended for home or unsupervised use.

Warnings:

- Keep out of reach of children.
- Do not swallow. If product is accidentally swallowed in quantities greater than would normally occur with a treatment gel, seek medical help or contact a Poison Control Center right away.
- Read directions carefully before using.

Dosage and Administration:

Shake well before use. This is a one minute or four minute fluoride gel for in-office patient use. It is normally used as a preventative caries treatment two times a year.

1. After thorough prophylaxis, fill two single or one dual tray, one third full with gel. Air dry teeth and insert trays into the mouth.
2. Instruct patient to bite down lightly but firmly for one minute (80% effectiveness) or four minutes (100% effectiveness).
3. Remove trays. Instruct patient to expectorate any excess gel and not to eat or drink for at least 30 minutes.

Other Information:

Store at controlled room temperature 59° to 86°F (15°-30°C).

Protect from freezing.

Inactive Ingredients:

Citric Acid, FD&C Red #40, FD&C Yellow #5 (tartrazine) as a color additive, Flavor, Magnesium Aluminum Silicate, Phosphoric Acid, Polysorbate 20, Sodium Benzoate, Sodium Saccharin, Titanium Dioxide, Water, Xanthan Gum, Xylitol.



APF GEL

THIXOTROPIC
with Xylitol

Acidulated Phosphate Fluoride
Treatment Gel
1.23% Fluoride Ion

GLUTEN FREE

RxOnly

Cotton Candy

Re-order#: TIG-24-45777

IMPORTANT: READ DIRECTIONS FOR PROPER USE

MADE IN USA

Net wt. 16 oz (454 g)

Active Ingredient:

Fluoride Ion 1.23% Fluoride Treatment Gel
Available from 2.09% Sodium Fluoride and Hydrofluoric Acid.

Purpose:

Indications and Usage:

- A stable thixotropic fluoride treatment gel used to help prevent dental decay.
- For Professional Office Use Only. This product is not intended for home or unsupervised consumer use.

Warnings:

- Keep out of reach of children.
- Do not swallow. If product is accidentally swallowed in quantities greater than would normally occur with a treatment gel, seek medical help or contact a Poison Control Center right away.
- Read directions carefully before using.

Dosage and Administration:

Shake well before use. This is a one minute or four minute fluoride gel for in-office patient use. It is normally used as a preventative caries treatment two times a year.

1. After thorough prophylaxis, fill two single or one dual tray, one third full with gel. Air dry teeth and insert trays into the mouth.
2. Instruct patient to bite down lightly but firmly for one minute (80% effectiveness) or four minutes (100% effectiveness).
3. Remove trays. Instruct patient to expectorate any excess gel and not to eat or drink for at least 30 minutes.

Other Information:

- Store at controlled room temperature 59° to 86°F (15°-30°C).
Protect from freezing.

Inactive Ingredients:

Citric Acid, FD&C Blue #1, Flavor, Magnesium Aluminum Silicate, Phosphoric Acid, Polysorbate 20, Sodium Benzoate, Sodium Saccharin, Titanium Dioxide, Water, Xanthan Gum, Xylitol.

Manufactured for:



99 Sherwood Avenue
Farmingdale, NY 11735
Tel: 631.293.2700
www.tigersupplyinc.com

99092287 Rev 03/2013

TIGER SUPPLY INC APF

sodium fluoride gel

Product Information

Product Type

HUMAN PRESCRIPTION DRUG

Item Code (Source)

NDC:69484-056

| | |
|--------------------------------|--------|
| Route of Administration | DENTAL |
|--------------------------------|--------|

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------------|
| SODIUM FLUORIDE (UNII: 8ZYQ1474W7) (FLUORIDE ION - UNII:Q80VPU408O) | FLUORIDE ION | 5.6 g in 454 g |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP) | |
| MAGNESIUM ALUMINUM SILICATE (UNII: 6M3P64V0NC) | |
| PHOSPHORIC ACID (UNII: E4GA8884NN) | |
| POLYSORBATE 20 (UNII: 7T1F30V5YH) | |
| SODIUM BENZOATE (UNII: OJ245FE5EU) | |
| SACCHARIN SODIUM (UNII: SB8ZUX40TY) | |
| WATER (UNII: 059QF0KO0R) | |
| XANTHAN GUM (UNII: TTV12P4NEE) | |
| XYLITOL (UNII: VCQ006KQ1E) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| FD&C BLUE NO. 1 (UNII: H3R47K3TBD) | |

Product Characteristics

| | | | |
|-----------------|--------------|---------------------|--|
| Color | blue | Score | |
| Shape | | Size | |
| Flavor | COTTON CANDY | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:69484-056-15 | 454 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product | 03/01/2013 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------------|--|----------------------|--------------------|
| unapproved drug other | | 03/01/2013 | |

Labeler - Tiger Supply Inc (168720709)