

**ACETAMINOPHEN- acetaminophen tablet**  
**CVS PHARMACY, INC**

-----  
**Rapid Release Gelcaps**  
**EXTRA STRENGTH**  
**Acetaminophen Gelcaps USP, 500 mg**  
**Pain reliever; Fever reducer**  
**Aspirin free**

**Active ingredient**

(in each Gelcap)

Acetaminophen, USP 500 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- headache
- muscular aches
- backache
- minor pain of arthritis
- toothache
- the common cold
- premenstrual and menstrual cramps
- temporarily reduces fever

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert**

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

liver disease.

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

### **Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of the reach of children.**

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

#### **■ do not take more than directed**

adults and children 12 years and over

- take 2 gels every 6 hours while symptoms last

- do not take more than 6 gelcaps in 24 hours, unless directed by a doctor
  - do not take more than 10 days unless directed by a doctor
- children under 12 years
- ask a doctor

### **Other information**

- store at 20°-25°C (68°-77°F). See USP Controlled Room Temperature
- avoid high humidity
- see end panel for lot number and expiration date

### **Inactive ingredients**

ammonium hydroxide, black iron oxide, black iron oxide irradiated, colloidal silicon dioxide, croscarmellose sodium, gelatin, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, hypromellose, iron oxide red, isopropyl alcohol, n-butyl alcohol, polyethylene glycol, povidone, pregelatinized starch, propylene glycol, shellac glaze, stearic acid, titanium dioxide, yellow iron oxide.

### **Questions or comments?**

call **1-877-770-3183** Mon-Fri 9:00 AM to 4:30 PM EST.

### **Acetaminophen, USP 500 mg Rapid Release Gelcaps**



COATING  
FREE AREA



**Important: Read all product information before using. Keep the carton for important information.**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**Drug Facts**

**Active Ingredient (in each gelcap)** **Purpose**  
Acetaminophen, USP 500 mg Pain reliever/fever reducer

**Uses**


- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches
  - backache
  - minor pain of arthritis
  - the common cold
  - toothache
- premenstrual and menstrual cramps
- temporarily reduces fever

**Warnings**

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:
- more than 4,000 mg of acetaminophen in 24 hours
  - with other drugs containing acetaminophen
  - 3 or more alcoholic drinks every day while using this product
- Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:
- skin redness
  - hives
  - itching
  - swelling of the face, lips, or throat
  - difficulty breathing
  - fever
  - skin rash
- If a skin reaction occurs, stop use and seek medical help right away.
- Do not use**
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
  - if you have ever had an allergic reaction to this product or any of its ingredients.
- Ask a doctor before use if you have liver disease.**
- Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

- Stop use and ask a doctor if**
- pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - new symptoms occur
  - redness or swelling is present
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.



COATING  
FREE AREA

**Drug Facts (continued)**  
**Directions**  
do not take more than directed  
adults and children 12 years and over: take 2 gelcaps every 6 hours while symptoms last. Do not take more than 8 gelcaps in 24 hours, unless directed by a doctor.  
children under 12 years: do not take more than 10 days unless directed by a doctor. ask a doctor.

**Other Information**  
Store at 20° - 25°C (68° - 77°F). See USP Controlled Room Temperature. avoid high humidity. see and panel for expiration date and lot number.  
**Inactive Ingredients** ammonium hydroxide, black iron oxide, black iron oxide trihydrate, colloidal silicon dioxide, croscarmellose sodium, gelatin, hydroxypropyl cellulose, hydroxypropyl methylcellulose, hypromellose, iron oxide red, isopropyl alcohol, n-butyl alcohol, polyethylene glycol, polydioxanone, pregelatinized starch, propylene glycol, stearic acid, titanium dioxide, yellow iron oxide.

**Questions or comments?**  
call 1-877-770-3183 Mon-Fri 8:00 AM to 4:30 PM EST.  
\*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Extra Strength Tylenol® Rapid Release Gels.

Distributed by: CVS Pharmacy, Inc.  
One CVS Drive, Woonsocket, RI 02895  
© 2018 CVS/pharmacy  
CVS.com • 1-800-SHOP CVS

Made in India  
V-32577  
CVS Quality  
Money Back Guarantee

CVS Health.  
EXTRA STRENGTH  
Acetaminophen  
Gelcaps USP, 500 mg  
Pain reliever, Fever reducer  
Rapid release  
Aspirin free  
225 GELCAPS

CVS Health.  
EXTRA STRENGTH  
Acetaminophen  
Gelcaps USP, 500 mg  
Pain reliever, Fever reducer  
Rapid release  
Aspirin free  
225 GELCAPS



Actual Size  
Actual Bottle Size  
on Top Panel



Compare to the active ingredient in Extra Strength Tylenol® Rapid Release Gels.  
Rapid Release Gelcaps

Package Contains One Bottle

Actual Size



COATING  
FREE AREA

|                     |                            |
|---------------------|----------------------------|
| END USER : 06RANUGS |                            |
| item                | : CVS APAP RR 100CT        |
| code #              | :                          |
| size                | : 2+3/32 X 2+3/32 X 4+3/16 |
| ref #               | : P2180604A                |
| material            | : .016 SBS                 |
| view                | : !                        |
| date                | : !                        |

COATING FREE AREA

R - 0 2

300000078

525000006

**Drug Facts (continued)**

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed

|                                       |   |
|---------------------------------------|---|
| adults and children 12 years and over | <ul style="list-style-type: none"> <li>■ take 2 gelscaps every 6 hours while symptoms last</li> <li>■ do not take more than 6 gelscaps in 24 hours, unless directed by a doctor</li> <li>■ do not take more than 10 days unless directed by a doctor</li> </ul> |
| children under 12 years               | <ul style="list-style-type: none"> <li>■ ask a doctor</li> </ul>  |

**Other information**

- Store at 20° - 25°C (68° - 77°F). See USP Controlled Room Temperature
- avoid high humidity
- see end panel for expiration date and lot number

**Inactive Ingredients:** ammonium hydroxide, black iron oxide, black iron oxide trihydrate, colloidal silicon dioxide, croscarmellose sodium, gelatin, hydroxypropyl cellulose, hydroxypropyl methylcellulose, hypromellose, iron oxide red, isopropyl alcohol, n-butyl alcohol, polyethylene glycol, povidone, propylparaben, propylene glycol, shellac, stearic acid, titanium dioxide, yellow iron oxide.

**Questions or comments?**  
call 1-877-770-3183 Mon-Fri 9:00 AM to 4:30 PM EST.

\*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Extra Strength Tylenol® Rapid Release Gels.

Distributed by: CVS Pharmacy, Inc.  
One CVS Drive, Woonsocket, RI 02895  
© 2018 CVS/pharmacy  
CVS.com® 1-800-SHOP CVS  
Made in India  
V-32577

**CVS Quality**  
Money Back Guarantee

Package Contains One Bottle

Actual Size

CVS Health. Rapid Release Gelscaps

**EXTRA STRENGTH**  
**Acetaminophen**  
Gelscaps USP, 500 mg  
Pain reliever; Fever reducer

- Rapid release
- Aspirin free

100 GELCAPS

CVS Health. Rapid Release Gelscaps

**EXTRA STRENGTH**  
**Acetaminophen**  
Gelscaps USP, 500 mg  
Pain reliever; Fever reducer

- Rapid release
- Aspirin free

100 GELCAPS

Actual Size on Bottom Panel

Compare to the active ingredient in Extra Strength Tylenol® Rapid Release Gels

**RAPID RELEASE**

#317561

0 50428 38730 6

Lot Exp.

COATING FREE AREA

**Important: Read all product information before using. Keep the carton for important information.**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Drug Facts**

| Active ingredient (in each gelscap) | Purpose                     |
|-------------------------------------|-----------------------------|
| Acetaminophen, USP 500 mg...        | Pain reliever/fever reducer |

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - backache
  - the common cold
  - menstrual and menstrual cramps
- temporarily reduce fever

**Warnings**

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- hives
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.





|          |                        |      |          |
|----------|------------------------|------|----------|
| NAME     | 01/15/18 R 1-15/18 R 2 | FORM | 010002   |
| REV      | 01/15/18               | DATE | 01/15/18 |
| REVISION | 1.000 000              | DATE | 01/15/18 |

CONTAINS  
FREE AREA

M - 0 2  
626000004  
300000074



**Drug Facts (continued)**

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Do not use and take a drink if you get pain or fever more than 10 days or fever gets worse or lasts more than 3 days.

If your symptoms occur suddenly or suddenly become worse, these could be signs of a serious reaction.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-325-1235) right away. Present medical information is critical for safety so tell us for children even if you do not notice any signs or symptoms.

**Directions**

Do not take more than directed.

|                                       |   |
|---------------------------------------|---|
| Adults and children 12 years and over | <ul style="list-style-type: none"> <li>Take 2 quick-release tablets with complete liquid.</li> <li>Do not take more than 2 tablets in 24 hours, unless directed by a doctor.</li> <li>Do not take more than 10 days unless directed by a doctor.</li> </ul> |
| Children under 12 years               | <ul style="list-style-type: none"> <li>Ask a doctor.</li> </ul>   |

**Other information**

Store at 20°-25° (68°-77°), the USP Controlled Room Temperature.

CONTAINS  
FREE AREA

300000074  
626000004  
M - 0 2



**Drug Facts (continued)**

Contains high alcohol content. Avoid alcohol while using this product.

**Warnings**

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Do not use and take a drink if you get pain or fever more than 10 days or fever gets worse or lasts more than 3 days.

If your symptoms occur suddenly or suddenly become worse, these could be signs of a serious reaction.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-325-1235) right away. Present medical information is critical for safety so tell us for children even if you do not notice any signs or symptoms.

**Directions**

Do not take more than directed.

|                                       |   |
|---------------------------------------|---|
| Adults and children 12 years and over | <ul style="list-style-type: none"> <li>Take 2 quick-release tablets with complete liquid.</li> <li>Do not take more than 2 tablets in 24 hours, unless directed by a doctor.</li> <li>Do not take more than 10 days unless directed by a doctor.</li> </ul> |
| Children under 12 years               | <ul style="list-style-type: none"> <li>Ask a doctor.</li> </ul>   |

**Other information**

Store at 20°-25° (68°-77°), the USP Controlled Room Temperature.

CONTAINS  
FREE AREA

300000074  
626000004  
M - 0 2





**CVS Health**

Pharmaceuticals  
Overseas

**EXTRA STRENGTH!**

**Acetaminophen**

24 GEL CAPS



**EXTRA STRENGTH!**

**Acetaminophen**

24 GEL CAPS



**CVS Health**

Pharmaceuticals  
Overseas

Compare to the active ingredient in Extra Strength Tylenol® Rapid Release Gels®

24 GEL CAPS

0 50428 28552 7



318884

**Drug Facts**

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Do not use and take a drink if you get pain or fever more than 10 days or fever gets worse or lasts more than 3 days.

If your symptoms occur suddenly or suddenly become worse, these could be signs of a serious reaction.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-325-1235) right away. Present medical information is critical for safety so tell us for children even if you do not notice any signs or symptoms.

**Directions**

Do not take more than directed.


|                                       |   |
|---------------------------------------|---|
| Adults and children 12 years and over | <ul style="list-style-type: none"> <li>Take 2 quick-release tablets with complete liquid.</li> <li>Do not take more than 2 tablets in 24 hours, unless directed by a doctor.</li> <li>Do not take more than 10 days unless directed by a doctor.</li> </ul> |
| Children under 12 years               | <ul style="list-style-type: none"> <li>Ask a doctor.</li> </ul>   |

**Other information**

Store at 20°-25° (68°-77°), the USP Controlled Room Temperature.

CONTAINS  
FREE AREA

300000074  
626000004  
M - 0 2







VALUE SIZE 600 GELCAPS



Compare to the active ingredient in Extra Strength Tylenol® Rapid Release Gels\*

Rapid Release Gelscaps

EXTRA STRENGTH

Acetaminophen

Gelcaps USP, 500 mg

Pain reliever; Fever reducer

Aspirin free



Actual Size



RAPID RELEASE



600 GELCAPS

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL, LINER, CAP & BRACKET OR MISSING

Drug Facts

Active ingredient (in each gelcap)

Acetaminophen, USP 500 mg.....Pain reliever/fever reducer

Uses

Temporarily relieves minor aches and pains due to:
- headache
- minor pain of arthritis
- toothache
- the common cold
- menstrual and muscular cramps
- temporary fevers due to

Warnings

This product contains acetaminophen. Some liver damage may occur if you take:
- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product
- Acetaminophen may cause severe skin reactions. Symptoms may include:
- if a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen
- if you have had an allergic reaction to this product or any of its ingredients
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have liver disease.

Stop use and ask a doctor if:

- you get worse or your condition does not improve after 3 days
- you have pain in your stomach area
- you have a rash or itching
- you are taking the blood thinning drug warfarin.

Keep out of reach of children.

- Do not take more than 10 days
- Do not take more than 3 days
- Do not take more than 10 days unless directed by a doctor

Directions

- Take 2 gelcaps every 6 hours while symptoms last
- Do not take more than 6 gelcaps in 24 hours, unless directed by a doctor
- Do not take more than 10 days unless directed by a doctor

Other information

- Contains 600 gelcaps. See USP Controlled Drug Transactions
- Contains 600 gelcaps. See USP Controlled Drug Transactions
- Contains 600 gelcaps. See USP Controlled Drug Transactions

Questions or comments?

Call 1-877-776-3183 Mon-Fri 9:00 AM to 6:00 PM EST.

\*All trademarks are property of their respective owners. This product is not affiliated with the trademarks of any other company.

© 2018 CVS/Pharmacy. All rights reserved. CVS Health, the CVS Health logo, and the CVS Health name are trademarks of CVS Health. All other trademarks are the property of their respective owners.

CVS/Pharmacy 1-800-SHOP-CVS Made in India. V-35877



#962743



0 50428 54660 4

Lot

Exp.

CVS/Pharmacy



Compare to the active ingredient in Extra Strength Tylenol® Rapid Release Gels\*

Rapid Release Gelscaps

EXTRA STRENGTH

Acetaminophen

Gelcaps USP, 500 mg

Pain reliever; Fever reducer

Aspirin free



Actual Size



400 GELCAPS

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

**Drug Facts**

**Active ingredient (in each gelcap)**

Acetaminophen, USP 500 mg.....Pain reliever/fever reducer

**Purpose**

**Uses** ■ temporarily relieves minor aches and pains due to:  
 ■ headache ■ muscular aches ■ backache ■ minor pain of arthritis  
 ■ toothache ■ the common cold ■ premenstrual and menstrual cramps  
 ■ temporarily reduces fever

**Warnings**

**Liver warnings:** This product contains acetaminophen. Severe liver damage may occur if you take:  
 ■ more than 4,000 mg of acetaminophen in 24 hours  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product  
**Always alert:** acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash ■ peeling  
 If a skin reaction occurs, stop use and seek medical help right away.

**Do not use** with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

**Ask a doctor before use** if you have ever had an allergic reaction to the product or any of its ingredients

**Ask a doctor or pharmacist before use** if you are taking the blood thinning drug warfarin.

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

(CONTINUED ON BACK OF LABEL)

\*All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Extra Strength Tylenol® Rapid Release Gels.

Distributed by: CVS Pharmacy, Inc.  
 One CVS Drive, Woonsocket, RI 02895

CVS.com® 1-800-SHOP CVS

Made in India



#870727

0 50428 43161 0

Lot Exp.  
 3000000 10/22



**Drug Facts (continued)**

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not take more than directed
- take 2 gelcaps every 6 hours while symptoms last
- do not take more than 6 gelcaps in 24 hours, unless directed by a doctor
- do not take more than 10 days unless directed by a doctor

children under 12 years ■ ask a doctor

**Other information**

- store at 20°-25°C (68°-77°F). See USP Controlled Room Temperature
- avoid high humidity
- see and point to expiration date and lot number

**Inactive ingredients:** ammonium hydroxide, black iron oxide, black iron oxide trihydrate, colloidal silicon dioxide, croscarmellose sodium, gelatin, hydroxypropyl cellulose, hydroxypropyl methyl cellulose, hypromellose, iron oxide red, isopropyl alcohol, n-butyl alcohol, polyethylene glycol, polydioxane, pregelatinized starch, polyethylene glycol, stearic acid, titanium dioxide, yellow iron oxide.

**Questions or comments?**  
 Call 1-877-776-3183 Mon-Fri 9:00 AM to 4:30 PM EST.

**ACETAMINOPHEN**

acetaminophen tablet

**Product Information**

|                                |                |                           |               |
|--------------------------------|----------------|---------------------------|---------------|
| <b>Product Type</b>            | HUMAN OTC DRUG | <b>Item Code (Source)</b> | NDC:69842-298 |
| <b>Route of Administration</b> | ORAL           |                           |               |

**Active Ingredient/Active Moiety**

| Ingredient Name  | Basis of Strength | Strength |
|--|-------------------|----------|
| ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN     | 500 mg   |

**Inactive Ingredients**

| Ingredient Name   | Strength |
|---|----------|
| GELATIN (UNII: 2G86QN327L)                              |          |
| FERRIC OXIDE RED (UNII: 1K09F3G675)                     |          |
| POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)     |          |
| HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH) |          |

|   |
|---|
| <b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302)           |
| <b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)            |
| <b>SHELLAC</b> (UNII: 46N107B710)                     |
| <b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)       |
| <b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)          |
| <b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)             |
| <b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)               |
| <b>BUTYL ALCOHOL</b> (UNII: 8PJ61P6TS3)               |
| <b>POVIDONE</b> (UNII: FZ989GH94E)                    |
| <b>STARCH, PREGELATINIZED CORN</b> (UNII: O8232NY3SJ) |
| <b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)                |
| <b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)            |
| <b>FERRIC OXIDE YELLOW</b> (UNII: EX438O2MRT)         |
| <b>AMMONIA</b> (UNII: 5138Q19F1X)                     |

### Product Characteristics

|                 |  |                     |          |
|-----------------|--|---------------------|----------|
| <b>Color</b>    | gray (Encapsulated with red opaque and blue gray opaque hard gelatin shells) | <b>Score</b>        | no score |
| <b>Shape</b>    | OVAL   | <b>Size</b>         | 19mm     |
| <b>Flavor</b>   |  | <b>Imprint Code</b> | G1       |
| <b>Contains</b> |  |                     |          |

### Packaging

| # | Item Code        | Package Description                                | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:69842-298-24 | 24 in 1 BOTTLE; Type 0: Not a Combination Product  | 04/01/2019           |                    |
| 2 | NDC:69842-298-05 | 50 in 1 BOTTLE; Type 0: Not a Combination Product  | 04/01/2019           |                    |
| 3 | NDC:69842-298-10 | 100 in 1 BOTTLE; Type 0: Not a Combination Product | 04/01/2019           |                    |
| 4 | NDC:69842-298-21 | 225 in 1 BOTTLE; Type 0: Not a Combination Product | 04/01/2019           |                    |
| 5 | NDC:69842-298-30 | 300 in 1 BOTTLE; Type 0: Not a Combination Product | 04/01/2019           |                    |
| 6 | NDC:69842-298-40 | 400 in 1 BOTTLE; Type 0: Not a Combination Product | 04/01/2019           |                    |
| 7 | NDC:69842-298-60 | 600 in 1 BOTTLE; Type 0: Not a Combination Product | 04/01/2019           |                    |
| 8 | NDC:69842-298-15 | 150 in 1 BOTTLE; Type 0: Not a Combination Product | 04/01/2019           |                    |

### Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M013                                     | 04/01/2019           |                    |

**Labeler** - CVS PHARMACY, INC (062312574)



