

RITE AID FAST RELIEF LAXATIVE- bisacodyl suppository suppository
Rite Aid Pharmacy

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

RITE AID® Fast Relief Laxative Suppositories 16 ct

Active Ingredient (in each Suppository)

| Active Ingredient (in each Suppository) | Purpose |
|---|--------------------|
| Bisacodyl, 10 mg..... | Stimulant Laxative |

Uses

Uses - For relief of occasional constipation -This product generally produces bowel movement in 15 minutes to 1 hour

Warnings

For rectal use only. May cause abdominal discomfort, faintness, rectal burning, and mild cramps.

Ask a doctor

Ask a doctor -before use if you have noticed a sudden change in bowel habits that periss over a period of two weeks -if you are pregnant or nursing a baby

Stop use and ask a doctor

Stop use and ask a doctor if you have rectal bleeding or fail to have bowel movement after using a laxative. This may indicate a serious condition.

Keep out of reach of children

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

| | |
|---|----------------------------|
| Adults and children 12 years of age and older Children under 6 | Children 6 to 12 years |
| One suppository once daily Ask doctor | 1/2 suppository once daily |

-Detach one suppository from the strip and remove from foil - Carefully insert one suppository well into the rectum

-Do not use more than once per day

Other information

Other information - Store at room temperature: 15° - 30°C (59° - 86°F). Do not exceed 30°C (86°F) - Individually sealed for your protection. Do not use if foil is torn or open

Inactive ingredient

Inactive ingredient hydrogenated vegetable oil

Purpose

| Active Ingredient (in each Suppository) | Purpose |
|---|--------------------|
| Bisacodyl, 10 mg..... | Stimulant Laxative |

Do not use

Do not use -more than one per day -for a period of longer than one week unless directed by a doctor -laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor -if sealed film around the suppository is torn, damaged, or open

Carton Label





RITE AID FAST RELIEF LAXATIVE

bisacodyl suppository suppository

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|----------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:11822-4601 |
| Route of Administration | RECTAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|------------------|
| BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y) | BISACODYL | 10 mg in 2000 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|------------------------------------|--------------------|
| PALM KERNEL OIL (UNII: B0S90M0233) | 1990 mg in 2000 mg |

Product Characteristics

| | | | |
|----------|--------|--------------|------|
| Color | white | Score | |
| Shape | BULLET | Size | 32mm |
| Flavor | | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:11822-4601-7 | 4 in 1 PACKAGE | 10/02/2014 | |
| 1 | NDC:11822-4601-6 | 40 mg in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|-------------------------|--|----------------------|--------------------|
| OTC monograph not final | part334 | 10/02/2014 | |

Labeler - Rite Aid Pharmacy (014578892)

Establishment

| Name | Address | ID/FEI | Business Operations |
|---------------|---------|-----------|-------------------------|
| Unipack, Inc. | | 009248480 | manufacture(11822-4601) |

Revised: 12/2021

Rite Aid Pharmacy