

**HEADACHE RELIEF EXTRA STRENGTH- acetaminophen, aspirin, caffeine tablet
SUPERVALU INC.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Equaline 44-159B

Active ingredients (in each tablet)

Acetaminophen 250 mg
Aspirin 250 mg (NSAID)*
Caffeine 65 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever
Pain reliever
Pain reliever aid

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - a cold
 - arthritis
 - muscular aches
 - toothache
 - premenstrual and menstrual cramps

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

if a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- with other drugs containing acetaminophen
- more than 4,000 mg in 24 hours, which is the maximum daily amount
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- take more or for a longer time than directed
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product

Caffeine warning: The recommended dose of this product contains about as much **caffeine** as a cup of coffee. Limit the use of **caffeine**-containing medications, foods, or beverages while taking this product because too much **caffeine** may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use

- if you ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have asthma
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- stomach bleeding warning applies to you
- you have liver disease

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug or are under a doctor's care for any serious condition

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- an allergic reaction occurs. Seek medical help right away.
- any new symptoms appear
- ringing in the ears or loss of hearing occurs
- pain gets worse or lasts for more than 10 days
- painful area is red or swollen
- fever gets worse or lasts for more than 3 days

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not use more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over
 - take 2 tablets every 6 hours
 - do not take more than 8 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, croscopovidone, hypromellose, microcrystalline cellulose, povidone, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

1-877-932-7948

Principal Display Panel

EQUALINE®

compare to

Excedrin® Extra Strength

active ingredients**

NDC 41163-159-08

extra strength

headache relief

acetaminophen

aspirin (*NSAID*) and caffeine

pain reliever/pain reliever aid

24 tablets

actual size

DOES NOT CONTAIN GLUTEN

**This product is not manufactured or distributed by Novartis AG, owner of the registered trademark Excedrin® Extra Strength.

**DISTRIBUTED BY SUPERVALU INC.
EDEN PRAIRIE, MN 55344 USA 1-877-932-7948**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS
BROKEN OR MISSING**

**KEEP OUTER PACKAGE FOR
COMPLETE PRODUCT INFORMATION**

Drug Facts
Active Ingredients (in each tablet)
Acetaminophen 250 mg Pain reliever
Aspirin 250 mg (NSAID)* Pain reliever
Caffeine 65 mg Pain reliever aid
*nonsteroidal anti-inflammatory drug

Uses
temporarily relieves minor aches and pains due to:
• headache • a cold • arthritis
• toothache • muscular aches
• premenstrual and menstrual cramps

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction, which may include:
• hives
• facial swelling • shock • asthma (wheezing)
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
• rash
• skin reddening • blisters
If a skin reaction occurs, stop use and seek medical help right away.
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
• with other drugs containing acetaminophen
• more than 4,000 mg in 24 hours, which is the maximum daily amount • 3 or more alcoholic drinks every day while using this product

compare to
Excedrin® Extra Strength
active ingredients**
NDC 41163-159-08

EQUALINE®

extra strength
headache relief
acetaminophen
aspirin (NSAID) and caffeine
pain reliever/pain reliever aid

24 tablets

actual size

Drug Facts (continued)
excursions permitted between 15°-30°C (59°-86°F)
• see end flap for expiration date and lot number

Inactive ingredients corn starch, croscopolone, hypromellose, microcrystalline cellulose, povidone, propylene glycol, sodium lauryl sulfate, sodium starch glycolate, stearic acid, titanium dioxide

QUESTIONS OR COMMENTS? 1-877-932-7948
DOES NOT CONTAIN GLUTEN
**This product is not manufactured or distributed by Novartis AG, owner of the registered trademark Excedrin® Extra Strength.

50844 REV0117E15908

**KEEP OUTER PACKAGE FOR
COMPLETE PRODUCT INFORMATION**

Drug Facts (continued)
taking any other drug or are under a doctor's care for any serious condition

Stop use and ask a doctor if
• you experience any of the following signs of stomach bleeding:
• feel faint
• have bloody or black stools
• vomit blood
• have stomach pain that does not get better
• an allergic reaction occurs. Seek medical help right away.
• any new symptoms appear
• ringing in the ears or loss of hearing occurs
• pain gets worse or lasts for more than 10 days
• painful area is red or swollen
• fever gets worse or lasts for more than 3 days

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
• do not use more than directed
• drink a full glass of water with each dose
• adults and children 12 years and over
• take 2 tablets every 6 hours
• do not take more than 8 tablets in 24 hours
• children under 12 years: ask a doctor

Other information
• store at 25°C (77°F);

EDEN PRAIRIE, MN 55344 USA 1-877-932-7948
**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

No Print Area
Lot no. & Exp. Date

0 41163 48840 3

DISTRIBUTED BY SUPERVALU INC.
EDEN PRAIRIE, MN 55344 USA 1-877-932-7948
**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

Drug Facts (continued)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
• take more or for a longer time than directed • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug
• are age 60 or older • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product
Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use
• if you have ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if
• you have asthma
• you have a history of stomach problems, such as heartburn • you are taking a diuretic • you have high blood pressure, heart disease, liver cirrhosis or kidney disease
• stomach bleeding warning applies to you • you have liver disease

Ask a doctor or pharmacist before use if you are
• taking a prescription drug for diabetes, gout, or arthritis

**KEEP OUTER PACKAGE FOR
COMPLETE PRODUCT INFORMATION**

Drug Facts (continued)
taking any other drug or are under a doctor's care for any serious condition

Stop use and ask a doctor if
• you experience any of the following signs of stomach bleeding:
• feel faint
• have bloody or black stools
• vomit blood
• have stomach pain that does not get better
• an allergic reaction occurs. Seek medical help right away.
• any new symptoms appear
• ringing in the ears or loss of hearing occurs
• pain gets worse or lasts for more than 10 days
• painful area is red or swollen
• fever gets worse or lasts for more than 3 days

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
• do not use more than directed
• drink a full glass of water with each dose
• adults and children 12 years and over
• take 2 tablets every 6 hours
• do not take more than 8 tablets in 24 hours
• children under 12 years: ask a doctor

Other information
• store at 25°C (77°F);

EDEN PRAIRIE, MN 55344 USA 1-877-932-7948
**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

No Print Area
Lot no. & Exp. Date

0 41163 48840 3

DISTRIBUTED BY SUPERVALU INC.
EDEN PRAIRIE, MN 55344 USA 1-877-932-7948
**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

Drug Facts (continued)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
• take more or for a longer time than directed • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug
• are age 60 or older • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product
Caffeine warning: The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use
• if you have ever had an allergic reaction to acetaminophen, aspirin, or any other pain reliever/fever reducer • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if
• you have asthma
• you have a history of stomach problems, such as heartburn • you are taking a diuretic • you have high blood pressure, heart disease, liver cirrhosis or kidney disease
• stomach bleeding warning applies to you • you have liver disease

Ask a doctor or pharmacist before use if you are
• taking a prescription drug for diabetes, gout, or arthritis

Equaline 44-159B

HEADACHE RELIEF EXTRA STRENGTH

acetaminophen, aspirin, caffeine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41163-159
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
CROSPVIDONE (UNII: 2S7830E561)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POVIDONE (UNII: FZ989GH94E)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	

Product Characteristics

Color	WHITE	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	44;159
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41163-159-08	1 in 1 CARTON	11/17/1992	
1		24 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41163-159-12	1 in 1 CARTON	11/17/1992	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part343	11/17/1992	

Labeler - SUPERVALU INC. (006961411)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(41163-159)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(41163-159)

Revised: 4/2017

SUPERVALU INC.