

4246 FIRST AID KIT - 4246 first aid kit
Honeywell Safety Products USA, INC

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

4246 First Aid Kit (Burn Jel, alcohol wipes, HC cr, Burn Relief pump, PVP wipes, EW, ASA, amm. Inh, Foille Burn, Aypanal - 340420FP)

Eyewash

Active ingredient

Sterile Water 99%

Eyewash

Purpose

Eyewash

Eyewash

Uses

- for flushing the eye to remove loose foreign material, air pollutants or chlorinated water

Eyewash

Warnings

For external use only Obtain immediate medical treatment for all open wounds in or near eyes. To avoid contamination, do not touch tip of container to any surface. Do not reuse. Once opened, discard.

Do not use

- if solution changes color or becomes cloudy
- if you have open wounds in or near the eyes, get medical help right away.

Stop use and ask a doctor if

- you experience eye pain
- changes in vision
- continued redness or irritation of the eye
- condition worsens or persists

Keep out of reach of children

- If swallowed, get medical help or contact a Poison Control Center right away.

Eyewash

Directions

- remove contacts before using
- twist top to remove
- flush the affected area as needed
- control rate of flow by pressure on the bottle
- if necessary, continue flushing with emergency eyewash or shower

Eyewash

Inactive ingredients

sodium chloride, sodium phosphate dibasic, sodium phosphate monobasic

Eyewash

Questions

1-800-430-5490

Aspirin

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)* *nonsteroidal anti-inflammatory drug

Aspirin

Purpose

Pain reliever/fever reducer

Aspirin

Uses

temporarily reduces fever and relieves minor aches and pains associated with:

- a cold
- headache
- toothache
- muscular aches
- backache
- minor pain of arthritis
- premenstrual and menstrual periods

Aspirin

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:are:

- age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- ringing in the ears or loss of hearing occurs
- any new symptoms appear

If pregnant or breast-feeding,

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last three months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

- In case of overdose, get medical help or contact Poison Control Center right away.

Aspirin**Directions**

- drink a full glass of water with each dose
- adults and children 12 years of age and older: take 1 or 2 tablets every 4 hours while symptoms last, not more than 12 tablets in 24 hours
- children under 12 years of age: consult a doctor

Aspirin**Other information**

- store at room temperature 15° - 30°C (59° - 86°F)
- TAMPER EVIDENT PACKETS
- DO NOT USE IF OPEN OR TORN

Aspirin***Inactive ingredients***

corn starch, croscarmellose sodium*, hypromellose*, microcrystalline cellulose*, mineral oil*, polyethylene glycol*, povidone, propylene glycol, silicon dioxide, stearic acid*, titanium dioxide*

*may contain these ingredients

Aspirin***Questions or Comments***

1-800-430-5490

Ammonia***Active ingredient***

Ammonia 15%

Ammonia***Purpose***

Respiratory stimulant

Ammonia***Uses***

- to prevent or treat fainting

Ammonia***Warnings*****For external use only****Do not use**

- if you have breathing problems such as asthma or emphysema

Stop use and ask a doctor if

- condition persists

Keep out of reach of children

- If swallowed get medical help or contact a Poison Control Center right away.

Ammonia***Directions***

- hold inhalant away from face and crush ampoule between thumb and forefinger at position indicated on sleeve.
- hold near nostrils for inhalation of volatile vapor

Ammonia***Other information***

- store at room temperature away from light

Ammonia***Inactive ingredient***

alcohol USP, FD&C red #40, lavender oil, lemon oil fcc, nutmeg oil, purified water

Ammonia***Questions or Comments?***

1-800-430-5490

Burn Jel***Active ingredient***

Lidocaine HCl 2.0 %

Burn Jel***Purpose***

External analgesic

Burn Jel***Uses***

temporarily relieves pain due to minor burns

Stop use and ask a doctor if

- the condition gets worse
- symptoms persist for more than 7 days
- condition clears up and recurs within a few days

Burn Jel***Warnings*****For external use only****Do not use**

- on large areas of the body, particularly over raw surfaces or blistered areas

When using this product

- avoid contact with eyes

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Burn Jel***Directions***

- adults and children 2 years of age and older; apply to affected area not more than 3 to 4 times daily
- children under 2 years of age: ask a doctor
- you may report a serious reaction to this product to 800-430-5490

Burn Jel

Other information

store at room temperature

do not use if opened or torn

Burn Jel

Inactive ingredients

carbopol 940, carbopol 1342, diazolidinyl urea, glycerin, melaleuca alternifolia (tea tree) leaf oil, methylparaben, octoxynol-9, propylene glycol, propylparaben, trolamine, water

Burn Jel

Questions

1-800-430-5490

Alcohol Wipes

Active ingredient

Isopropyl alcohol 70%

Alcohol Wipes

Purpose

First aid antiseptic

Alcohol Wipes

Uses

first aid to help prevent infection in minor cuts, scrapes, and burns

Keep out of the reach of children

If swallowed, get medical help or contact a Poison Control center right away

Alcohol Wipes

Warnings

For external use only

Flammable, keep away from fire and flame

Do not use

- in or near eyes
- over large areas of the body

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

When using this product

- do not use longer than 1 week unless directed by a doctor

Stop use and consult a doctor if

- condition persists or gets worse

Alcohol Wipes

Directions

- clean the affected area
- may be covered with a sterile bandage
- apply wipe to affected area 1 to 3 times daily
- discard wipe after single use

Alcohol Wipes

Other information

- store at room temperature 15⁰ to 25⁰ C (59⁰ to 77⁰ F)
- do not use if packet is torn or opened

Alcohol Wipe

Inactive ingredient

water

Alcohol Wipe

Questions

1-800-430-5490

Hydrocortisone

Active ingredient (in each gram)

Hydrocortisone acetate (equivalent to Hydrocortisone 1%)

Hydrocortisone

Purpose

Anti-itch cream

Hydrocortisone

Uses

for the temporary relief of itching associated with minor skin irritations and rashes

Hydrocortisone

Warnings

For external use only

Ask a doctor before use if

- you are using any other hydrocortisone product

When using the product

- avoid contact with eyes
- do not begin use of any other hydrocortisone product unless you have consulted a doctor
- do not use for the treatment of diaper rash

Stop use and ask a doctor if

- condition worsens
- condition persists for more than 7 days
- condition clears up and recurs within a few days

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Hydrocortisone***Directions***

- **adults and children 2 years and older**
- clean the affected area
- apply to the area not more than 3 to 4 times daily
- children under 2 years of age: consult a doctor

Hydrocortisone***Other information***

- store at room temperature (do not freeze)

Hydrocortisone***Inactive ingredients***

cetyl alcohol, citric acid, diazolidinyl urea, edetate disodium, glycerin, glyceryl monostearate, methylparaben, mineral oil, polyethylene glycol, propylene glycol, propylparaben, purified water, stearic acid, trolamine

Hydrocortisone***Questions or Comments?***

1-800-430-5490

PVP Wipe***Active ingredient***

Povidone-iodine solution USP, 10% (equivalent to 1% titratable iodine)

PVP Wipe***Purpose***

First aid antiseptic

PVP Wipe***Uses***

- first aid to help prevent the risk of infection in minor cuts, scrapes, and burns

Stop use and ask a doctor if

- conditions persists or gets worse
- irritation and redness develops

PVP Wipe

Warnings

For external use only

Do not use

- in the eyes
- over large areas of the body
- on individuals who are allergic or sensitive to iodine

Ask a doctor before use if you have

- deep or puncture wounds,
- animal bites
- serious burns

When using this product

- do not use longer than one week unless directed by a doctor

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away

PVP Wipe

Directions

Reverse cardboard sleeve, then crush at dot between thumb and forefinger. Allow solution to saturate tip and apply solution to injury.

- clean affected area
- apply to affected area 1 to 3 times daily
- may be covered with a sterile bandage
- discard swab after single use

PVP

Other information

- store at room temperature away from light
- keep from freezing or excessive heat
- do not use if package is torn or open

PVP

Inactive ingredients

citric acid, disodium phosphate, nonoxynol-9, sodium hydroxide, water

PVP

Questions and Comments?

1-800-430-5490

Burn Relief

Active ingredient

Lidocaine HCL 2%

Burn Relief Spray

Purpose

External analgesic

Burn Spray

Uses

temporarily relieves pain due to minor burns

Burn Relief Spray

Warnings

For external use only

Do not use

- over large areas of the body, particularly over raw surfaces or blistered areas

When using this product

- avoid contact with the eyes

Stop use and ask a doctor if

- condition worsens
- symptoms persist for more than 7 days
- condition clears up and occurs again within a few days

Keep out of the reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Burn Relief

Directions

adults and children 2 years of age and older: apply to affected area not more than 3 to 4 times daily

children under 2 years of age: ask a doctor

Burn Relief Spray

Other information

- store at room temperature

Burn Reelief Spray

Inactive ingredients

diazolidinyl urea, edetate disodium, glycerin, hypromellose, methylparaben, octoxynol 9, propylene glycol, propylparaben, purified water, tea tree oil, trolamine

Burn Relief Spray

Questions or Comments?

1-800-430-5490

Aypanal

Active ingredient

Acetaminophen 325 mg

Aypanal

Purpose

Pain reliever/fever reducer

Aypanal

Uses

- temporarily relieves minor aches and pains due to the common cold and headache
- temporarily reduces fever

Aypanal

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin rash occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

- liver disease

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin

Stop using and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

If pregnant or breast-feeding

- ask a health professional before use.

Keep out of reach of children

Keep out of reach of children.

Overdose warning: In case of accidental overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Aypanal

Directions

- **do not take more than directed (see overdose warning)**
- adults and children 12 years of age or older
- take two tablets every 4-6 hours while symptoms last
- do not take more than 12 tablets in 24 hours
- children 6 to under 12 years of age
- take 1 tablet every 4-6 hours while symptoms last
- do not take more than 5 tablets in 24 hours
- children under 6 years
- consult a doctor

Aypanal

Other information

- store at room temperature 15⁰ to 30⁰ C (59⁰ - 86⁰ F)
- TAMPER EVIDENT PACKETS- DO NOT USE IF OPEN OR TORN

Aypanal

Inactive ingredients

corn starch, microcrystalline cellulose, povidone, sodium starch glycolate, stearic acid

Aypanal

Questions or Comments?

1-80-430-5490

Foille Ointment

Active ingredient

Benzocaine 5.0% (w/w)
Chloroxylenol 0.1% (w/w)

Foille Ointment

Purpose

External analgesic

Antiseptic

Foille Ointment

Uses

- For the temporary relief of pain associated with burns, sunburn, minor cuts, scrapes, insect bites, and minor skin irritations.
- First aid to help prevent infection in minor cuts, scrapes and burns.

Foille Ointment

Warnings

For external use only.

When using this product

- avoid contact with the eyes

Stop use and ask a doctor if

- condition worsens, or symptoms persist for more than 7 days or clear up and occur again within a few days.
- do not apply over large areas of the body. In case of deep or puncture wounds, animal bites, or serious burns, consult a doctor.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Foille Ointment

Directions

- clean the affected area.
- adults and children 2 years of age and older: Apply to affected area not more than 3 to 4 times daily.
- children under 2 years of age: consult a physician.

Foille Ointment

Other information

- avoid contact with clothing. Foille may stain certain fabrics.

Foille Ointment

Inactive ingredients

beeswax, benzyl alcohol, calcium disodium EDTA, calcium hydroxide, ceresin, eugenol, hydrogenated vegetable oil, maleic anhydride, mono- and di-glycerides, PEG-32, purified water, sodium borate, sodium lauryl sulfate, zea mays (corn) oil.

4246

68100REC Kit Contents

1 3/4 X 3 PLAS 100/BOX

1 1X3 PLASTIC 100/BOX

1 KING SIZE 2 X 3 PLASTIC 50/BX

1 AMMONIA INHALANTS 10 PER

1 GAUZE COMPRESS, 1728 SQ IN 1

4 INSTANT COLD PACK 4" X 6"

3 BANDAGE COMP, 4" OFFSET, 1 PER

1 BURN JEL 1/8 OZ, 6 PER

1 ALCOHOL PREP PADS 10P
1 HYDROCORTISON,1.0%,1/32 OZ,10P
2 ADHESIVE TAPE W/P 1/2"X 5 YD
2 O/H PUMP BURN RELIEF 2 OZ ID G
2 GAUZE BANDAGE 1" x 2 YDS
1 FIRST AID GUIDE ASHI
1 EMERGENCY SURVIVAL BLANKET
6 TAPE ADHESIVE 1"X 5 YD PLSTC
2 GZE PADS STERILE 2"X 2" 10'S
2 GZE PADS STERILE 3"X 3" 10'S
2 GZE PADS STERILE 2"X 2" 25'S
5 CO-FLEX BAND 1"X5YD TAN 2/PACK
2 CO-FLEX BANDAGE 4"X5YDS TAN
1 PVP PREP PADS MEDIUM 100/BX
1 ASPIRIN IND PK 5 GR 2/ENV 100
1 212 4 OZ TUBE SKIN COND, 1 EA ID P
2 1 OZ, BUFF EYEWASH
1 SCISSOR BDGE 4" RED PLS HDL
1 KIT TWEEZER 3 1/2" SLANTED
1 180 EMPTY BLANK NO LOGO
1 TONGUE BLADES SR WRAPPED 6'S
1 LBL STOCK 6-3/8"X4"
1 LBL STOCK 4"X2-7/8"
1 LBL STOCK 3"x1-7/8"
2 PR LRG NITRILE GLVES ZIP BAG
2 SELF-ADH WRAP 3 X 5 YDS NORTH REV E
2 TRI BNDG NON WOVEN 40"X40"X56"
2 FOILLE BURN/F A OINT 1/2 OZ
8 EYE PADS STD OVAL STERILE
5 GAUZE PADS 4"X4" 12PLY
1 WOVEN KNUCKLE 8'S
1 FINGERTIP "T" 8/BX
1 TRNQT W/WEB STP&BKL 1
1 AYPANAL NON-ASP 25/2

Honeywell

TAMPER-EVIDENT CAP.
TAPA CON SELLO DE SEGURIDAD.
BOUCHON INDICATEUR D'INFRACTION.

eyesaline®

LAVAOJOS
EYESALINE

Solución
Isotónico Estéril

EYESALINE
EYEWASH

Sterile
Isotonic Solution

LAVAGE
OCULAIRE
EYESALINE

La Solution
Isotonique Stérile

16 fl. oz. (473 mL)

NPN: 80057528
64809 1 45033 17

Drug Facts (for USA only)

Active ingredient Sterile water 99%
Purpose Eyewash
Uses for flushing the eye to remove loose foreign material, air pollutants, or chlorinated water.
Warnings For external use only - Obtain immediate medical treatment for all open wounds in or near the eyes. To avoid contamination, do not touch tip of container to any surface. Do not reuse. Once opened, discard.
Do not use if solution changes color or becomes cloudy
if you have open wounds in or near the eyes, get medical help right away
Stop use and consult a doctor if:
• you experience eye pain • changes in vision
• continued redness or irritation of the eye
• condition worsens or persists
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.
Directions
• remove contacts before using • twist top to remove
• flush the affected area as needed
• control rate of flow by pressure on the bottle
• if necessary, continue flushing with emergency eyewash or shower
Inactive ingredients sodium chloride, sodium phosphate dibasic, sodium phosphate monobasic
Questions? Call 1-800-430-5490
Honeywell Safety Products USA, Inc. Smithfield, RI. 02917

LABEL #32-0045/0 Rev. J REORDER / NUEVO PEDIDO / REAPPROVISIONNEMENT #32-000454-0000

space for lot code and supplier part number

PEEL / PELAR / PELEL


Datos de medicamento (Para EE.UU. solamente)

Ingrediente Activo Agua estéril 99%
Propósito Lavaojos
Usos para el lavado de ojo para quitar las partículas sueltas y extrañas, los contaminantes aéreos, o agua de cloruro.
Advertencias Para el uso externo sólo - Obtenga tratamiento médico inmediato para todas las heridas abiertas en o cerca de los ojos. Para evitar la contaminación, no toque la punta del envase con ninguna superficie. No vuelva a usar. Vez abierto, descarte.
No se use si la solución se enturbia o cambia de color
si tiene heridas abiertas en o cerca del ojo, obtenga ayuda médica de inmediato
Deje de usar y consulte a un médico si:
• experimenta dolor de ojo • cambio de visión
• rojez continuo o irritación del ojo
• la condición empeora o persiste
Manténgase fuera del alcance de los niños. En caso de ingestión accidental, obtenga atención médica o llame a un Centro de control de envenenamiento inmediatamente.
Instrucciones
• quite los lentes de contacto antes de usar la solución
• tuerza la tapa para quitar
• enjuague el área afectada según se necesite
• controle el chorro haciendo presión en la botella
• si es necesario, sigue enjuagado con un lavaojos o ducha de emergencia
Ingredientes inactivos cloruro de sodio, fosfato de sodio dibásico, fosfato de sodio monobásico
¿Preguntas? Llame al 1-800-430-5490
Honeywell Safety Products USA, Inc. Smithfield, RI. 02917

Information

Usages Pour le rinçage des yeux afin d'enlever un corps étranger, des polluants atmosphériques ou de l'eau chlorée.
Advertissements Pour usage externe seulement - Obtenir immédiatement des soins médicaux pour toutes les plaies ouvertes dans ou près des yeux. Pour éviter toute contamination, ne pas toucher la pointe du récipient à n'importe quelle surface. Ne pas réutiliser. Une fois ouvert, jeter.
Ne pas utiliser si la solution a changé de couleur ou si elle est brouillée
si vous avez des plaies ouvertes aux yeux ou à proximité, consultez immédiatement un médecin
Cesser d'utiliser la solution et consulter un médecin si vous ressentez une douleur oculaire si votre vision change
rougeur ou irritation persistante des yeux
condition empire ou persiste
Garder hors de la portée des enfants. En cas d'ingestion, communiquer immédiatement avec un médecin ou avec un centre antipoison.
Mode d'emploi enlever les verres de contact avant l'utilisation dévisser le bouchon pour l'enlever rincer la zone touchée selon les besoins ajuster le débit d'écoulement de la solution en augmentant ou en réduisant la pression exercée sur le contenant si nécessaire, continuer de rincer avec une solution de rinçage oculaire d'urgence ou une douche
Ingédients eau stérile, chlorure de sodium, phosphate dibasique de sodium, phosphate monobasique de sodium
Des questions? Faites le 1-800-430-5490
Honeywell Safety Products USA, Inc. Smithfield, RI. 02917

Aspirin
Principal Display Panel



7722-01 Rev C

Drug Facts (continued)
Active ingredient (in each tablet): Aspirin 325mg (NSAID)
Purpose: Pain reliever / fever reducer
Nonsteroidal anti-inflammatory drug
Uses: temporarily reduces fever and relieves minor aches and pains associated with:
 - cold - headache - toothache - menstrual aches - backache - minor pain of arthritis
Warnings:
 - aspirin allergy: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, changes in behavior with nausea and vomiting occur. Consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.
Warnings (continued):
 - aspirin allergy: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, changes in behavior with nausea and vomiting occur. Consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.



6 96351 53168 2
 Distributed by
 Honeywell Safety
 Products, USA, Inc.
 Smithfield, RI 02917

ASPIRIN 325 MG
Honeywell
Pain Reliever / Fever Reducer
(NSAID)
25 Packets (2 Tablets per Packet)
Total quantity 50 Tablets
PACKAGE NOT CHILD-RESISTANT
 35222AA

ASPIRIN 325 MG
25 Packets (2 Tablets per Packet)
Total quantity 50 Tablets
 35222AA

Drug Facts (continued)
Allergy alert: Aspirin may cause a severe allergic reaction which may include:
 • hives • facial swelling • asthma (wheezing) • shock
Stomach bleeding warning: This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you:
 • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed
Do not use if you are allergic to aspirin or any other pain reliever/fever reducer.
Ask a doctor before use if: - stomach bleeding warning applies to you - you have a history of stomach problems such as heartburn - you have high blood pressure, heart disease, liver cirrhosis, or kidney disease - you are taking a diuretic - you have asthma
Ask a doctor or pharmacist before use if you are: taking a prescription drug for diabetes, gout or arthritis
Stop use and ask a doctor if: - you experience any of the following signs of stomach bleeding:
 • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better
 • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days
 • redness or swelling is present in the painful area • ringing in the ears or loss of hearing occurs
 • any new symptoms appear
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center immediately.



Drug Facts (continued)
Directions - drink a full glass of water with each dose
 • adults and children 12 years of age and older: take 1 - 2 tablets every 4 hours while symptoms last, not more than 12 tablets in 24 hours
 • children under 12 years of age: consult a doctor
Other information - Store at room temperature 15° - 30° C (59° - 86° F)
 • **KEEPER EVIDENCE PACKETS - DO NOT USE IF OPEN OR TORN**
Inactive ingredients - corn starch, croscarmellose sodium, hypromellose, microcrystalline cellulose, mineral oil, polyethylene glycol, povidone, propylene glycol, silicon dioxide, stearic acid, titanium dioxide
 *may contain these ingredients
Questions or comments? 1-800-430-5490

ASPIRIN 325 MG
25 Packets (2 Tablets per Packet)
Total quantity 50 Tablets

7722-01 Rev C.pdf 1 9/5/2018 2:18:46 PM



Ammonia
Principal Display Panel

796006 Rev. E Unit Carton Printing Plate for "A" size carton.




796006 Rev. E (page 3 of 3)


Principal Display Panel



**Alcohol Wipe
Principal Display Panel**





**Hydrocortisone
Principal Display Panel**



PVP
Principal Display Panel

796042 Rev. B Unit Carton Printing Plate for "A" size carton.



Principal Display Panel

032204RH.pdf 1 8/27/2018 3:57:29 PM

032204 Rev. H North Burn Relief Spray
Two Color: Red PMS 032 & Black

	Honeywell 032204  BURN RELIEF SPRAY Lidocaine HCl <i>External analgesic</i> 2 FL OZ (59 mL) Package Not Child Resistant ANSI Z308.1-2003 	Drug Facts <table border="1"> <tr> <td>Active ingredient</td> <td>Purpose</td> </tr> <tr> <td>Lidocaine HCl 2%</td> <td>External analgesic</td> </tr> </table> Use temporarily relieves pain due to minor burns Warnings For external use only Do not use over large areas of the body, particularly over raw surfaces or blistered areas. When using this product avoid contact with the eyes Stop use and ask a doctor if <ul style="list-style-type: none"> condition worsens symptoms last more than 7 days condition clears up and occurs again within a few days Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. Directions <ul style="list-style-type: none"> adults and children 2 years of age and older: apply to affected area not more than 3 to 4 times daily children under 2 years of age: ask a doctor Other information store at room temperature Inactive ingredients diazolidinyl urea, edetate disodium, glycerin, hypromellose, methylparaben, octoxynol 9, propylene glycol, propylparaben, purified water, tea tree oil, triolamine Questions or comments? 1-800-430-5490	Active ingredient	Purpose	Lidocaine HCl 2%	External analgesic
	Active ingredient	Purpose				
Lidocaine HCl 2%	External analgesic					
	8 21812 01499 9 Mfg. for: Honeywell Safety Products USA, Inc. Smithfield, RI 02917 032204 Rev. H					

032204K Rev. F (Page 3 of 3)

Principal Display Panel

AYPANAL
25 Packets (2 Tablets Per Packet)
Quantity 50 Tablets

Questions or comments? 1-800-430-5490

Inactive ingredients corn starch, microcrystalline cellulose, polydene, sodium starch glycolate, stearic acid

Other information - store at room temperature 15-30°C (59°-86°F)
- TABLETS: DO NOT USE IF OPEN OR TORN

Drug Facts (continued)

children under 6 years	consult a doctor
12 years and older	do not take more than 5 tablets in 24 hours
adults and children	do not take more than 12 tablets in 24 hours

Directions - do not take more than directed (see overdose warning)

Overdose warning In case of accidental overdose, get medical help or contact a poison control center right away. From a medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Keep out of reach of children

If germs or yeast - feeling ask a health professional before use.

- fever gets worse or lasts more than 3 days - redness or swelling is present - new symptoms occur

- pain gets worse or lasts more than 5 days in children under 12 years

- pain gets worse or lasts more than 10 days in adults

Stop and ask a doctor if - pain gets worse or lasts more than 10 days in adults

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Do not use with any other drug containing acetaminophen (prescription or nonprescription).

Warnings - rash - skin redness - blisters - pain - if a skin reaction occurs, stop use and seek medical help right away.

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

Drug Facts (continued)

Honeywell

AYPANAL
25 Packets (2 Tablets Per Packet)
Quantity 50 Tablets

Acetaminophen 325 mg
Pain Reliever-Fever Reducer

25 Packets (2 Tablets Per Packet)
Quantity 50 Tablets

PACKAGE NOT CHILD-RESISTANT

UNIT NO. 35225AP

Distributed by
Honeywell Safety
Products USA, Inc.
Smithfield, RI 02917



6 69635 35170 5

Drug Facts

Active ingredient (in each tablet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer

Uses - temporarily relieves minor aches and pains due to the common cold and headache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg acetaminophen in 24 hours, which is the maximum daily amount
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adults has 3 or more alcoholic drinks every day while using this product

77225-01 Rev.B

Foille Ointment Principal Display Panel

Drug Facts

Active ingredients Purpose

Benzocaine 5.0% (w/w) External Analgesic
Chloroxylenol 0.1% (w/w) Antiseptic

Uses

- For the temporary relief of pain associated with burns, sunburn, minor cuts, scrapes, insect bites, and minor skin irritations.
- First aid to help prevent infection in minor cuts, scrapes and burns.

Warnings

- For external use only.
- When using this product:
 - Avoid contact with the eyes.
- **Stop use and ask a doctor if**
 - condition worsens, or symptoms persist for more than 7 days or clear up and occur again within a few days.
 - Do not apply over large areas of the body. In case of deep or puncture wounds, animal bites, or serious burns, consult a doctor.

Keep out of reach of children.
If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- Clean the affected area.
- Adults and children 2 years of age and older: Apply to affected area not more than 3 to 4 times daily.
- Children under 2 years of age: consult a physician.



MEDICATED FIRST AID OINTMENT

Fast, Soothing Relief Of Pain Due To:
Cuts & Scrapes • Minor Burns
Sunburn • Insect Bites

NDC 10157-9302-4



MEDICATED FIRST AID OINTMENT

Fast, Soothing Relief Of Pain Due To:
Cuts & Scrapes • Minor Burns
Sunburn • Insect Bites

NET WT.
1 oz (28g)

Drug Facts (continued)

Other information

- Avoid contact with clothing. Foille may stain certain fabrics.

Inactive ingredients

beeswax, benzyl alcohol, calcium disodium EDTA, calcium hydroxide, cerasin, eugenol, hydrogenated vegetable oil, maleic anhydrid mono- and di-glycerides, PEG-3-2, purified water, sodium borate, sodium lauryl sulfate, zea mays (corn) oil.



For the temporary relief of pain due to scrapes, cuts, minor burns, sunburn, non-poisonous insect bites, and minor skin irritation. Foille First Aid Ointment stops pain on contact and lets you resume normal activities right away. Foille has a medicated oil-based formula that helps heal and prevent infection.

SATISFACTION GUARANTEED
Blistex

©2009 Blistex Inc.
P.O. Box 5392
Oak Brook, IL 60522-5392
#39013

♻️ Carton is 100% Recyclable.

4246 Kit Label
68100REC

FIRST AID KIT



Louisville, Kentucky
(800) 428-3254

Distributed by Honeywell Safety Products USA, Inc. Smithfield, RI 02917

4246 FIRST AID KIT

4246 first aid kit kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0498-4246
--------------	----------------	--------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-4246-01	1 in 1 KIT; Type 0: Not a Combination Product	10/18/2018	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	2 TUBE	28 g
Part 2	10 PACKET	9 g
Part 3	2 BOTTLE	60 mL
Part 4	50 PACKET	100
Part 5	10 AMPULE	3 mL
Part 6	2 BOTTLE, SPRAY	118 mL
Part 7	10 POUCH	4 mL
Part 8	100 POUCH	30 mL
Part 9	6 PACKET	21 g
Part 10	10 PACKET	9 g
Part 11	25 PACKET	50

Part 1 of 11

BLISTEX FOILLE MEDICATED FIRST AID

benzocaine and chloroxylenol ointment

Product Information

Item Code (Source)	NDC:10157-9302
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLOROXYLENOL (UNII: 0F32U78V2Q) (CHLOROXYLENOL - UNII:0F32U78V2Q)	CHLOROXYLENOL	0.1 g in 100 g
BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	5 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
YELLOW WAX (UNII: 2ZA36H0S2V)	
CALCIUM HYDROXIDE (UNII: PF5DZW74VN)	
CERESIN (UNII: Q1LS2UJO3A)	
POLYETHYLENE GLYCOL 1500 (UNII: 1212Z7S33A)	
SODIUM BORATE (UNII: 91MBZ8H3QO)	
BENZYL ALCOHOL (UNII: LKG8494WBH)	
WATER (UNII: 059QF0KO0R)	

SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
CORN OIL (UNII: 8470G57WFM)	
EDETATE CALCIUM DISODIUM ANHYDROUS (UNII: 8U5D034955)	
EUGENOL (UNII: 3T8H1794QW)	
MALEIC ANHYDRIDE (UNII: V5877ZJZ25)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		14 g in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part348	03/05/2013	

Part 2 of 11

HYDROCORTISONE

anti-itch cream

Product Information

Item Code (Source)	NDC:0498-0801
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROCORTISONE ACETATE (UNII: 3X7931PO74) (HYDROCORTISONE - UNII:W14X0X7BPJ)	HYDROCORTISONE ACETATE	1 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
EDETATE DISODIUM (UNII: 7FLD91C86K)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
TROLAMINE (UNII: 9O3K93S3TK)	
METHYL PARABEN (UNII: A218C7H9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
DIAZOLIDINYL UREA (UNII: H5RIZ3MPW4)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	

PROPYLPARABEN (UNII: Z8IX2SC1OH)

GLYCERIN (UNII: PDC6A3C0OX)

WATER (UNII: 059QF0KO0R)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0801-35	0.9 g in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part348	10/15/2019	

Part 3 of 11

EYESALINE EMERGENCY EYEWASH

purified water liquid

Product Information

Item Code (Source)	NDC:0498-0100
Route of Administration	OPHTHALMIC

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
WATER (UNII: 059QF0KO0R) (WATER - UNII:059QF0KO0R)	WATER	98.6 mL in 100 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM PHOSPHATE, MONOBASIC, MONOHYDRATE (UNII: 593YOG76RN)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM PHOSPHATE, DIBASIC (UNII: GR686LBA74)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0100-01	30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

Part 4 of 11

ASPIRIN

aspirin tablet

Product Information

Item Code (Source)	NDC:0498-0114
--------------------	---------------

Route of Administration	ORAL
-------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSE 2208 (100 MPA.S) (UNII: B1QE5P712K)	
MINERAL OIL (UNII: T5L8T28FGP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	FR21
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0114-01	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	09/18/2018	

Part 5 of 11

AMMONIA INHALENT

ammonia inhalent inhalant

Product Information

Item Code (Source)	NDC:0498-3334
Route of Administration	RESPIRATORY (INHALATION)

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AMMONIA (UNII: 5138Q19F1X) (AMMONIA - UNII:5138Q19F1X)	AMMONIA	0.045 g in 0.3 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-3334-00	0.3 mL in 1 AMPULE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

Part 6 of 11

BURN RELIEF

lidocaine hydrochloride spray

Product Information

Item Code (Source)	NDC:0498-0221
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	24.64 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
GLYCERIN (UNII: PDC6A3C0OX)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
TROLAMINE (UNII: 9O3K93S3TK)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
DIAZOLIDINYL UREA (UNII: H5RIZ3MPW4)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
OCTOXYNOL-9 (UNII: 7JPC6Y25QS)	
TEA TREE OIL (UNII: VIF565UC2G)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0221-59	59 mL in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC mono graph not final	part348	09/18/2018	

Part 7 of 11**ALCOHOL WIPE**

isopropyl alcohol swab

Product Information

Item Code (Source)	NDC:0498-0143
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	0.7 mL in 1 mL
--	-------------------	----------------

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0143-04	0.4 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	09/18/2018	

Part 8 of 11

PVP IODINE WIPE

povidone-iodine 10% swab

Product Information

Item Code (Source)	NDC:0498-0121
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POVIDONE-IODINE (UNII: 85H0HZU99M) (IODINE - UNII:9679TC07X4)	IODINE	10 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
NONOXYNOL-9 (UNII: 48Q180SH9T)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0121-00	0.3 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

Part 9 of 11

BURN JEL

gel for burns gel

Product Information

Item Code (Source)	NDC:0498-0203
Route of Administration	TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	2 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
TEA TREE OIL (UNII: VIF565UC2G)	
DIAZOLIDINYL UREA (UNII: H5RIZ3MPW4)	
TROLAMINE (UNII: 9O3K93S3TK)	
METHYL PARABEN (UNII: A2I8C7HI9T)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
CARBOMER HOMOPOLYMER TYPE C (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: 4Q93RCW27E)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYL PARABEN (UNII: Z8IX2SC1OH)	
OCTOXYNOL-9 (UNII: 7JPC6Y25QS)	
CARBOMER HOMOPOLYMER TYPE B (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: HHT01ZNK31)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0203-00	3.5 g in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part348	09/18/2018	

Part 10 of 11

HYDROCORTISONE

anti-itch cream ointment

Product Information

Item Code (Source) NDC:0498-0800

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROCORTISONE ACETATE (UNII: 3X7931PO74) (HYDROCORTISONE - UNII:W14X0X7BPJ)	HYDROCORTISONE ACETATE	1 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
EDETATE DISODIUM (UNII: 7FLD91C86K)	
CETYL ALCOHOL (UNII: 936JST6JCN)	
TROLAMINE (UNII: 9O3K93S3TK)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
DIAZOLIDINYL UREA (UNII: H5RIZ3MPW4)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
LIGHT MINERAL OIL (UNII: N6K5787QVP)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0800-35	0.9 g in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part348	03/06/2013	10/15/2019

Part 11 of 11

AYPANAL NON-ASPIRIN

acetaminophen tablet

Product Information

Item Code (Source) NDC:0498-2001

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE (UNII: FZ989GH94E)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	circle;U
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-2001-01	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	01/02/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		10/18/2018	

Labeler - Honeywell Safety Products USA, INC (079287321)

Establishment

Name	Address	ID/FEI	Business Operations
Blistex Inc.		005126354	manufacture(10 157-9302)

Establishment

Name	Address	ID/FEI	Business Operations
James Alexander		040756421	manufacture(0498-3334)

Establishment

Name	Address	ID/FEI	Business Operations
Honeywell Safety Products USA, INC		079287321	pack(0498-4246)

Establishment

Name	Address	ID/FEI	Business Operations
Ultra Seal Corporation		085752004	manufacture(0498-0114, 0498-2001)

Establishment

Name	Address	ID/FEI	Business Operations
Water-Jel Technologies		155522589	manufacture(0498-0221, 0498-0203, 0498-0800, 0498-0801)

Establishment

Name	Address	ID/FEI	Business Operations
Honeywell Safety Products USA, Inc.		167518617	manufacture(0498-0100)

Establishment

Name	Address	ID/FEI	Business Operations
Changzhou Mao kang Medical		421317073	manufacture(0498-0143)

Establishment

Name	Address	ID/FEI	Business Operations
Sion Medical Biotext		532775194	manufacture(0498-0121)