

METZ SANI-SHIELD IODINE BARRIER TEAT DIP- iodine solution
Surpass Chemical Company, Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Metz Sani-Shield Iodine Barrier Teat Dip

METZ SANI-SHIELD

AIDS IN CONTROLLING THE SPREAD OF BACTERIA THAT MAY CAUSE
MASTITIS PRODUCT #: 630850

DIRECTIONS:

POST-DIP: Immediately after milker is removed, dip the teats with undiluted SANI-SHIELD. Do not wipe. If the cow will be returned to a below freezing temperature environment, allow the dip to air dry before discharging her from the milking area.

When a cow is being dried off, dip the teats for 3 or 4 days after last milking.

BEFORE THE NEXT MILKING: Just prior to the next milking, remove the barrier film by pre-dipping with NCDS IODINE TEAT DIP following label directions for pre-dipping. Dry teats thoroughly with a clean single service paper towel before milking.

CAUTION:

Only use as directed. Do not take internally. Avoid contact with eyes, skin, clothing, and food. Avoid breathing vapors.

Always use full strength Teat Dip. Discard solution if it becomes faded or noticeably dirty. Do not use this dip to sanitize milking equipment or utensils.

Wash dip cups thoroughly before use. Do not return dip from the cups back to the product container. If teats are chapped or sore, consult your veterinarian before beginning or continuing to use this product.

PRECAUTIONARY STATEMENTS

Prevention: Do not breathe dust/fume/gas/mist/vapors/spray Wash face, hands and any exposed skin thoroughly after handling Do not eat, drink or smoke when using this product

Response: Get medical advice/attention if you feel unwell

Disposal: Dispose of contents/container to an approved waste disposal plant

FIRST AID:

IF SPLASHED IN EYES: Rinse immediately with plenty of water, also under the eyelids,

for at least 15 minutes. Get medical attention if irritation occurs.

IF ON SKIN (or hair): Wash off immediately with plenty of water for at least 15 minutes. Get medical attention if symptoms occur.

IF INHALED: Move to fresh air. If symptoms persist, call a physician.

IF SWALLOWED: Drink plenty of water. Do NOT induce vomiting. Never give anything by mouth to an unconscious person. Consult a physician if necessary.

INGREDIENTS:

ACTIVE: Iodine complex (Provides 1% titratable iodine. Equivalent to 10,000 ppm).

12% Emollients

KEEP OUT OF REACH OF CHILDREN!

Surpass Chemical Company, Inc.

Phone: (518) 434-8101

24 HOUR

EMERGENCY ASSISTANCE:

www.surpasschemical.com

Fax: (518) 434-

2798

CHEMTREC 1-800-424-9300

1254 Broadway Albany NY, 12204

Email: customerservice@surpasschemical.com

KEEP OUT OF REACH OF CHILDREN!

BEFORE USING THIS MATERIAL, READ AND UNDERSTAND THE CURRENT SAFETY DATA SHEET.



METZ SANI-SHIELD IODINE BARRIER TEAT DIP

KEEP OUT OF REACH OF CHILDREN!

PRODUCT #: 630850

AIDS IN CONTROLLING THE SPREAD OF BACTERIA THAT MAY CAUSE MASTITIS



NPE FREE

LOT NUMBER
NET CONTENTS

CAUTION

Only use as directed. Do not take internally. Avoid contact with eyes, skin, clothing, and food. Avoid breathing vapors. Always use full strength Teat Dip. Discard solution if it becomes faded or noticeably dirty. Do not use this dip to sanitize milking equipment or utensils.

Wash dip cups thoroughly before use. Do not return dip from the cups back to the product container. If teats are chapped or sore, consult your veterinarian before beginning or continuing to use this product.

INGREDIENTS

ACTIVE: Iodine complex (Provides 1% titratable iodine. Equivalent to 10,000 ppm).
12% Emollients

DIRECTIONS

POST-DIP: Immediately after milker is removed, dip the teats with undiluted SANI-SHIELD. Do not wipe. If the cow will be returned to a below freezing temperature environment, allow the dip to air dry before discharging her from the milking area. When a cow is being dried off, dip the teats for 3 or 4 days after last milking.

BEFORE NEXT MILKING: Just prior to the next milking, remove the barrier film by pre-dipping with NCDS IODINE TEAT DIP following label directions for pre-dipping. Dry teats thoroughly with a clean single service paper towel before milking.

FIRST AID

IF SPLASHED IN EYES: Rinse immediately with plenty of water, also under the eyelids, for at least 15 minutes. Get medical attention if irritation occurs.

IF ON SKIN (or hair): Wash off immediately with plenty of water for at least 15 minutes. Get medical attention if symptoms occur.

IF INHALED: Move to fresh air. If symptoms persist, call a physician.

IF SWALLOWED: Drink plenty of water. Do NOT induce vomiting. Never give anything by mouth to an unconscious person. Consult a physician if necessary.



Surpass Chemical Company, Inc.
www.surpasschemical.com
1254 Broadway Albany NY, 12204

Phone: (518) 434-8101

Email: customerservice@surpasschemical.com

Fax: (518) 434-2798

BEFORE USING THIS MATERIAL, READ AND UNDERSTAND THE CURRENT SAFETY DATA SHEET.

24 HOUR EMERGENCY ASSISTANCE:
CHEMTREC 1-800-424-9300



METZ SANI-SHIELD IODINE BARRIER TEAT DIP

KEEP OUT OF REACH OF CHILDREN!

PRODUCT #: 630850

AIDS IN CONTROLLING THE SPREAD OF BACTERIA THAT MAY CAUSE MASTITIS



NPE FREE

LOT NUMBER
NET CONTENTS

CAUTION

Only use as directed. Do not take internally. Avoid contact with eyes, skin, clothing, and food. Avoid breathing vapors. Always use full strength Teat Dip. Discard solution if it becomes faded or noticeably dirty. Do not use this dip to sanitize milking equipment or utensils.

Wash dip cups thoroughly before use. Do not return dip from the cups back to the product container. If teats are chapped or sore, consult your veterinarian before beginning or continuing to use this product.

INGREDIENTS

ACTIVE: Iodine complex (Provides 1% titratable iodine. Equivalent to 10,000 ppm).
12% Emollients

DIRECTIONS

POST-DIP: Immediately after milker is removed, dip the teats with undiluted SANI-SHIELD. Do not wipe. If the cow will be returned to a below freezing temperature environment, allow the dip to air dry before discharging her from the milking area. When a cow is being dried off, dip the teats for 3 or 4 days after last milking.

BEFORE NEXT MILKING: Just prior to the next milking, remove the barrier film by pre-dipping with NCDS IODINE TEAT DIP following label directions for pre-dipping. Dry teats thoroughly with a clean single service paper towel before milking.

FIRST AID

IF SPLASHED IN EYES: Rinse immediately with plenty of water, also under the eyelids, for at least 15 minutes. Get medical attention if irritation occurs.

IF ON SKIN (or hair): Wash off immediately with plenty of water for at least 15 minutes. Get medical attention if symptoms occur.

IF INHALED: Move to fresh air. If symptoms persist, call a physician.

IF SWALLOWED: Drink plenty of water. Do NOT induce vomiting. Never give anything by mouth to an unconscious person. Consult a physician if necessary.



Surpass Chemical Company, Inc.
www.surpasschemical.com
1254 Broadway Albany NY, 12204

Phone: (518) 434-8101

Email: customerservice@surpasschemical.com

Fax: (518) 434-2798

BEFORE USING THIS MATERIAL, READ AND UNDERSTAND THE CURRENT SAFETY DATA SHEET.

24 HOUR EMERGENCY ASSISTANCE:
CHEMTREC 1-800-424-9300



METZ SANI-SHIELD IODINE BARRIER TEAT DIP

AIDS IN CONTROLLING THE SPREAD OF BACTERIA THAT MAY CAUSE MASTITIS

CAUTION

Only use as directed. Do not take internally. Avoid contact with eyes, skin, clothing, and food. Avoid breathing vapors. Always use full strength Teat Dip. Discard solution if it becomes faded or noticeably dirty. Do not use this dip to sanitize milking equipment or utensils. Wash dip cups thoroughly before use. Do not return dip from the cups back to the product container. If teats are chapped or sore, consult your veterinarian before beginning or continuing to use this product.

INGREDIENTS

ACTIVE: Iodine complex (Provides 1% titratable iodine. Equivalent to 10,000 ppm).
12% Emollients

DIRECTIONS

POST-DIP: Immediately after milker is removed, dip the teats with undiluted SANI-SHIELD. Do not wipe. If the cow will be returned to a below freezing temperature environment, allow the dip to air dry before discharging her from the milking area. When a cow is being dried off, dip the teats for 3 or 4 days after last milking.

BEFORE NEXT MILKING: Just prior to the next milking, remove the barrier film by pre-dipping with NCDS IODINE TEAT DIP following label directions for pre-dipping. Dry teats thoroughly with a clean single service paper towel before milking.

FIRST AID

IF SPLASHED IN EYES: Rinse immediately with plenty of water, also under the eyelids, for at least 15 minutes. Get medical attention if irritation occurs.

IF ON SKIN (or hair): Wash off immediately with plenty of water for at least 15 minutes. Get medical attention if symptoms occur.

IF INHALED: Move to fresh air. If symptoms persist, call a physician.

IF SWALLOWED: Drink plenty of water. Do NOT induce vomiting. Never give anything by mouth to an unconscious person. Consult a physician if necessary.

KEEP OUT OF REACH OF CHILDREN!

NPE FREE

PRODUCT #: 630850



Surpass Chemical Company, Inc.
www.surpasschemical.com

1254 Broadway Albany NY, 12204

Phone: (518)434-8101

Email: customerservice@surpasschemical.com

Fax: (518) 434-2798

24 HOUR EMERGENCY ASSISTANCE:
CHEMTREC 1-800-424-9300

BEFORE USING THIS MATERIAL, READ AND UNDERSTAND THE CURRENT SAFETY DATA SHEET.

LOT

NUMBER:

NET

CONTENTS:

METZ SANI-SHIELD IODINE BARRIER TEAT DIP

iodine solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:86067-0122
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
iodine (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	iodine	0.04724 kg in 1 kg

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	0.87076 kg in 1 kg
C9-11 PARETH-8 (UNII: 80E6PSE1XL)	0.02 kg in 1 kg
ALOE VERA FLOWER (UNII: 575DY8C1ER)	0.0005 kg in 1 kg
GLYCERIN (UNII: PDC6A3C0OX)	0.05 kg in 1 kg
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	0.0025 kg in 1 kg
XANTHAN GUM (UNII: TTV12P4NEE)	0.003 kg in 1 kg
BUTYLENE GLYCOL (UNII: 3XUS85K0RA)	0.006 kg in 1 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:86067-0122-2	4 in 1 BOX		
1	NDC:86067-0122-1	3.87 kg in 1 JUG		
2	NDC:86067-0122-3	19.05 kg in 1 PAIL		
3	NDC:86067-0122-4	58.06 kg in 1 DRUM		
4	NDC:86067-0122-5	116.12 kg in 1 DRUM		
5	NDC:86067-0122-6	213.19 kg in 1 DRUM		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		05/15/2023	

Labeler - Surpass Chemical Company, Inc. (002075133)

Establishment

Name	Address	ID/FEI	Business Operations
Surpass Chemical Company, Inc.		002075133	manufacture, api manufacture

Revised: 5/2024

Surpass Chemical Company, Inc.