PLANTAGO MAJOR- plantago major pellet Boiron

Disclaimer: This homeopathic product has not been evaluated by the Food and Drug Administration for safety or efficacy. FDA is not aware of scientific evidence to support homeopathy as effective.

Plantago major 200CK

Plantago major 200CK

(**contains 0.443 mg of the active ingredient per pellet)

Children bedwetting or dental pains*

Stop use and ask a doctor if symptoms persist for more than 3 days or worsen

If pregnant or breast-feeding ask a health professional before use

Keep out of reach of children

Do not use if pellet dispenser seal is broken.

Contains approx 80 pellets.

How to dispense pellets? Turn tube upside down. Twist until 5 pellets are dispensed into cap. Carefully remove the cap and use it to pour pellets under the tongue.

*CLAIMS BASED ON TRADITIONAL HOMEOPATHIC PRACTICE NOT ACCEPTED MEDICAL EVIDENCE. NOT FDA EVALUATED.

*C,K,CK, and X are homeopathic dilutions: see BoironUSA.com/info for details.

lactose, sucrose

Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

1-800-BOIRON-1 (1-800-264-7661), BoironUSA.com Info@boiron.com Distributed by Boiron, Inc. Newtown Square, PA 19073



Children bedwetting or dental pains.*

Lot: Exp:

3 06960 58131 4

Contains approx. 80 pellets. US Peel for Drugs Facts and instructions for use.

NAL HOMEOPATHIC PRACTICE, DENCE. NOT FDA EVALUATED.

Drug Facts

Active ingredient**: See product name on front panel (contains 0.443 mg of the active ingredient per pellet).


Uses: See symptoms on front panel.

Warnings: Stop use and ask a doctor if symptoms persist for more than 3 days or worsen. If pregnant or breastfeeding, ask a health professional before use. Keep out of reach of children.

Directions: ■ Adults and children: At the onset of symptoms, dissolve 5 pellets under the tongue 3 times a day until symptoms are relieved or as directed by a doctor.

Other information: ■ Do not use if pellet dispenser seal is broken.

Drug Facts (continued) **Inactive ingredients:** lactose, sucrose



PLANTAGO MAJOR

plantago major pellet

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0220-4088	
Route of Administration	ORAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
PLANTAGO MAJOR (UNII: W2469WNO6U) (PLANTAGO MAJOR - UNII: W2469WNO6U)	PLANTAGO MAJOR	200 [kp_C] in 200 [kp_C]	

Inactive Ingredients		
Ingredient Name	Strength	
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)		
SUCROSE (UNII: C151H8M554)		

Product Characteristics			
Color	white	Score	
Shape	ROUND	Size	4mm

Flavor	Imprint Code	
Contains		

ı	Packaging				
	# Item Code	Package Description	Marketing Start Date	Marketing End Date	
	1 NDC:0220- 4088-41	200 [kp_C] in 1 TUBE; Type 0: Not a Combination Product	03/03/1983		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved homeopathic		03/03/1983	
homeopathic		03/03/1903	

Labeler - Boiron (282560473)

Registrant - Boiron, Inc. (014892269)

Establishment			
Name	Address	ID/FEI	Business Operations
Boiron		282560473	manufacture(0220-4088)

Revised: 11/2023 Boiron