MINOXIDIL- minoxidil aerosol, foam P & L Development, LLC

Minoxidil Topical Solution for Men

Drug Facts

Active ingredient

Minoxidil, USP 5% w/v (without propellant)

Purpose

Hair regrowth treatment for men

Use

to regrow hair on the top of the scalp (vertex only, see pictures on side of carton)

Warnings

For external use only. For use by men only.

Flammable: Avoid fire. flame, or smoking during and immediately following application

Do not use if

- you are a woman
- your amount of hair loss is different than that shown on the side of this carton or your hair loss is on the front of the scalp. Minoxidil topical Aerosol, 5% (Foam) is not intended for frontal baldness or receding hairline.
- you have no family history of hair loss
- your hair loss is sudden and/or patchy
- you do not know the reason for your hair loss
- you are under 18 years of age. Do not use on babies and children.
- your scalp is red, inflamed, infected, irritated, or painful
- you use other medicines on the scalp

Ask a doctor before use if you have

heart disease

When using this product

- do not apply on other parts of the body
- avoid contact with the eyes. In case of accidental contact, rinse eyes with large amounts of cool tap water.
- some people have experienced changes in hair color and/or texture
- it takes time to regrow hair. Results may occur at 2 months with twice a day usage.

For some men, you may need to use this product for at least 4 months before you see results.

• the amount of hair regrowth is different for each person. This product will not work for all men.

Stop use and ask a doctor if

- chest pain, rapid heartbeat, faintness, or dizziness occurs
- sudden, unexplained weight gain occurs
- your hands or feet swell
- scalp irritation or redness occurs
- unwanted facial hair growth occurs
- you do not see hair regrowth in 4 months

May be harmful if used when pregnant or breast-feeding.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- apply half a capful 2 times a day to the scalp in the hair loss area
- massage into scalp with fingers, then wash hands well
- see enclosed leaflet for complete directions on how to use
- using more or more often will not improve results
- continued use is necessary to increase and keep your hair regrowth, or hair loss will begin again

Other information

- hair growth has been shown in clinical study of men (mostly white) aged 18-49 years who used it for 4 month
- see hair loss pictures on side of this carton
- before use, read all information on carton and enclosed leaflet
- keep the carton. It contains important information.
- store at controlled room temperature 20° to 25°C (68° to 77°F)
- contents under pressure. Do not puncture or incinerate container.
- Do not expose to heat or store at temperatures above 120° F (49° C).

Inactive ingredients

butane, butylated hydroxytoluene, cetyl alcohol, citric acid glycerin, isobutane, lactic acid, polysorbate 60, propane, purified water, SD alcohol 40-B, stearyl alcohol

Questions?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal display panel

*Compare to the active ingredient in Men's Rogaine®

men's hair regrowth treatment

Minoxidil Topical aerosol, 5%

hair regrowth treatment

- Clinically proven to help regrow hair
- reactivates hair follicles to stimulate regrowth
- unscented

easy-to-use foam

NOT FOR USE BY WOMEN

THREE MONTH SUPPLY

THREE 60 G (2.11 OZ) CANS


mL (FL OZ) BOTTLE

*This product is not manufactured by, or distributed by Johnson & Johnson Comsumer Inc., distributor of Men's Rogaine®

Distributed by: PL Developments

200 Hicks Street, Westbury, NY 11590

Package label



WELLNESS BASICS Men's Hair regrowth Treatment

MINOXIDIL

minoxidil aerosol, foam

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:59726-125

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name
Basis of Strength
MINOXIDIL (UNII: 5965120SH1) (MINOXIDIL - UNII:5965120SH1)
MINOXIDIL
50 mg in 1 mL

Inactive Ingredients

Ingredient Name
Strength

BUTANE (UNII: 6LV4FOR43R)

BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)

CETYL ALCOHOL (UNII: 936JST6JCN)

CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)

GLYCERIN (UNII: PDC6A3C0OX)

ISOBUTANE (UNII: BXR49TP611)

LACTIC ACID (UNII: 33X04XA5AT)	
POLYSORBATE 60 (UNII: CAL22UVI4M)	
PROPANE (UNII: T75W9911L6)	
WATER (UNII: 059QF0KO0R)	
ALCOHOL (UNII: 3K9958V90M)	
STEARYL ALCOHOL (UNII: 2KR89I4H1Y)	

Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:59726-125- 60	1 in 1 PACKAGE	10/01/2023		
1		60 mL in 1 CAN; Type 0: Not a Combination Product			
2	NDC:59726-125- 70	3 in 1 PACKAGE	10/01/2023		
2		60 mL in 1 CAN; Type 0: Not a Combination Product			

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
ANDA	ANDA208092	10/01/2023			

Labeler - P & L Development, LLC (800014821)

Revised: 10/2023 P & L Development, LLC