# VICKS VAPOCOOL SORE THROAT- benzocaine and menthol liquid The Procter & Gamble Manufacturing Company

-----

# VICKS VAPOCOOL SORE THROAT BENZOCAINE & MENTHOL - ORAL ANESTHETICS Spray

**Drug Facts** 

#### **Active ingredients**

Benzocaine (5 %) Menthol (1 %)

#### **Purpose**

Oral anesthetic

Oral anesthetic

#### Uses

temporarily relieves occasional minor irritation and pain due to

- sore throat
- sore mouth

#### **Warnings**

# Methemoglobinemia Warning-

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

## Allergy alert:

do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics. If a skin reaction occurs, stop use and seek medical help right away.

# Sore Throat Warning -

Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 12 years of age.

#### Do not use

- for teething
- in children under 2

#### When using this product

do not exceed recommended dosage

#### Ask a doctor before use if you have

- a severe sore throat accompanied by difficulty in breathing or that lasts more than 2 days
- a sore throat accompanied by fever, headache, rash, swelling, nausea or vomiting

#### Stop use and ask a doctor if

 sore mouth symptoms do not improve in 7 days, or if irritation, pain, or redness persists or worsens

## If pregnant or breast-feeding,

ask a health professional before use.

# Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- adults and children 12 years and over: Apply to the affected area (one spray). Gargle, swish around in the mouth, or allow to remain in place at least 1 minute and then spit out. Use up to 4 times daily or as directed by a doctor.
- children under 12 years, do not use.

#### Other information

• do not exceed 25°C. Do not refrigerate.

# Inactive ingredients

alcohol, D&C Yellow No. 10, FD&C Blue No. 1, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sucralose

#### **Ouestions?**

#### TAMPER EVIDENT: Do not use if printed shrinkband is missing or broken.

**Made in Canada** Distributed by **Procter & Gamble,** Cincinnati, OH 45202

#### PRINCIPAL DISPLAY PANEL - 177 ml Bottle Label

**VICKS**®

VapoC00L

#### **SORE THROAT**

BENZOCAINE & MENTHOL - ORAL ANESTHETICS

COOLING + NUMBING

Soothes Throat Pain Fast-acting

WINTERFROST

6 FL OZ (177 ml)



	monator (1 70) Ordi dilodilodo
	Uses temporarily relieves occasional minor irritation and pain due to • sore throat • sore mouth
	Warnings Methemoglobinemia Warning – Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops: • pale, gray or blue colored skin (cyanosis • headache • rapid heart rate • shortness of breatf • dizziness or lightheadedness • fatigue or lack of energy
	Allergy alert: do not use this product if you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine, or other "caine" anesthetics. If a skin reaction occurs, stop use and seek medical help right away.
by high fever, headache, naus	re or persistent sore throat or sore throat accompanied ea, and vomiting may be serious. Consult doctor an 2 days or administer to children under 12 years of age
Do not use • for teething	in children under 2
When using this product do	not exceed recommended dosage
difficulty in breathing or that I	ou have • a severe sore throat accompanied by asts more than 2 days by fever, headache, rash, swelling, nausea or vomiting
Stop use and ask a doctor i if irritation, pain, or redness p	<ul> <li>sore mouth symptoms do not improve in 7 days, or ersists or worsens</li> </ul>
reach of children.	g, ask a health professional before use. <b>Keep out of</b> cal help or contact a Poison Control Center right away
(one spray). Gargle, swish aro	d children 12 years and over: Apply to the affected area und in the mouth, or allow to remain in place at least se up to 4 times daily or as directed by a doctor. not use.
Other information •	do not exceed 25°C. Do not refrigerate.

**Purpose** 

**Drug Facts** Active ingredients

Benzocaine (5 %) Menthol (1 %) .....

#### benzocaine and menthol liquid

### **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:37000-928

Route of Administration ORAL

# Active Ingredient/Active Moiety Ingredient Name BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5) BENZOCAINE (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A) MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A) MENTHOL 1 g in 100 mL

Inactive Ingredients			
Ingredient Name	Strength		
GLYCERIN (UNII: PDC6A3C0OX)			
ALCOHOL (UNII: 3K9958V90M)			
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
WATER (UNII: 059QF0KO0R)			
SACCHARIN SODIUM (UNII: SB8ZUX40TY)			

Product Characteristics					
Color	turquoise	Score			
Shape		Size			
Flavor	MENTHOL	Imprint Code			
Contains					

Packaging				
# Item Code	Package Description	Marketing Start Date	Marketing End Date	
1 NDC:37000- 928-06	177 mL in 1 BOTTLE, PUMP; Type 0: Not a Combination Product	10/09/2019		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M022	10/09/2019		