

ALBUTEROL SULFATE HFA- albuterol sulfate aerosol, metered
H.J. Harkins Company, Inc.

1171 ALBUTEROL (PROAIR)

1.1 Bronchospasm

Albuterol Sulfate HFA Inhalation Aerosol is indicated for the treatment or prevention of bronchospasm in patients 4 years of age and older with reversible obstructive airway disease.

1.2 Exercise-Induced Bronchospasm

Albuterol Sulfate HFA Inhalation Aerosol is indicated for the prevention of exercise-induced bronchospasm in patients 4 years of age and older.

2.1 Bronchospasm

For treatment of acute episodes of bronchospasm or prevention of symptoms associated with bronchospasm, the usual dosage for adults and children 4 years and older is two inhalations repeated every 4 to 6 hours. More frequent administration or a larger number of inhalations is not recommended. In some patients, one inhalation every 4 hours may be sufficient.

2.2 Exercise-Induced Bronchospasm

The usual dosage for adults and children 4 years of age or older is two inhalations 15 to 30 minutes before exercise.

2.3 Administration Information

Administer Albuterol Sulfate HFA by oral inhalation only. Shake well before each spray. To maintain proper use of this product and to prevent medication build-up and blockage, it is important to follow the cleaning directions carefully.

Priming: Prime the inhaler before using for the first time and in cases where the inhaler has not been used for more than 2 weeks by releasing three sprays into the air, away from the face.

Cleaning: As with all HFA-containing albuterol inhalers, to maintain proper use of this product and to prevent medication build-up and blockage, it is important to clean the plastic mouthpiece regularly. The inhaler may cease to deliver medication if the plastic actuator mouthpiece is not properly cleaned and dried. To clean: Wash the plastic mouthpiece with warm running water for 30 seconds, shake off excess water, and air dry thoroughly at least once a week. If the patient has more than one Albuterol Sulfate HFA inhaler, the patient should wash each one separately to prevent attaching the wrong canister to the wrong plastic actuator. In this way, the patient can be sure to always know the correct number of remaining doses. Never attach a canister of medication from any other inhaler to the Albuterol Sulfate HFA actuator and never attach the Albuterol Sulfate HFA canister to an actuator from any other inhaler. If the mouthpiece becomes blocked, washing the mouthpiece will remove the blockage. If it is necessary to use the inhaler before it is completely dry, shake off excess water, replace canister, spray twice into the air away from face, and take the prescribed dose. After such use, the mouthpiece should be rewashed and allowed to air dry thoroughly. [see FDA-APPROVED PATIENT LABELING (17.9)].

Dose Counter: Albuterol Sulfate HFA has a dose counter attached to the actuator. When the patient receives the inhaler, a black dot will appear in the viewing window until it has been primed 3 times, at which point the number 200 will be displayed. The dose counter will count down each time a spray is released. When the dose counter reaches 20, the color of the numbers will change to red to remind the patient to contact their pharmacist for a refill of medication or consult their physician for a prescription refill. When the dose counter reaches 0, the background will change to solid red. Albuterol Sulfate HFA inhaler should be discarded when the dose counter displays 0 or after the expiration date on the product, whichever comes first.

Albuterol Sulfate HFA is an inhalation aerosol. Albuterol Sulfate HFA is supplied as an 8.5 g/200 actuations pressurized aluminum canister with a red plastic actuator with a dose counter and white dust cap each in boxes of one. Each actuation delivers 120 mcg of albuterol sulfate from the canister valve and 108 mcg of albuterol sulfate from the actuator mouthpiece (equivalent to 90 mcg of albuterol base).

Albuterol Sulfate HFA Inhalation Aerosol is contraindicated in patients with a history of hypersensitivity to albuterol and any other Albuterol Sulfate HFA Inhalation Aerosol components. Rare cases of hypersensitivity reactions, including urticaria, angioedema, and rash have been reported after the use of albuterol sulfate [see WARNINGS AND PRECAUTIONS (5.6)].

5.1 Paradoxical Bronchospasm

Albuterol Sulfate HFA Inhalation Aerosol can produce paradoxical bronchospasm that may be life threatening. If paradoxical bronchospasm occurs, Albuterol Sulfate HFA Inhalation Aerosol should be discontinued immediately and alternative therapy instituted. It should be recognized that paradoxical bronchospasm, when associated with inhaled formulations, frequently occurs with the first use of a new canister.

5.2 Deterioration of Asthma

Asthma may deteriorate acutely over a period of hours or chronically over several days or longer. If the patient needs more doses of Albuterol Sulfate HFA Inhalation Aerosol than usual, this may be a marker of destabilization of asthma and requires re-evaluation of the patient and treatment regimen, giving special consideration to the possible need for anti-inflammatory treatment, e.g., corticosteroids.

5.3 Use of Anti-inflammatory Agents

The use of beta-adrenergic-agonist bronchodilators alone may not be adequate to control asthma in many patients. Early consideration should be given to adding anti-inflammatory agents, e.g., corticosteroids, to the therapeutic regimen.

5.4 Cardiovascular Effects

Albuterol Sulfate HFA Inhalation Aerosol, like other beta-adrenergic agonists, can produce clinically significant cardiovascular effects in some patients as measured by pulse rate, blood pressure, and/or symptoms. Although such effects are uncommon after administration of Albuterol Sulfate HFA Inhalation Aerosol at recommended doses, if they occur, the drug may need to be discontinued. In addition, beta-agonists have been reported to produce ECG changes, such as flattening of the T wave, prolongation of the QTc interval, and ST segment depression. The clinical significance of these findings is unknown. Therefore, Albuterol Sulfate HFA Inhalation Aerosol, like all sympathomimetic amines, should be used with caution in patients with cardiovascular disorders, especially coronary insufficiency, cardiac arrhythmias, and hypertension.

5.5 Do Not Exceed Recommended Dose

Fatalities have been reported in association with excessive use of inhaled sympathomimetic drugs in patients with asthma. The exact cause of death is unknown, but cardiac arrest following an unexpected development of a severe acute asthmatic crisis and subsequent hypoxia is suspected.

5.6 Immediate Hypersensitivity Reactions

Immediate hypersensitivity reactions may occur after administration of albuterol sulfate, as demonstrated by rare cases of urticaria, angioedema, rash, bronchospasm, anaphylaxis, and oropharyngeal edema. The potential for hypersensitivity must be considered in the clinical evaluation of patients who experience immediate hypersensitivity reactions while receiving Albuterol Sulfate HFA Inhalation Aerosol.

5.7 Coexisting Conditions

Albuterol Sulfate HFA Inhalation Aerosol, like all sympathomimetic amines, should be used with caution in patients with cardiovascular disorders, especially coronary insufficiency, cardiac arrhythmias, and hypertension; in patients with convulsive disorders, hyperthyroidism, or diabetes

mellitus; and in patients who are unusually responsive to sympathomimetic amines. Clinically significant changes in systolic and diastolic blood pressure have been seen in individual patients and could be expected to occur in some patients after use of any beta-adrenergic bronchodilator. Large doses of intravenous albuterol have been reported to aggravate preexisting diabetes mellitus and ketoacidosis.

5.8 Hypokalemia

As with other beta-agonists, Albuterol Sulfate HFA Inhalation Aerosol may produce significant hypokalemia in some patients, possibly through intracellular shunting, which has the potential to produce adverse cardiovascular effects. The decrease is usually transient, not requiring supplementation.

Use of Albuterol Sulfate HFA may be associated with the following:

Paradoxical bronchospasm [see WARNINGS AND PRECAUTIONS (5.1)]

Cardiovascular Effects [see WARNINGS AND PRECAUTIONS (5.4)]

Immediate hypersensitivity reactions [see WARNINGS AND PRECAUTIONS (5.6)]

Hypokalemia [see WARNINGS AND PRECAUTIONS (5.8)]

6.1 Clinical Trials Experience

A total of 1090 subjects were treated with Albuterol Sulfate HFA Inhalation Aerosol, or with the same formulation of albuterol as in Albuterol Sulfate HFA Inhalation Aerosol, during the worldwide clinical development program.

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Adult and Adolescents 12 Years of Age and Older: The adverse reaction information presented in the table below concerning Albuterol Sulfate HFA Inhalation Aerosol is derived from a 6-week, blinded study which compared Albuterol Sulfate HFA Inhalation Aerosol (180 mcg four times daily) with a double-blinded matched placebo HFA-Inhalation Aerosol and an evaluator-blinded marketed active comparator HFA-134a albuterol inhaler in 172 asthmatic patients 12 to 76 years of age. The table lists the incidence of all adverse events (whether considered by the investigator drug related or unrelated to drug) from this study which occurred at a rate of 3% or greater in the Albuterol Sulfate HFA Inhalation Aerosol treatment group and more frequently in the Albuterol Sulfate HFA Inhalation Aerosol treatment group than in the matched placebo group. Overall, the incidence and nature of the adverse events reported for Albuterol Sulfate HFA Inhalation Aerosol and the marketed active comparator HFA-134a albuterol inhaler were comparable.

Adverse Experience Incidences (% of Patients) in a Six-Week Clinical Trial*

Body System/ Adverse Event (as Preferred Term)	Albuterol Sulfate HFA Inhalation Aerosol (N = 58)	Marketed active comparator HFA-134a albuterol inhaler (N = 56)	Matched Placebo HFA-134a Inhalation Aerosol (N = 58)
Body as a Whole Headache	7	5	2
Cardiovascular Tachycardia	3	2	0
Musculoskeletal Pain	3	0	0
Nervous System Dizziness	3	0	0
Respiratory System Pharyngitis	14	7	9
Rhinitis	5	4	2

*This table includes all adverse events (whether considered by the investigator drug related or

unrelated to drug) which occurred at an incidence rate of at least 3.0% in the Albuterol Sulfate HFA Inhalation Aerosol group and more frequently in the Albuterol Sulfate HFA Inhalation Aerosol group than in the placebo HFA Inhalation Aerosol group.

Adverse events reported by less than 3% of the patients receiving Albuterol Sulfate HFA Inhalation Aerosol but by a greater proportion of Albuterol Sulfate HFA Inhalation Aerosol patients than the matched placebo patients, which have the potential to be related to Albuterol Sulfate HFA Inhalation Aerosol, included chest pain, infection, diarrhea, glossitis, accidental injury (nervous system), anxiety, dyspnea, ear disorder, ear pain, and urinary tract infection.

In small cumulative dose studies, tremor, nervousness, and headache were the most frequently occurring adverse events.

Pediatric Patients 4 to 11 Years of Age: Adverse events reported in a 3-week pediatric clinical trial comparing the same formulation of albuterol as in Albuterol Sulfate HFA Inhalation Aerosol (180 mcg albuterol four times daily) to a matching placebo HFA inhalation aerosol occurred at a low incidence rate (no greater than 2% in the active treatment group) and were similar to those seen in adult and adolescent trials.

6.2 Postmarketing Experience

The following adverse reactions have been identified during postapproval use of Albuterol Sulfate HFA. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. Reports have included rare cases of aggravated bronchospasm, lack of efficacy, asthma exacerbation (reported fatal in one case), muscle cramps, and various oropharyngeal side-effects such as throat irritation, altered taste, glossitis, tongue ulceration, and gagging.

The following adverse events have been observed in postapproval use of inhaled albuterol: urticaria, angioedema, rash, bronchospasm, hoarseness, oropharyngeal edema, and arrhythmias (including atrial fibrillation, supraventricular tachycardia, extrasystoles). In addition, albuterol, like other sympathomimetic agents, can cause adverse reactions such as: angina, hypertension or hypotension, palpitations, central nervous system stimulation, insomnia, headache, nervousness, tremor, muscle cramps, drying or irritation of the oropharynx, hypokalemia, hyperglycemia, and metabolic acidosis.

Other short-acting sympathomimetic aerosol bronchodilators should not be used concomitantly with Albuterol Sulfate HFA Inhalation Aerosol. If additional adrenergic drugs are to be administered by any route, they should be used with caution to avoid deleterious cardiovascular effects.

7.1 Beta-Blockers

Beta-adrenergic-receptor blocking agents not only block the pulmonary effect of beta-agonists, such as Albuterol Sulfate HFA Inhalation Aerosol, but may produce severe bronchospasm in asthmatic patients. Therefore, patients with asthma should not normally be treated with beta-blockers. However, under certain circumstances, e.g., as prophylaxis after myocardial infarction, there may be no acceptable alternatives to the use of beta-adrenergic-blocking agents in patients with asthma. In this setting, consider cardioselective beta-blockers, although they should be administered with caution.

7.2 Diuretics

The ECG changes and/or hypokalemia which may result from the administration of non-potassium sparing diuretics (such as loop or thiazide diuretics) can be acutely worsened by beta-agonists, especially when the recommended dose of the beta-agonist is exceeded. Although the clinical significance of these effects is not known, caution is advised in the coadministration of beta-agonists with non-potassium sparing diuretics. Consider monitoring potassium levels.

7.3 Digoxin

Mean decreases of 16% and 22% in serum digoxin levels were demonstrated after single dose intravenous and oral administration of albuterol, respectively, to normal volunteers who had received

digoxin for 10 days. The clinical significance of these findings for patients with obstructive airway disease who are receiving albuterol and digoxin on a chronic basis is unclear. Nevertheless, it would be prudent to carefully evaluate the serum digoxin levels in patients who are currently receiving digoxin and Albuterol Sulfate HFA Inhalation Aerosol.

7.4 Monoamine Oxidase Inhibitors or Tricyclic Antidepressants

Albuterol Sulfate HFA Inhalation Aerosol should be administered with extreme caution to patients being treated with monoamine oxidase inhibitors or tricyclic antidepressants, or within 2 weeks of discontinuation of such agents, because the action of albuterol on the cardiovascular system may be potentiated. Consider alternative therapy in patients taking MAO inhibitors or tricyclic antidepressants.

8.1 Pregnancy

Pregnancy Exposure Registry

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to asthma medications during pregnancy. For more information, contact the Mothers To Baby Pregnancy Studies conducted by the Organization of Teratology Information Specialists at 1-877-311-8972 or visit <http://mothertobaby.org/pregnancy-studies/>.

Risk Summary

There are no randomized clinical studies of use of albuterol during pregnancy. Available data from published epidemiological studies and postmarketing case reports of pregnancy outcomes following inhaled albuterol use do not consistently demonstrate a risk of major birth defects or miscarriage. There are clinical considerations with use of albuterol in pregnant women [see Clinical Considerations]. In animal reproduction studies, when albuterol sulfate was administered subcutaneously to pregnant mice there was evidence of cleft palate at less than and up to 9 times the maximum recommended human daily inhalation dose (MRHDID) [see Data].

The estimated background risk of major birth defects and miscarriage for the indicated population(s) are unknown. In the U.S. general population, the estimated risk of major birth defects and miscarriage in clinically recognized pregnancies is 2 to 4% and 15 to 20%, respectively.

Clinical Considerations

Disease-Associated Maternal and/or Embryo/Fetal Risk

In women with poorly or moderately controlled asthma, there is an increased risk of preeclampsia in the mother and prematurity, low birth weight, and small for gestational age in the neonate. Pregnant women should be closely monitored and medication adjusted as necessary to maintain optimal control.

Labor or Delivery

Because of the potential for beta-agonist interference with uterine contractility, use of Albuterol Sulfate HFA Inhalation Aerosol for relief of bronchospasm during labor should be restricted to those patients in whom the benefits clearly outweigh the risk. Albuterol Sulfate HFA Inhalation Aerosol has not been approved for the management of pre-term labor. Serious adverse reactions, including pulmonary edema, have been reported during or following treatment of premature labor with beta2-agonists, including albuterol.

Data

Animal Data

In a mouse reproduction study, subcutaneously administered albuterol sulfate produced cleft palate formation in 5 of 111 (4.5%) fetuses at an exposure nine-tenths of the MRHDID for adults (on a mg/m² basis at a maternal dose of 0.25 mg/kg) and in 10 of 108 (9.3%) fetuses at approximately 9 times the MRHDID (on a mg/m² basis at a maternal dose of 2.5 mg/kg). Similar effects were not observed at approximately one-eleventh the MRHDID for adults (on a mg/m² basis at a maternal dose of 0.025 mg/kg). Cleft palate also occurred in 22 of 72 (30.5%) fetuses from females treated subcutaneously with

isoproterenol (positive control).

In a rabbit reproduction study, orally administered albuterol sulfate induced cranioschisis in 7 of 19 fetuses (37%) at approximately 750 times the MRHDID (on a mg/m² basis at a maternal dose of 50 mg/kg).

In a rat reproduction study, an albuterol sulfate/HFA-134a formulation administered by inhalation did not produce any teratogenic effects at exposures approximately 80 times the MRHDID (on a mg/m² basis at a maternal dose of 10.5 mg/kg).

A study in which pregnant rats were dosed with radiolabeled albuterol sulfate demonstrated that drug-related material is transferred from the maternal circulation to the fetus.

8.2 Lactation

Risk Summary

There are no available data on the presence of albuterol in human milk, the effects on the breastfed child, or the effects on milk production. However, plasma levels of albuterol after inhaled therapeutic doses are low in humans, and if present in breast milk, albuterol has a low oral bioavailability [see Clinical Pharmacology (12.3)].

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for albuterol and any potential adverse effects on the breastfed child from albuterol or from the underlying maternal condition.

8.4 Pediatric Use

The safety and effectiveness of Albuterol Sulfate HFA Inhalation Aerosol for the treatment or prevention of bronchospasm in children 12 years of age and older with reversible obstructive airway disease is based on one 6-week clinical trial in 116 patients 12 years of age and older with asthma comparing doses of 180 mcg four times daily with placebo, and one single-dose crossover study comparing doses of 90, 180, and 270 mcg with placebo in 58 patients [see CLINICAL STUDIES (14.1)]. The safety and effectiveness of Albuterol Sulfate HFA Inhalation Aerosol for treatment of exercise-induced bronchospasm in children 12 years of age and older is based on one single-dose crossover study in 24 adults and adolescents with exercise-induced bronchospasm comparing doses of 180 mcg with placebo [see CLINICAL STUDIES (14.2)].

The safety of Albuterol Sulfate HFA Inhalation Aerosol in children 4 to 11 years of age is based on one 3-week clinical trial in 50 patients 4 to 11 years of age with asthma using the same formulation of albuterol as in Albuterol Sulfate HFA Inhalation Aerosol comparing doses of 180 mcg four times daily with placebo. The effectiveness of Albuterol Sulfate HFA Inhalation Aerosol in children 4 to 11 years of age is extrapolated from clinical trials in patients 12 years of age and older with asthma and exercise-induced bronchospasm, based on data from a single-dose study comparing the bronchodilatory effect of Albuterol Sulfate HFA 90 mcg and 180 mcg with placebo in 55 patients with asthma and a 3-week clinical trial using the same formulation of albuterol as in Albuterol Sulfate HFA Inhalation Aerosol in 95 asthmatic children 4 to 11 years of age comparing a dose of 180 mcg albuterol four times daily with placebo [see CLINICAL STUDIES (14.1)].

The safety and effectiveness of Albuterol Sulfate HFA Inhalation Aerosol in pediatric patients below the age of 4 years have not been established.

8.5 Geriatric Use

Clinical studies of Albuterol Sulfate HFA Inhalation Aerosol did not include sufficient numbers of patients aged 65 and over to determine whether they respond differently from younger patients. Other reported clinical experience has not identified differences in responses between elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or other drug therapy [see WARNINGS AND PRECAUTIONS

(5.4,5.7)].

All beta2-adrenergic agonists, including albuterol, are known to be substantially excreted by the kidney, and the risk of toxic reactions may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.


The expected symptoms with overdosage are those of excessive beta-adrenergic stimulation and/or occurrence or exaggeration of any of the symptoms listed under ADVERSE REACTIONS, e.g., seizures, angina, hypertension or hypotension, tachycardia with rates up to 200 beats per minute, arrhythmias, nervousness, headache, tremor, dry mouth, palpitation, nausea, dizziness, fatigue, malaise, and insomnia.

Hypokalemia may also occur. As with all sympathomimetic medications, cardiac arrest and even death may be associated with abuse of Albuterol Sulfate HFA Inhalation Aerosol.

Treatment consists of discontinuation of Albuterol Sulfate HFA Inhalation Aerosol together with appropriate symptomatic therapy. The judicious use of a cardioselective beta-receptor blocker may be considered, bearing in mind that such medication can produce bronchospasm. There is insufficient evidence to determine if dialysis is beneficial for overdosage of Albuterol Sulfate HFA Inhalation Aerosol.

The oral median lethal dose of albuterol sulfate in mice is greater than 2,000 mg/kg (approximately 6,800 times the maximum recommended daily inhalation dose for adults on a mg/m² basis and approximately 3,200 times the maximum recommended daily inhalation dose for children on a mg/m² basis). In mature rats, the subcutaneous median lethal dose of albuterol sulfate is approximately 450 mg/kg (approximately 3,000 times the maximum recommended daily inhalation dose for adults on a mg/m² basis and approximately 1,400 times the maximum recommended daily inhalation dose for children on a mg/m² basis). In young rats, the subcutaneous median lethal dose is approximately 2,000 mg/kg (approximately 14,000 times the maximum recommended daily inhalation dose for adults on a mg/m² basis and approximately 6,400 times the maximum recommended daily inhalation dose for children on a mg/m² basis). The inhalation median lethal dose has not been determined in animals.

The active ingredient of Albuterol Sulfate HFA Inhalation Aerosol is albuterol sulfate, a racemic salt, of albuterol. Albuterol sulfate has the chemical name α 1-[(tert-butylamino) methyl]-4-hydroxy-m-xylene- α,α' -diol sulfate (2:1) (salt), and has the following chemical structure:



The molecular weight of albuterol sulfate is 576.7, and the empirical formula is (C₁₃H₂₁NO₃)₂•H₂SO₄. Albuterol sulfate is a white to off-white crystalline powder. It is soluble in water and slightly soluble in ethanol. Albuterol sulfate is the official generic name in the United States, and salbutamol sulfate is the World Health Organization recommended generic name. Albuterol Sulfate HFA Inhalation Aerosol is a pressurized metered-dose aerosol unit with a dose counter. Albuterol Sulfate HFA is for oral inhalation only. It contains a microcrystalline suspension of albuterol sulfate in propellant HFA-134a (1, 1, 1, 2-tetrafluoroethane) and ethanol.

Prime the inhaler before using for the first time and in cases where the inhaler has not been used for more than 2 weeks by releasing three sprays into the air, away from the face. After priming, each actuation delivers 108 mcg albuterol sulfate, from the actuator mouthpiece (equivalent to 90 mcg of albuterol base). Each canister provides 200 actuations (inhalations).

This product does not contain chlorofluorocarbons (CFCs) as the propellant.

12.1 Mechanism of Action

Albuterol sulfate is a beta2-adrenergic agonist. The pharmacologic effects of albuterol sulfate are attributable to activation of beta2-adrenergic receptors on airway smooth muscle. Activation of beta2-adrenergic receptors leads to the activation of adenylyl cyclase and to an increase in the intracellular concentration of cyclic-3', 5'-adenosine monophosphate (cyclic AMP). This increase of cyclic AMP is associated with the activation of protein kinase A, which in turn inhibits the phosphorylation of myosin and lowers intracellular ionic calcium concentrations, resulting in muscle relaxation. Albuterol relaxes the smooth muscle of all airways, from the trachea to the terminal bronchioles. Albuterol acts as a functional antagonist to relax the airway irrespective of the spasmogen involved, thus protecting against all bronchoconstrictor challenges. Increased cyclic AMP concentrations are also associated with the inhibition of release of mediators from mast cells in the airway. While it is recognized that beta2-adrenergic receptors are the predominant receptors on bronchial smooth muscle, data indicate that there are beta-receptors in the human heart, 10% to 50% of which are cardiac beta2-adrenergic receptors. The precise function of these receptors has not been established [see WARNINGS AND PRECAUTIONS (5.4)].

Albuterol has been shown in most controlled clinical trials to have more effect on the respiratory tract, in the form of bronchial smooth muscle relaxation, than isoproterenol at comparable doses while producing fewer cardiovascular effects. However, inhaled albuterol, like other beta-adrenergic agonist drugs, can produce a significant cardiovascular effect in some patients, as measured by pulse rate, blood pressure, symptoms, and/or electrocardiographic changes [see WARNINGS AND PRECAUTIONS (5.4)].

12.2 Pharmacokinetics

The systemic levels of albuterol are low after inhalation of recommended doses. In a crossover study conducted in healthy male and female volunteers, high cumulative doses of Albuterol Sulfate HFA Inhalation Aerosol (1,080 mcg of albuterol base administered over one hour) yielded mean peak plasma concentrations (C_{max}) and systemic exposure (AUC_{inf}) of approximately 4,100 pg/mL and 28,426 pg/mL*hr, respectively compared to approximately 3,900 pg/mL and 28,395 pg/mL*hr, respectively following the same dose of an active HFA-134a albuterol inhaler comparator. The terminal plasma half-life of albuterol delivered by Albuterol Sulfate HFA Inhalation Aerosol was approximately 6 hours. Comparison of the pharmacokinetic parameters demonstrated no differences between the products.

The pharmacokinetic profile of Albuterol Sulfate HFA Inhalation Aerosol was evaluated in a two-way cross-over study in 11 healthy pediatric volunteers, 4 to 11 years of age. A single dose administration of Albuterol Sulfate HFA Inhalation Aerosol (180 mcg albuterol base) yielded a least square mean (SE) C_{max} and $AUC_{0-\infty}$ of 1,100 (1.18) pg/mL and 5,120 (1.15) pg/mL*hr, respectively. The least square mean (SE) terminal plasma half-life of albuterol delivered by Albuterol Sulfate HFA Inhalation Aerosol was 166 (7.8) minutes.

Metabolism and Elimination: Information available in the published literature suggests that the primary enzyme responsible for the metabolism of albuterol in humans is SULT1A3 (sulfotransferase). When racemic albuterol was administered either intravenously or via inhalation after oral charcoal administration, there was a 3- to 4-fold difference in the area under the concentration-time curves between the (R)- and (S)-albuterol enantiomers, with (S)-albuterol concentrations being consistently higher. However, without charcoal pretreatment, after either oral or inhalation administration the differences were 8- to 24-fold, suggesting that the (R)-albuterol is preferentially metabolized in the gastrointestinal tract, presumably by SULT1A3.

The primary route of elimination of albuterol is through renal excretion (80% to 100%) of either the parent compound or the primary metabolite. Less than 20% of the drug is detected in the feces. Following intravenous administration of racemic albuterol, between 25% and 46% of the (R)-albuterol fraction of the dose was excreted as unchanged (R)-albuterol in the urine.

Geriatric, Pediatric, Hepatic/Renal Impairment: No pharmacokinetic studies for Albuterol Sulfate HFA Inhalation Aerosol have been conducted in neonates or elderly subjects.

The effect of hepatic impairment on the pharmacokinetics of Albuterol Sulfate HFA Inhalation Aerosol has not been evaluated.

The effect of renal impairment on the pharmacokinetics of albuterol was evaluated in 5 subjects with creatinine clearance of 7 to 53 mL/min, and the results were compared with those from healthy volunteers. Renal disease had no effect on the half-life, but there was a 67% decline in albuterol clearance. Caution should be used when administering high doses of Albuterol Sulfate HFA Inhalation Aerosol to patients with renal impairment [see USE IN SPECIFIC POPULATIONS (8.5)].

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

In a 2-year study in Sprague-Dawley rats, albuterol sulfate caused a dose-related increase in the incidence of benign leiomyomas of the mesovarium at and above dietary doses of 2 mg/kg (approximately 15 times and 6 times the maximum recommended human daily inhalation dose (MRHDID) for adults and children, respectively, on a mg/m² basis). In another study this effect was blocked by the coadministration of propranolol, a non-selective beta-adrenergic antagonist. In an 18-month study in CD-1 mice, albuterol sulfate showed no evidence of tumorigenicity at dietary doses of up to 500 mg/kg (approximately 1,900 times and 740 times the MRHDID for adults and children, respectively, on a mg/m² basis). In a 22-month study in Golden Hamsters, albuterol sulfate showed no evidence of tumorigenicity at dietary doses of up to 50 mg/kg (approximately 250 times and 100 times the MRHDID for adults and children, respectively, on a mg/m² basis).

Albuterol sulfate was not mutagenic in the Ames test or a mutation test in yeast. Albuterol sulfate was not clastogenic in a human peripheral lymphocyte assay or in an AH1 strain mouse micronucleus assay.

Reproduction studies in rats demonstrated no evidence of impaired fertility at oral doses up to 50 mg/kg (approximately 380 times the MRHDID for adults on a mg/m² basis).

13.2 Animal Toxicology and/or Pharmacology

Preclinical: Intravenous studies in rats with albuterol sulfate have demonstrated that albuterol crosses the blood-brain barrier and reaches brain concentrations amounting to approximately 5% of the plasma concentrations. In structures outside the blood-brain barrier (pineal and pituitary glands), albuterol concentrations were found to be 100 times those in the whole brain.

Studies in laboratory animals (minipigs, rodents, and dogs) have demonstrated the occurrence of cardiac arrhythmias and sudden death (with histologic evidence of myocardial necrosis) when β -agonists and methylxanthines were administered concurrently. The clinical significance of these findings is unknown.

Propellant HFA-134a is devoid of pharmacological activity except at very high doses in animals (380 - 1300 times the maximum human exposure based on comparisons of AUC values), primarily producing ataxia, tremors, dyspnea, or salivation. These are similar to effects produced by the structurally related chlorofluorocarbons (CFCs), which have been used extensively in metered-dose inhalers.



In animals and humans, propellant HFA-134a was found to be rapidly absorbed and rapidly eliminated, with an elimination half-life of 3 - 27 minutes in animals and 5 - 7 minutes in humans. Time to maximum plasma concentration (T_{max}) and mean residence time are both extremely short leading to a transient appearance of HFA-134a in the blood with no evidence of accumulation.

14.1 Bronchospasm Associated with Asthma

Adult and Adolescent Patients 12 Years of Age and Older: In a 6-week, randomized, double-blind, placebo-controlled trial, Albuterol Sulfate HFA Inhalation Aerosol (58 patients) was compared to a matched placebo HFA inhalation aerosol (58 patients) in asthmatic patients 12 to 76 years of age at a dose of 180 mcg albuterol four times daily. An evaluator-blind marketed active comparator HFA-134a albuterol inhaler arm (56 patients) was included.

Serial FEV₁ measurements, shown below as percent change from test-day baseline at Day 1 and at Day

43, demonstrated that two inhalations of Albuterol Sulfate HFA Inhalation Aerosol produced significantly greater improvement in FEV₁ over the pre-treatment value than the matched placebo, as well as a comparable bronchodilator effect to the marketed active comparator HFA-134a albuterol inhaler.



In this study, 31 of 58 patients treated with Albuterol Sulfate HFA Inhalation Aerosol achieved a 15% increase in FEV1 within 30 minutes post-dose on Day 1. In these patients, the median time to onset, median time to peak effect, and median duration of effect were 8.2 minutes, 47 minutes, and approximately 3 hours, respectively. In some patients, the duration of effect was as long as 6 hours.

In a placebo-controlled, single-dose, crossover study, Albuterol Sulfate HFA Inhalation Aerosol, administered at albuterol doses of 90, 180 and 270 mcg, produced bronchodilator responses significantly greater than those observed with a matched placebo HFA inhalation aerosol and comparable to a marketed active comparator HFA-134a albuterol inhaler.

Pediatric Patients 4 to 11 Years of Age: In a 3-week, randomized, double-blind, placebo-controlled trial, the same formulation of albuterol as in Albuterol Sulfate HFA Inhalation Aerosol (50 patients) was compared to a matched placebo HFA inhalation aerosol (45 patients) in asthmatic children 4 to 11 years of age at a dose of 180 mcg albuterol four times daily. Serial FEV1 measurements, expressed as the maximum percent change from test-day baseline in percent predicted FEV1 at Day 1 and at Day 22 observed within two hours post-dose, demonstrated that two inhalations of HFA albuterol sulfate produced significantly greater improvement in FEV1 over the pre-treatment value than the matched placebo.

In this study, 21 of 50 pediatric patients treated with the same formulation of albuterol as in Albuterol Sulfate HFA Inhalation Aerosol achieved a 15% increase in FEV1 within 30 minutes post-dose on Day 1. In these patients, the median time to onset, median time to peak effect and median duration of effect were 10 minutes, 31 minutes, and approximately 4 hours, respectively. In some pediatric patients, the duration of effect was as long as 6 hours.

In a placebo-controlled, single-dose, crossover study in 55 pediatric patients 4 to 11 years of age, Albuterol Sulfate HFA Inhalation Aerosol, administered at albuterol doses of 90 and 180 mcg, was compared with a matched placebo HFA inhalation aerosol. Serial FEV1 measurements, expressed as the baseline-adjusted percent predicted FEV1 observed over 6 hours post-dose, demonstrated that one and two inhalations of Albuterol Sulfate HFA Inhalation Aerosol produced significantly greater bronchodilator responses than the matched placebo.

14.2 Exercise-Induced Bronchospasm

In a randomized, single-dose, crossover study in 24 adults and adolescents with exercise-induced bronchospasm (EIB), two inhalations of Albuterol Sulfate HFA taken 30 minutes before exercise prevented EIB for the hour following exercise (defined as maintenance of FEV1 within 80% of post-dose, pre-exercise baseline values) in 83% (20 of 24) of patients as compared to 25% (6 of 24) of patients when they received placebo.

Some patients who participated in these clinical trials were using concomitant steroid therapy.

Albuterol Sulfate HFA Inhalation Aerosol is supplied as a pressurized aluminum canister with a red plastic actuator with a dose counter and white dust cap each in boxes of one. Each canister contains 8.5 g of the formulation and provides 200 actuations (NDC 0093-3174-31). Each actuation delivers 120 mcg of albuterol sulfate from the canister valve and 108 mcg of albuterol sulfate from the actuator mouthpiece (equivalent to 90 mcg of albuterol base).

SHAKE WELL BEFORE USE. Store between 15° and 25°C (59° and 77°F). Contents under pressure. Do not puncture or incinerate. Protect from freezing temperatures and prolonged exposure to direct sunlight. Exposure to temperatures above 120°F may cause bursting. For best results, canister should be at room temperature before use. Avoid spraying in eyes. Keep out of reach of children.

See FDA-APPROVED PATIENT LABELING (17.9) for priming and cleaning instructions.

The red actuator supplied with Albuterol Sulfate HFA Inhalation Aerosol should not be used with the canister from any other inhalation aerosol products. The Albuterol Sulfate HFA Inhalation Aerosol canister should not be used with the actuator from any other inhalation aerosol products.

Albuterol Sulfate HFA inhaler has a dose counter attached to the actuator. Patients should never try to alter the numbers for the dose counter or tamper with the pin mechanism inside the actuator. Discard the Albuterol Sulfate HFA inhaler when the counter displays 0 or after the expiration date on the product, whichever comes first. The labeled amount of medication in each actuation cannot be assured after the counter displays 0, even though the canister is not completely empty and will continue to operate. Never immerse the canister into water to determine how full the canister is ("float test").

Albuterol Sulfate HFA Inhalation Aerosol does not contain chlorofluorocarbons (CFCs) as the propellant.

See FDA-APPROVED PATIENT LABELING (17.9)

Patients should be given the following information:

17.1 Frequency of Use

The action of Albuterol Sulfate HFA Inhalation Aerosol should last for 4 to 6 hours. Do not use Albuterol Sulfate HFA Inhalation Aerosol more frequently than recommended. Instruct patients to not increase the dose or frequency of doses of Albuterol Sulfate HFA Inhalation Aerosol without consulting the physician. If patients find that treatment with Albuterol Sulfate HFA Inhalation Aerosol becomes less effective for symptomatic relief, symptoms become worse, and/or they need to use the product more frequently than usual, they should seek medical attention immediately.

17.2 Priming and Cleaning

Priming: Priming is essential to ensure appropriate albuterol content in each actuation. Instruct patients to prime the inhaler before using for the first time and in cases where the inhaler has not been used for more than 2 weeks by releasing three sprays into the air, away from the face.

Cleaning: To ensure proper dosing and prevent actuator orifice blockage, instruct patients to wash the red plastic actuator mouthpiece and dry thoroughly at least once a week. Instruct patients that if they have more than one Albuterol Sulfate HFA inhaler, they should wash each one at separate times to prevent attaching the wrong canister to the wrong plastic actuator. In this way, they can be sure they will always know the correct number of remaining doses. Patients should be instructed to never attach a canister of medicine from any other inhaler to the Albuterol Sulfate HFA actuator and never attach the Albuterol Sulfate HFA canister to an actuator from any other inhaler. Patients should not remove the canister from the actuator except during cleaning because reattachment may release a dose into the air and the dose counter will count down each time a spray is released. Detailed cleaning instructions are included in the illustrated Information for the Patient leaflet.

17.3 Dose Counter

Patients should be informed that Albuterol Sulfate HFA has a dose counter attached to the actuator. When the patient receives the inhaler, a black dot will appear in the viewing window until it has been primed 3 times, at which point the number 200 will be displayed. The dose counter will count down each time a spray is released. The dose-counter window displays the number of sprays left in the inhaler in units of two (e.g., 200, 198, 196, etc). When the counter displays 20, the color of the numbers will change to red to remind the patient to contact their pharmacist for a refill of medication or consult their physician for a prescription refill. When the dose counter reaches 0, the background will change to solid red. Patients should be informed to discard Albuterol Sulfate HFA inhaler when the dose counter displays 0 or after the expiration date on the product, whichever comes first.

17.4 Paradoxical Bronchospasm

Inform patients that Albuterol Sulfate HFA Inhalation Aerosol can produce paradoxical bronchospasm. Instruct patients to discontinue Albuterol Sulfate HFA Inhalation Aerosol if paradoxical bronchospasm occurs.

17.5 Concomitant Drug Use

While patients are taking Albuterol Sulfate HFA Inhalation Aerosol, other inhaled drugs and asthma

medications should be taken only as directed by a physician.

17.6 Common Adverse Events

Common adverse effects of treatment with inhaled albuterol include palpitations, chest pain, rapid heart rate, tremor, or nervousness.

17.7 Pregnancy

Patients who are pregnant or nursing should contact their physician about the use of Albuterol Sulfate HFA Inhalation Aerosol.

There is a pregnancy exposure registry that monitors pregnancy outcomes in women exposed to asthma medications during pregnancy [see Use in Specific Populations (8.1)].

17.8 General Information on Use

Effective and safe use of Albuterol Sulfate HFA Inhalation Aerosol includes an understanding of the way that it should be administered.

Shake well before each spray.

Use Albuterol Sulfate HFA Inhalation Aerosol only with the actuator supplied with the product. Discard the Albuterol Sulfate HFA inhaler when the dose counter displays 0 or after the expiration date on the product, whichever comes first. Never immerse the canister in water to determine how full the canister is ("float test").

In general, the technique for administering Albuterol Sulfate HFA Inhalation Aerosol to children is similar to that for adults. Children should use Albuterol Sulfate HFA Inhalation Aerosol under adult supervision, as instructed by the patient's physician.

Albuterol Sulfate HFA Inhalation Aerosol

Read this Patient Information before you start using Albuterol Sulfate HFA and each time you get a refill. There may be new information. This information does not take the place of talking to your doctor about your medical condition or your treatment.

What is Albuterol Sulfate HFA?

Albuterol Sulfate HFA is a prescription medicine used in people 4 years of age and older to:
treat or prevent bronchospasm in people who have reversible obstructive airway disease
prevent exercise induced bronchospasm

It is not known if Albuterol Sulfate HFA is safe and effective in children under 4 years of age.

Who should not use Albuterol Sulfate HFA?

Do not use Albuterol Sulfate HFA if you are allergic to albuterol sulfate or any of the ingredients in Albuterol Sulfate HFA. See the end of this leaflet for a complete list of ingredients in Albuterol Sulfate HFA.

What should I tell my doctor before I use Albuterol Sulfate HFA?

Before you use Albuterol Sulfate HFA, tell your doctor if you:

- have heart problems
- have high blood pressure (hypertension)
- have convulsions (seizures)
- have thyroid problems
- have diabetes
- have low potassium levels in your blood
- are pregnant or plan to become pregnant. It is not known if Albuterol Sulfate HFA will harm your unborn

baby. Talk to your doctor if you are pregnant or plan to become pregnant. are breastfeeding or plan to breastfeed. It is not known if Albuterol Sulfate HFA passes into your breast milk. Talk to your doctor about the best way to feed your baby if you are using Albuterol Sulfate HFA. Tell your doctor about all the medicines you take, including prescription and non-prescription medicines, vitamins, and herbal supplements.

Albuterol Sulfate HFA and other medicines may affect each other and cause side effects. Albuterol Sulfate HFA may affect the way other medicines work, and other medicines may affect the way Albuterol Sulfate HFA works.

Especially tell your doctor if you take:

- other inhaled medicines or asthma medicines
- beta blocker medicines
- diuretics
- digoxin
- monoamine oxidase inhibitors
- tricyclic antidepressants

Ask your doctor or pharmacist for a list of these medicines if you are not sure.

Know the medicines you take. Keep a list of them to show your doctor and pharmacist when you get a new medicine.

How should I use Albuterol Sulfate HFA?

For detailed instructions, see "Instructions for Use" at the end of this Patient Information.

Use Albuterol Sulfate HFA exactly as your doctor tells you to use it.

If your child needs to use Albuterol Sulfate HFA, watch your child closely to make sure your child uses the inhaler correctly. Your doctor will show you how your child should use Albuterol Sulfate HFA.

Each dose of Albuterol Sulfate HFA should last up to 4 hours to 6 hours.

Do not increase your dose or take extra doses of Albuterol Sulfate HFA without first talking to your doctor.

Get medical help right away if Albuterol Sulfate HFA no longer helps your symptoms.

Get medical help right away if your symptoms get worse or if you need to use your inhaler more often.

While you are using Albuterol Sulfate HFA, do not use other inhaled rescue medicines and asthma medicines unless your doctor tells you to do so.

Call your doctor if your asthma symptoms like wheezing and trouble breathing become worse over a few hours or days. Your doctor may need to give you another medicine (for example, corticosteroids) to treat your symptoms.

What are the possible side effects of Albuterol Sulfate HFA?

Albuterol Sulfate HFA may cause serious side effects, including:

- worsening trouble breathing, coughing and wheezing (paradoxical bronchospasm). If this happens stop using Albuterol Sulfate HFA and call your doctor or get emergency help right away. Paradoxical bronchospasm is more likely to happen with your first use of a new canister of medicine.

- heart problems including faster heart rate and higher blood pressure

- possible death in people with asthma who use too much Albuterol Sulfate HFA

- allergic reactions. Call your doctor right away if you have the following symptoms of an allergic reaction:

 - itchy skin

 - swelling beneath your skin or in your throat

 - rash

 - worsening trouble breathing

low potassium levels in your blood
worsening of other medical problems in people who also use Albuterol Sulfate HFA including increases in blood sugar

The most common side effects of Albuterol Sulfate HFA include:

your heart feels like it is pounding or racing (palpitations)
chest pain
fast heart rate
shakiness
nervousness
headache
dizziness
sore throat
runny nose

Tell your doctor if you have any side effect that bothers you or that does not go away.

These are not all of the possible side effects of Albuterol Sulfate HFA. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store Albuterol Sulfate HFA?

Store Albuterol Sulfate HFA at room temperature between 59° F and 77° F (15° C and 25° C).

Avoid exposure to extreme heat and cold.

Shake the Albuterol Sulfate HFA canister well before use.

Do not puncture the Albuterol Sulfate HFA canister.

Do not store the Albuterol Sulfate HFA canister near heat or a flame. Temperatures above 120° F may cause the canister to burst.

Do not throw the Albuterol Sulfate HFA canister into a fire or an incinerator.

Avoid spraying Albuterol Sulfate HFA in your eyes.

Keep Albuterol Sulfate HFA and all medicines out of the reach of children.

General Information about the safe and effective use of Albuterol Sulfate HFA

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use Albuterol Sulfate HFA for a condition for which it was not prescribed. Do not give Albuterol Sulfate HFA to other people, even if they have the same symptoms that you have. It may harm them.

This Patient Information summarizes the most important information about Albuterol Sulfate HFA. If you would like more information, talk with your doctor. You can ask your pharmacist or doctor for information about Albuterol Sulfate HFA that is written for health professionals.

For more information, call 1-888-482-9522.

What are the ingredients in Albuterol Sulfate HFA?

Active ingredient: albuterol sulfate

Inactive ingredients: propellant HFA-134a and ethanol.

ASHFAPL-002

Rev. 1/2018

Instructions for Use

Albuterol Sulfate HFA
Inhalation Aerosol

Read this Instructions for Use before you start using Albuterol Sulfate HFA and each time you get a refill. There may be new information. This information does not take the place of talking to your doctor about your medical condition or your treatment.

The Parts of Your Albuterol Sulfate HFA Inhaler Device:

There are 2 main parts of your Albuterol Sulfate HFA inhaler device including a:


red plastic actuator that sprays the medicine from the canister. See Figure A.

protective dust cap that covers the mouthpiece of the actuator. See Figure A.

There is also a metal canister that holds the medicine. See Figure A.

There is also a dose counter attached to the back of the actuator with a viewing window that shows you how many sprays of medicine you have left. See Figure B.

You will see a black dot in the viewing window on the actuator until the device has been primed 3 times. See Figure B and "Priming Your Albuterol Sulfate HFA Device" below.



Do not use the Albuterol Sulfate HFA actuator with a canister of medicine from any other inhaler. Do not use an Albuterol Sulfate HFA canister with an actuator from any other inhaler, including another Albuterol Sulfate HFA inhaler.

Priming Your Albuterol Sulfate HFA Device:

Your Albuterol Sulfate HFA device must be primed before you use it for the first time or if your device has not been used for more than 14 days in a row. Do not prime your Albuterol Sulfate HFA device every day.

Remove your Albuterol Sulfate HFA device from its package.

Remove the protective dust cap from the mouthpiece.

Shake the inhaler well, and spray it into the air away from your face. See Figure C.




Figure C

Shake and spray the inhaler like this 2 more times to finish priming it.

The dose counter on the actuator should display the number 200 after you prime the actuator for the first time. See Figure D.




Figure D

Each Time You Use Your Albuterol Sulfate HFA Device:

Make sure the canister fits firmly in the plastic actuator.

Look into the mouthpiece to make sure there are no foreign objects there, especially if the cap has not been used to cover the mouthpiece.

Reading the Dose Counter on Your Albuterol Sulfate HFA Actuator

The dose counter will count down each time a spray is released. The dose counter window shows the number of sprays left in your inhaler in units of 2 sprays. For example, there are 190 sprays left if the arrow is exactly opposite the number 190, or 189 sprays left if the arrow points between 190 and 188. See Figure D.

When the dose counter reaches 0, it will continue to show 0 and you should replace your Albuterol Sulfate HFA device.

The dose counter cannot be reset and is permanently attached to the actuator. Never change the numbers for the dose counter or touch the pin inside the actuator.

Do not remove the canister from the plastic actuator except during cleaning. Reattaching the canister to the actuator may accidentally release a dose of Albuterol Sulfate HFA into the air. The dose counter will count down each time a spray is released.

Using Your Albuterol Sulfate HFA Device:

Step 1. Shake the inhaler well before each spray. Take the cap off the mouthpiece of the actuator.

Step 2. Hold the inhaler with the mouthpiece down. See Figure E.



Figure E

Step 3. Breathe out through your mouth and push as much air from your lungs as you can. Put the mouthpiece in your mouth and close your lips around it. See Figure F.

Step 4. Push the top of the canister all the way down while you breathe in deeply and slowly through your mouth. See Figure F.



Figure F

Step 5. Right after the spray comes out, take your finger off the canister. After you have breathed in all the way, take the inhaler out of your mouth and close your mouth.


Step 6. Hold your breath as long as you can, up to 10 seconds, then breathe normally.

If your doctor has told you to use more sprays, wait 1 minute and shake the inhaler again. Repeat Steps 2 through Step 6.

Step 7. Put the cap back on the mouthpiece after every time you use the inhaler. Make sure the cap snaps firmly into place.

Cleaning Your Albuterol Sulfate HFA Device:

It is very important to keep the plastic actuator clean so the medicine will not build-up and block the spray. See Figure G and Figure H.



Do not try to clean the metal canister or let it get wet. The inhaler may stop spraying if it is not cleaned correctly.

If you have more than 1 Albuterol Sulfate HFA inhaler, wash each device at separate times to prevent putting the wrong canister together with the wrong plastic actuator. This way you can be sure you will always know the correct number of remaining doses of Albuterol Sulfate HFA.

Wash the actuator at least 1 time each week as follows:

Take the canister out of the actuator, and take the cap off the mouthpiece.

Hold the actuator under the faucet and run warm water through it for about 30 seconds. See Figure I.



Figure I

Turn the actuator upside down and run warm water through the mouthpiece for about 30 seconds. See Figure J.




Figure J

Shake off as much water from the actuator as you can. Look into the mouthpiece to make sure any medicine build-up has been completely washed away. If there is any build-up, repeat the washing instructions.

Let the actuator air-dry completely, such as overnight. See Figure K.



Figure K

When the actuator is dry, put the canister in the actuator and make sure it fits firmly. Shake the inhaler well and spray it twice into the air away from your face. Put the cap back on the mouthpiece.

If you need to use your inhaler before the actuator is completely dry:

Shake as much water off the actuator as you can.

Put the canister in the actuator and make sure it fits firmly.

Shake the inhaler well and spray it twice into the air away from your face.

Take your Albuterol Sulfate HFA dose as prescribed.

Follow the Cleaning Instructions above.

Replacing Your Albuterol Sulfate HFA Device

When the dose counter on the actuator says the number 20, the color of the numbers will change to red. The red numbers are to remind you to refill your prescription or ask your doctor for another prescription for Albuterol Sulfate HFA. When the dose counter reaches 0, the background color will change to solid red.

Throw the Albuterol Sulfate HFA inhaler away as soon as the dose counter says 0 or after the expiration date on the Albuterol Sulfate HFA packaging, whichever comes first. You should not keep using the inhaler after 200 sprays even though the canister may not be completely empty. You cannot be sure you will receive any medicine after using 200 sprays.

Do not use the inhaler after the expiration date on the Albuterol Sulfate HFA packaging.

This Patient Information and Instructions for Use has been approved by the U.S. Food and Drug Administration.

Marketed by Teva Pharmaceuticals USA, Inc.
North Wales, PA 19454

Manufactured by IVAX Pharmaceuticals Ireland
Waterford, Ireland

Revised 1/2018

ASHFAIFU-002



CAUTION: federal Law PROHIBITS the transfer of this drug to anyone other than the person whom prescribed and prohibits dispensing without a prescription, unless OTC. See outsert for add'l Rx info. KEEP OUT OF REACH OF CHILDREN Store in a cool, dry place at 68-77 F unless printed otherwise.

76519-1171-00

ALBUT SUL HFA(PROAIR) 90mcg INH 8.5gm

Compare: Proair

Exp. 00/00 Lot#: AB00CD

Mfg. TEVA PHARM 00093-3174-31

ACCOUNT: 00-0000

Use As Directed by Physician

ALBUT SUL HFA(PROAIR) 90mcg INH
NDC: 76519-1171-00 QTY: 8.5gm
Exp. 00/00 Lot#: AB00CD
MFG NDC 00093-3174-31

ALBUT SUL HFA(PROAIR) 90mcg INH
NDC: 76519-1171-00 QTY: 8.5gm
Exp. 00/00 Lot#: AB00CD
MFG NDC 00093-3174-31

ALBUT SUL HFA(PROAIR) 90mcg INH
NDC: 76519-1171-00 QTY: 8.5gm
Exp. 00/00 Lot#: AB00CD
MFG NDC 00093-3174-31

ALBUT SUL HFA(PROAIR) 90mcg INH
NDC: 76519-1171-00 QTY: 8.5gm
Exp. 00/00 Lot#: AB00CD
MFG NDC 00093-3174-31

Repack: H.J. Harkins Co., Inc. Grover Beach, CA 93433

ALBUTEROL SULFATE HFA

albuterol sulfate aerosol, metered

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:76519-1171
Route of Administration	RESPIRATORY (INHALATION)		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALBUTEROL SULFATE (UNII: 021SEF3731) (ALBUTEROL - UNII:QF8SVZ843E)	ALBUTEROL	90 ug in 1 g

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76519-1171-0	8.5 g in 1 INHALER; Type 0: Not a Combination Product	03/25/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA021457	03/25/2019	

Labeler - H.J. Harkins Company, Inc. (147681894)

Establishment

Name	Address	ID/FEI	Business Operations
H.J. Harkins Company, Inc.		147681894	relabel(76519-1171) , repack(76519-1171) , manufacture(76519-1171)

Revised: 4/2019

H.J. Harkins Company, Inc.