

**ALL DAY PAIN RELIEF- naproxen sodium tablet, coated
WALGREENS**

**608R WALGREEN 0363-9608 NAPROXEN SODIUM CAPLETS ALL DAY PAIN
RELIEF**

DRUG FACTS

Naproxen Sodium 220 mg

(naproxen 200 mg)(NSAID)*

*nonsteroidal anti-inflammatory drug

Pain reliever/Fever reducer

- temporarily relieves minor aches and pains due to:
 - o minor pain of arthritis
 - o muscular aches
 - o backache
 - o menstrual cramps
 - o headache
 - o toothache
 - o the common cold

- temporarily reduces fever

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

hives
facial swelling
asthma (wheezing)
shock
skin reddening
rash
blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

are age 60 or older
have had stomach ulcers or bleeding problems
take blood thinning (anticoagulant) or steroid drug
take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
have 3 or more alcoholic drinks every day while using this product.
take more or for a longer time than directed.

Heart attack and stroke warning: NSAID's, except aspirin, increases the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:

feel faint

vomit blood

have bloody or black stools

have stomach pain that does not get better

you have symptoms of heart problems or stroke:

chest pain

trouble breathing

weakness in one part or side of body

slurred speech

leg swelling

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

do not take more than directed
the smallest effective dose should be used

drink a full glass of water with each dose

Adults and children 12 years and older:

take 1 caplet every 8 to 12 hours while symptoms last
for the first dose, you may take 2 caplets within the first hour
do not exceed 2 caplets in any 8-to 12-hour period
do not exceed 3 caplets in a 24-hour period

Children under 12 years:

ask a doctor

Other information

each caplet contains: sodium 20 mg
store at 20-25°C (68-77°F).

Avoid high humidity and excessive heat above 40°C (104°F)

Colloidal silicon dioxide, croscarmellose sodium, FD&C blue #2 lake, Hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, titanium dioxide.

Questions or comments? call **1-877-290-4008**

NDC 0363-9608-09

Walgreens

All Day Pain Relief

NAPROXEN SODIUM TABLETS USP,
220 mg / PAIN RELIEVER /
FEVER REDUCER (NSAID)

• Strength to last 12 hours

90 CAPLETS

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Uses ■ temporarily relieves minor aches and pains due to:
● minor pain of arthritis ● muscular aches ● backache
● menstrual cramps ● headache ● toothache ● the common cold ■ temporarily reduces fever

Warnings Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin.
Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. **If pregnant or breast-feeding**, ask a health professional before use.

Keep out of reach of children.

Directions ■ do not take more than directed ■ the smallest effective dose should be used ■ drink a full glass of water with each dose Adults and children 12 years and older ■ take 1 caplet every 8 to 12 hours while symptoms last ■ for the first dose you may take 2 caplets within the first hour ■ do not exceed 2 caplets in any 8- to 12-hour period ■ do not exceed 3 caplets in a 24-hour period Children under 12 years ■ ask a doctor

Other information ■ **TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING** ■ each caplet contains: sodium 20 mg ■ store at 20-25°C (68-77°F). ■ avoid high humidity and excessive heat above 40°C (104°F).

Questions or comments? call 1-877-290-4008 608R REV 0924
DISTRIBUTED BY: WALGREEN CO., 200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED walgreens.com ©2021 Walgreen Co.
MADE IN INDIA ITEM 988355 W00000-0000-0 W30RG1221-F

Lot No.:
Exp Date:

Varnish Omit Area

Adhesive Area

Drug Facts (continued)

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you are taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke: ● chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period

Children under 12 years: ■ ask a doctor

Other information

- each caplet contains: sodium 20 mg
- store at 20-25°C (68-77°F).
- avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, F&G Blue #2 lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinylpyrrolidone, titanium dioxide

Questions or comments? call 1-877-290-4008

Inside

Walgreens

Compare to the active ingredient in ALEVE® Caplets*

EASY OPEN

All Day Pain Relief

NAPROXEN SODIUM TABLETS USP, 220 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

12 Hour • Strength to last 12 hours

90 CAPLETS

ACTUAL SIZE

PACKAGE NOT CHILD RESISTANT

NDC 0363-0608-90

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Drug Facts

Active ingredient (in each caplet)

Naproxen sodium 220 mg (NSAID)* Pain reliever
(naproxen 220 mg) Fever reducer
nonsteroidal anti-inflammatory drug

Purposes

Pain reliever
Fever reducer

Uses ■ temporarily relieves minor aches and pains due to:

- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ facial swelling (wheezing) ■ shock ■ skin reddening ■ rash ■ asthma (wheezing) ■ shock ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older

- have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed
- Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Drug Facts (continued under label)

Walgreens Pharmacist Recommended. Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Aleve®.

DISTRIBUTED BY: WALGREEN CO. ITEM 210503 W00000-0000-Q
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
Walgreens.com ©2022 Walgreen Co.
MADE IN INDIA 68R 022 REV-0423
W3ORG0322-F



Lot No.:

Varnish Omit Area

Exp Date:



Adhesive Area

Drug Facts (continued)

Do not use

■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ right before or after heart surgery

Ask a doctor before use if ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

■ under a doctor's care for any serious condition
■ taking aspirin for heart attack or stroke, because naproxen may decrease this benefit or aspirin ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding:
● feel faint ● vomit blood ● have bloody or black stools
● have stomach pain that does not get better
■ you have symptoms of heart problems or stroke ● chest pain
● trouble breathing ● weakness in one part or side of body
● slurred speech ● leg swelling
■ pain gets worse or lasts more than 10 days
■ you have difficulty swallowing
■ it feels like the pill is stuck in your throat
■ redness or swelling is present in the painful area
■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions ■ do not take more than directed

■ the smallest effective dose should be used

■ drink a full glass of water with each dose

Adults and children 12 years and older

■ take 1 caplet every 8 to 12 hours with symptoms best
■ for the first dose you may take 2 caplets within the first hour
■ do not exceed 2 caplets in any 8- to 12-hour period
■ do not exceed 3 caplets in a 24-hour period

Children under 12 years

■ ask a doctor

Other information

■ each caplet contains: sodium 20 mg ■ store at 20°-25°C (68°-77°F).
■ avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients: colloidal silicon dioxide, croscarmellose sodium, FD&C Blue #2 Lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, titanium dioxide

Questions or comments? call 1-877-290-4008

Inside

Varnish Omit Area

Lot No.:

Exp Date:



Drug Facts (continued under label)

Walgreens Pharmacist Recommended. Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Aleve®.

DISTRIBUTED BY: WALGREEN CO. ITEM 517569 W000000-00000-0
200 WILLMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
Walgreens.com ©2022 Walgreen Co.
MADE IN INDIA 608R REV 0921 3
W3ORG1221-F REV-0423



Drug Facts

Active ingredient (in each caplet)

Naproxen sodium 220 mg
(naproxen 200 mg) (NSAID)*
*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/
fever reducer

Uses

■ temporarily relieves minor aches and pains due to:
● minor pain of arthritis ● backache ● menstrual cramps
● headache ● toothache ● the common cold
■ temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause severe allergic reaction, especially in people allergic to aspirin. Symptoms include:
■ hives ■ facial swelling ■ asthma (wheezing)
■ shock ■ skin redness ■ rash ■ blisters
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

VALUE SIZE NDC 0363-9608-61

Walgreens

Compare to the active ingredient in ALEVE® Caplets*

All Day Pain Relief
NAPROXEN SODIUM TABLETS USP, 220 mg /
PAIN RELIEVER / FEVER REDUCER (NSAID)

12 Hour • Strength to last 12 hours

600
CAPLETS

ACTUAL SIZE



TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING

Varnish Omit

W3ORG1221-F



THE BOTTLE IS CUT, TORN, BROKEN OR MISSING
DO NOT USE THIS PRODUCT IF
TAMPER EVIDENT - DO NOT USE THIS PRODUCT IF
DO NOT OPENED FOR SEALING PURPOSES

NDC 0363-9608-09

Walgreens

Compare to the active ingredient
in ALEVE® Caplets*



Varnish Omit

E1891

All Day Pain Relief

NAPROXEN SODIUM TABLETS USP,
220 mg / PAIN RELIEVER /
FEVER REDUCER (NSAID)

12 Hour
• Strength to last 12 hours

90
CAPLETS

ACTUAL SIZE

Varnish Omit

Drug Facts (continued)

Adults and children 12 years and older

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period

Children under 12 years:

- ask a doctor

Other information

- each caplet contains: sodium 20 mg
- store at 20-25°C (68-77°F).
- avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients croscarmellose sodium, croscarmellose sodium, FD&C Blue #2 Lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, titanium dioxide

Questions or comments? call 1-877-290-4008

*Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.
†This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Aleve®.

Drug Facts (continued)

Ask a doctor before use if ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are ■ under a doctor's care for any serious condition ■ taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin ■ taking any other drug

When using this product ■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ you have difficulty swallowing ■ it feels like the pill is stuck in your throat ■ redness or swelling is present in the painful area ■ any new symptoms appear

■ pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Drug Facts

Active ingredient (in each caplet)	Purposes
Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	

Uses ■ temporarily relieves minor aches and pains due to: ■ minor pain of arthritis ■ muscular aches ■ backache ■ menstrual cramps ■ headache ■ toothache ■ the common cold ■ temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 5 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

DISTRIBUTED BY: WALGREEN CO.
200 WILMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
©2022 Walgreen Co. MADE IN INDIA
ITEM 988355 W00000-0000-0



Lot No.:
Exp Date: Print & Varnish Omit

60881 REV 0021

Lot No.:
Exp Date:

NDC 0363-9608-05

Walgreens

All Day Pain Relief

NAPROXEN SODIUM TABLETS USP,
220 mg / PAIN RELIEVER /
FEVER REDUCER (NSAID)

• Strength to last 12 hours

50 CAPLETS

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Uses ■ temporarily relieves minor aches and pains due to:
 • minor pain of arthritis ■ muscular aches ■ backache
 • menstrual cramps ■ headache ■ toothache ■ the common cold ■ temporarily reduces fever

Warnings Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away. **Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed **Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed. If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** **Directions** ■ do not take more than directed ■ the smallest effective dose should be used ■ drink a full glass of water with each dose. Adults and children 12 years and older ■ take 1 caplet every 8 to 12 hours while symptoms last ■ for the first dose you may take 2 caplets within the first hour ■ do not exceed 2 caplets in any 8- to 12-hour period ■ do not exceed 3 caplets in a 24-hour period ■ ask a doctor.

Other information ■ **TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN, OR MISSING** ■ each caplet contains: sodium 220 mg ■ store at 20-25°C (68-77°F) ■ avoid high humidity and excessive heat above 40°C (104°F) ■ 608R REV 0921.

Questions or comments? call 1-877-290-4008
 DISTRIBUTED BY: WALGREEN CO. ITEM 517531
 200 WILMOT RD., DEERFIELD, IL 60015 W00000-0000-0
 100% SATISFACTION GUARANTEED W3ORG1221-F
 walgreens.com ©2021 Walgreen Co. MADE IN INDIA
 40/25/02 Code No.: 608RUGS515
 Lot No.:

Exp. Date:

Varnish Omit Area

Varnish Omit

W3ORG1221-F
REV-1022

NDC 0363-9608-05

Walgreens

Compare to the active ingredient
in ALEVE® Caplets*

All Day Pain Relief

NAPROXEN SODIUM TABLETS USP,
220 mg / PAIN RELIEVER /
FEVER REDUCER (NSAID)

12 Hour

• Strength to last 12 hours

50
CAPLETS

ACTUAL SIZE

*Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by Bayer Healthcare LLC, owner of the registered trademark Aleve®.

TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING



ACTUAL SIZE



Varnish Omit

Drug Facts (continued)

pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke ■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

■ under a doctor's care for any serious condition
 ■ taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
 ■ taking any other drug

When using this product

■ take with food or milk if stomach upset occurs

Stop use and ask a doctor if

■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better ■ you have symptoms of heart problems or stroke: ■ chest pain ■ trouble breathing ■ weakness in one part or side of body ■ slurred speech ■ leg swelling ■ pain gets worse or lasts more than 10 days ■ fever gets worse or lasts more than 3 days ■ you have difficulty swallowing ■ it feels like the pill is stuck in your throat ■ redness or swelling is present in the painful area ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

■ do not take more than directed
 ■ the smallest effective dose should be used
 ■ drink a full glass of water with each dose
 Adults and children 12 years and older
 ■ take 1 caplet every 8 to 12 hours while symptoms last
 ■ for the first dose you may take 2 caplets within the first hour
 ■ do not exceed 2 caplets in any 8- to 12-hour period
 ■ do not exceed 3 caplets in a 24-hour period

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

DISTRIBUTED BY: WALGREEN CO.
 200 WILMOT RD., DEERFIELD, IL 60015
 100% SATISFACTION GUARANTEED
 walgreens.com ©2022 Walgreen Co.
 MADE IN INDIA

ITEM 517531 W00000-0000-0



Lot No.:

Exp. Date:

Print & Varnish Omit

Drug Facts

Active Ingredient (in each caplet) Naproxen sodium 220 mg (naproxen 220 mg) (NSAID)*
Purposes Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Uses

■ temporarily relieves minor aches and pains due to:
 • minor pain of arthritis ■ muscular aches
 • backache ■ menstrual cramps ■ headache
 • toothache ■ the common cold
 ■ temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters. If an allergic reaction occurs, stop use and seek medical help right away.


Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer ■ right before or after heart surgery

Ask a doctor before use if ■ the stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood

608R REV 0921



All Day Pain Relief

NAPROXEN SODIUM TABLETS,
220 mg / PAIN RELIEVER /
FEVER REDUCER (NSAID)

NDC 0363-9608-27

220 mg / PAIN RELIEVER / FEVER REDUCER (NSAID)

12 HOUR CAPLETS

• Strength to last 12 hours

270 CAPLETS

READ AND KEEP CARTON FOR COMPLETE WARNINGS AND INFORMATION

Uses ■ temporarily relieves minor aches and pains due to: ■ minor pain of arthritis ■ muscular aches ■ backache ■ menstrual cramps ■ headache ■ toothache ■ the common cold ■ temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock ■ skin reddening ■ rash ■ blisters If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood-thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 2 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Do not take more than directed
■ the **smallest effective dose should be used**
■ drink a full glass of water with each dose

adults and children 12 years and older:
■ take 1 tablet every 8 to 12 hours while symptoms last
■ for the first dose you may take 2 tablets within the first hour
■ do not exceed 2 tablets in any 8- to 12-hour period
■ do not exceed 3 tablets in a 24-hour period
■ children under 12 years: ■ ask a doctor

Other information ■ TAMPER EVIDENT, DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING
■ each tablet contains **sodium 20 mg**
■ store at 20-25°C (68-77°F)
■ avoid high humidity and excessive heat above 40° C (104°F)

Questions or comments? 1-877-250-4008

DISTRIBUTED BY: WALGREEN CO.
200 WILLMOT RD., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com ©2019 Walgreen Co.

ITEM 1792933
W00000-0000-0
608R REV 0519
ORG0619-F

Lot No:

Exp Date:

Varnish Omit Area

naproxen sodium tablet, coated

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-9608
Route of Administration	ORAL		

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Ingredient Name	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
FD&C BLUE NO. 2--ALUMINUM LAKE (UNII: 4AQJ3LG584)	

Color	blue	Score	no score
Shape	OVAL	Size	12mm
Flavor		Imprint Code	144
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-9608-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	06/30/2019	
2	NDC:0363-9608-20	200 in 1 BOTTLE; Type 0: Not a Combination Product	06/30/2019	
3	NDC:0363-9608-27	270 in 1 BOTTLE; Type 0: Not a Combination Product	06/30/2019	
4	NDC:0363-9608-61	600 in 1 BOTTLE; Type 0: Not a Combination Product	06/30/2019	
5	NDC:0363-9608-09	1 in 1 CARTON	06/26/2021	
5		90 in 1 BOTTLE; Type 0: Not a Combination Product		
6	NDC:0363-9608-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	04/01/2022	
7	NDC:0363-9608-42	1 in 1 CARTON	06/30/2019	
7		24 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090545	06/26/2019	

Labeler - WALGREENS (008965063)**Registrant** - TIME CAP LABORATORIES, INC (037052099)**Establishment**

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LTD		925822975	manufacture(0363-9608)

Revised: 3/2024

WALGREENS