


HYDROXYZINE HYDROCHLORIDE- hydroxyzine hydrochloride tablet, film coated
Mutual Pharmaceutical Company, Inc.

HydrOXYzine HYDROCHLORIDE TABLETS USP

Rx only

DESCRIPTION

HydrOXYzine hydrochloride is designated chemically as 1-(p-chlorobenzhydryl) 4-[2-(2-hydroxy-ethoxy)-ethyl] piperazine dihydrochloride.



Each tablet, for oral administration, contains 10 mg, 25 mg, or 50 mg of hydrOXYzine hydrochloride.

Inactive ingredients: anhydrous lactose, colloidal silicon dioxide, croscarmellose sodium, FD&C Blue No. 2 Aluminum lake (not contained in the 25 mg or 50 mg pink tablets), hypromellose, magnesium stearate, polyethylene glycol, and titanium dioxide. The 25 mg includes hydroxypropyl cellulose; and the 50 mg includes polysorbate 80. The lavender 10 mg includes D&C Red No. 27 Aluminum lake; FD&C Red No. 40 Aluminum lake; FD&C Yellow No. 6 Aluminum lake; polydextrose and triacetin. The 25 mg and 50 mg include D&C Red No. 7 Calcium lake.

CLINICAL PHARMACOLOGY

HydrOXYzine hydrochloride is unrelated chemically to the phenothiazines, reserpine, meprobamate, or the benzodiazepines.

HydrOXYzine is not a cortical depressant, but its action may be due to a suppression of activity in certain key regions of the subcortical area of the central nervous system. Primary skeletal muscle relaxation has been demonstrated experimentally. Bronchodilator activity, and antihistaminic and analgesic effects have been demonstrated experimentally and confirmed clinically. An antiemetic effect, both by the apomorphine test and the veriloid test, has been demonstrated. Pharmacological and clinical studies indicate that hydrOXYzine in therapeutic dosage does not increase gastric secretion or acidity and in most cases has mild antisecretory activity. HydrOXYzine is rapidly absorbed from the gastrointestinal tract and hydrOXYzine hydrochloride clinical effects are usually noted within 15 to 30 minutes after oral administration.

INDICATIONS

For symptomatic relief of anxiety and tension associated with psychoneurosis and as an adjunct in organic disease states in which anxiety is manifested.

Useful in the management of pruritus due to allergic conditions such as chronic urticaria and atopic and contact dermatoses, and in histamine-mediated pruritus.

As a sedative when used as premedication and following general anesthesia, **hydrOXYzine may potentiate meperidine and barbiturates**, so their use in pre-anesthetic adjunctive therapy should be modified on an individual basis. Atropine and other belladonna alkaloids are not affected by the drug. HydrOXYzine is not known to interfere with the action of digitalis in any way and it may be used concurrently with this agent.

The effectiveness of hydrOXYzine as an antianxiety agent for long term use, that is more than 4 months, has not been assessed by systematic clinical studies. The physician should reassess periodically the usefulness of the drug for the individual patient.

CONTRAINDICATIONS

HydrOXYzine, when administered to the pregnant mouse, rat, and rabbit, induced fetal abnormalities in the rat and mouse at doses substantially above the human therapeutic range. Clinical data in human beings are inadequate to establish safety in early pregnancy. Until such data are available, hydrOXYzine is contraindicated in early pregnancy.

HydrOXYzine is contraindicated for patients who have shown a previous hypersensitivity to it.

WARNINGS

Nursing Mothers

It is not known whether this drug is excreted in human milk. Since many drugs are so excreted, hydrOXYzine should not be given to nursing mothers.

PRECAUTIONS

THE POTENTIATING ACTION OF HYDROXYZINE MUST BE CONSIDERED WHEN THE DRUG IS USED IN CONJUNCTION WITH CENTRAL NERVOUS SYSTEM DEPRESSANTS SUCH AS NARCOTICS, NON-NARCOTIC ANALGESICS AND BARBITURATES. Therefore when central nervous system depressants are administered concomitantly with hydrOXYzine their dosage should be reduced.

Since drowsiness may occur with use of this drug, patients should be warned of this possibility and cautioned against driving a car or operating dangerous machinery while taking hydrOXYzine. Patients should be advised against the simultaneous use of other CNS depressant drugs, and cautioned that the effect of alcohol may be increased.

Geriatric Use

A determination has not been made whether controlled clinical studies of hydrOXYzine included sufficient numbers of subjects aged 65 and over to define a difference in response from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal or cardiac function, and of concomitant disease or other drug therapy.

The extent of renal excretion of hydrOXYzine has not been determined. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selections.

Sedating drugs may cause confusion and over sedation in the elderly; elderly patients generally should

be started on low doses of hydrOXYzine and observed closely.

ADVERSE REACTIONS

Side effects reported with the administration of hydrOXYzine hydrochloride are usually mild and transitory in nature.

Anticholinergic: Dry mouth.

Central Nervous System: Drowsiness is usually transitory and may disappear in a few days of continued therapy or upon reduction of the dose. Involuntary motor activity including rare instances of tremor and convulsions have been reported, usually with doses considerably higher than those recommended. Clinically significant respiratory depression has not been reported at recommended doses.

OVERDOSAGE

The most common manifestation of hydrOXYzine overdosage is hypersedation. As in the management of overdosage with any drug, it should be borne in mind that multiple agents may have been taken.

If vomiting has not occurred spontaneously, it should be induced. Immediate gastric lavage is also recommended. General supportive care, including frequent monitoring of the vital signs and close observation of the patient, is indicated. Hypotension, though unlikely, may be controlled with intravenous fluids and levarterenol, or metaraminol. Do not use epinephrine as hydrOXYzine counteracts its pressor action.

There is no specific antidote. It is doubtful that hemodialysis would be of any value in the treatment of overdosage with hydrOXYzine. However, if other agents such as barbiturates have been ingested concomitantly, hemodialysis may be indicated. There is no practical method to quantitate hydrOXYzine in body fluids or tissue after its ingestion or administration.

DOSAGE

For symptomatic relief of anxiety and tension associated with psychoneurosis and as an adjunct in organic disease states in which anxiety is manifested: in adults, 50-100 mg q.i.d.; children under 6 years, 50 mg daily in divided doses and over 6 years, 50-100 mg daily in divided doses.

For use in the management of pruritus due to allergic conditions such as chronic urticaria and atopic and contact dermatoses, and in histamine-mediated pruritus: in adults, 25 mg t.i.d. or q.i.d.; children under 6 years, 50 mg daily in divided doses and over 6 years, 50-100 mg daily in divided doses.

As a sedative when used as a premedication and following general anesthesia: 50-100 mg in adults, and 0.6 mg/kg in children.

When treatment is initiated by the intramuscular route of administration, subsequent doses may be administered orally.

As with all medications, the dosage should be adjusted according to the patient's response to therapy.

HOW SUPPLIED

HydrOXYzine hydrochloride tablets, USP are supplied as follows:

HydrOXYzine hydrochloride tablets, 10 mg, lavender, round, film-coated, debossed MP 3

Bottles of 100 NDC 53489-126-01

Bottles of 500 NDC 53489-126-05

Bottles of 1000 NDC 53489-126-10

HydroXYzine hydrochloride tablets, 25 mg, purple, round, film-coated, debossed MP 7

Bottles of 100 NDC 53489-127-01

Bottles of 500 NDC 53489-127-05

Bottles of 1000 NDC 53489-127-10

HydroXYine hydrochloride tablets, 50 mg, purple, round, film-coated, debossed MP 13

Bottles of 100 NDC 53489-128-01

Bottles of 500 NDC 53489-128-05

Bottles of 1000 NDC 53489-128-10

HydroXYine hydrochloride tablets, 25 mg, pink, round, film-coated, debossed MP 7

Bottles of 100 NDC 53489-594-01

Bottles of 500 NDC 53489-594-05

Bottles of 1000 NDC 53489-594-10

HydroXYine hydrochloride tablets, 50 mg, pink, round, film-coated, debossed MP 13

Bottles of 100 NDC 53489-595-01

Bottles of 500 NDC 53489-595-05

Bottles of 1000 NDC 53489-595-10

Store at 20° to 25° C (68° to 77°F).

[See USP Controlled Room Temperature]

DISPENSE IN TIGHT, LIGHT-RESISTANT CONTAINER.

Manufactured By:

MUTUAL PHARMACEUTICAL COMPANY, INC.

Philadelphia, PA 19124 USA

Revised: September 2005NP

HYDROXYZINE HYDROCHLORIDE

hydroxyzine hydrochloride tablet, film coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:53489-126
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Hydroxyzine hydrochloride (UNII: 76755771U3) (Hydroxyzine - UNII:30S50YM8OG)		10 mg

Inactive Ingredients

Ingredient Name	Strength
Anhydrous lactose ()	
colloidal silicon dioxide ()	
croscarmellose sodium ()	
FD&C Blue No. 2 Aluminum lake ()	

hypromellose ()	
magnesium stearate (UNII: 70097M6I30)	
polyethylene glycol ()	
titanium dioxide (UNII: 15FIX9V2JP)	
D&C Red No. 27 Aluminum lake ()	
FD&C Red No. 40 Aluminum lake ()	
FD&C Yellow No. 6 Aluminum lake ()	
polydextrose ()	
triacetin (UNII: XHX3C3X673)	

Product Characteristics

Color	PURPLE (Lavender)	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	MP;3
Contains			
Coating	true	Symbol	false

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53489-126-01	100 in 1 BOTTLE, PLASTIC		
2	NDC:53489-126-05	500 in 1 BOTTLE, PLASTIC		
3	NDC:53489-126-10	1000 in 1 BOTTLE, PLASTIC		

HYDROXYZINE HYDROCHLORIDE

hydroxyzine hydrochloride tablet, film coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:53489-127
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Hydroxyzine hydrochloride (UNII: 76755771U3) (Hydroxyzine - UNII:30S50YM8OG)		25 mg

Inactive Ingredients

Ingredient Name	Strength
Anhydrous lactose ()	
colloidal silicon dioxide ()	
croscarmellose sodium ()	
FD&C Blue No. 2 Aluminum lake ()	
hypromellose ()	
magnesium stearate (UNII: 70097M6I30)	

polyethylene glycol ()	
titanium dioxide (UNII: 15FIX9V2JP)	
hydroxypropyl cellulose ()	
D&C Red No. 7 Calcium lake ()	

Product Characteristics

Color	PURPLE	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	MP;7
Contains			
Coating	true	Symbol	false

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53489-127-01	100 in 1 BOTTLE, PLASTIC		
2	NDC:53489-127-05	500 in 1 BOTTLE, PLASTIC		
3	NDC:53489-127-10	1000 in 1 BOTTLE, PLASTIC		

HYDROXYZINE HYDROCHLORIDE

hydroxyzine hydrochloride tablet, film coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:53489-128
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Hydroxyzine hydrochloride (UNII: 76755771U3) (Hydroxyzine - UNII:30S50YM8OG)		50 mg

Inactive Ingredients

Ingredient Name	Strength
Anhydrous lactose ()	
colloidal silicon dioxide ()	
croscarmellose sodium ()	
FD&C Blue No. 2 Aluminum lake ()	
hypromellose ()	
magnesium stearate (UNII: 70097M6I30)	
polyethylene glycol ()	
titanium dioxide (UNII: 15FIX9V2JP)	
polysorbate 80 ()	
D&C Red No. 7 Calcium lake ()	

Product Characteristics

Color	PURPLE	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	MP;13
Contains			
Coating	true	Symbol	false

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53489-128-01	100 in 1 BOTTLE, PLASTIC		
2	NDC:53489-128-05	500 in 1 BOTTLE, PLASTIC		
3	NDC:53489-128-10	1000 in 1 BOTTLE, PLASTIC		

HYDROXYZINE HYDROCHLORIDE

hydroxyzine hydrochloride tablet, film coated

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:53489-594
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Hydroxyzine hydrochloride (UNII: 76755771U3) (Hydroxyzine - UNII:30S50YM8OG)		25 mg

Inactive Ingredients

Ingredient Name	Strength
Anhydrous lactose ()	
colloidal silicon dioxide ()	
croscarmellose sodium ()	
hypromellose ()	
magnesium stearate (UNII: 70097M6I30)	
polyethylene glycol ()	
titanium dioxide (UNII: 15FIX9V2JP)	
hydroxypropyl cellulose ()	
D&C Red No. 7 Calcium lake ()	

Product Characteristics

Color	PINK	Score	no score
Shape	ROUND	Size	6mm
Flavor		Imprint Code	MP;7

Contains				
Coating	true	Symbol	false	
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53489-594-01	100 in 1 BOTTLE, PLASTIC		
2	NDC:53489-594-05	500 in 1 BOTTLE, PLASTIC		
3	NDC:53489-594-10	1000 in 1 BOTTLE, PLASTIC		

HYDROXYZINE HYDROCHLORIDE

hydroxyzine hydrochloride tablet, film coated

Product Information			
Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:53489-595
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	Hydroxyzine hydrochloride (UNII: 76755771U3) (Hydroxyzine - UNII:30S50YM8OG)		50 mg

Inactive Ingredients	
Ingredient Name	Strength
Anhydrous lactose ()	
colloidal silicon dioxide ()	
croscarmellose sodium ()	
hypromellose ()	
magnesium stearate (UNII: 70097M6I30)	
polyethylene glycol ()	
titanium dioxide (UNII: 15FIX9V2JP)	
polysorbate 80 ()	
D&C Red No. 7 Calcium lake ()	

Product Characteristics			
Color	PINK	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	MP;13
Contains			
Coating	true	Symbol	false

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53489-595-01	100 in 1 BOTTLE, PLASTIC		
2	NDC:53489-595-05	500 in 1 BOTTLE, PLASTIC		
3	NDC:53489-595-10	1000 in 1 BOTTLE, PLASTIC		

Labeler - Mutual Pharmaceutical Company, Inc.

Revised: 1/2007

Mutual Pharmaceutical Company, Inc.