COLD MAX DAY AND NIGHT- acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, and phenylephrine hydrochloride AmeriSource Bergen

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

GNP-1150-2022-0802

COLD MAX DAY

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - cough
 - sinus congestion and pressure
- helps clear nasal passages
- promotes nasal and sinus drainage
- temporarily reduces fever

COLD MAX NIGHT

Drug Facts

Active ingredients (in each caplet)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg	Antihistamine
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - runny nose and sneezing
 - cough
 - sinus congestion and pressure
- helps clear nasal passages
- relieves cough to help you sleep
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

liver disease

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

do not exceed recommended dosage

In addition, when using Cold Max Night:

- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- do not take Day and Night caplets at the same time
- do not take more than a total of 10 caplets in 24 hours

adults and children 12 years and over	 take 2 caplets every 4 hours swallow whole – do not crush, chew, or dissolve
children under 12 years	ask a doctor

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

Cold Max Day

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, crospovidone, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

Cold Max Night

acesulfame potassium, colloidal silicon dioxide, croscarmellose sodium, FD&C blue #1 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide

PRINCIPAL DISPLAY PANEL

Good Neighbor Pharmacy®

NDC 46122-411-62

DAY & NIGHT

Cold Max

Acetaminophen, Phenylephrine HCl, Dextromethorphan HBr, Chlorpheniramine Maleate*
Pain Reliever/Fever Reducer, Nasal Decongestant, Cough Suppressant, Antihistamine*
*Antihistamine in Nighttime only

Day

- Head & Body Aches
- Fever & Sore Throat
- Nasal Congestion
- Cough

COOL TASTE


12 CAPLETS

Night

- Head & Body Aches
- Fever & Sore Throat
- Nasal Congestion
- Cough
- Runny Nose*

COOL TASTE

8 CAPLETS





glycol, polyvinyl alcohol, povidone, pregelatinized starch, propylene glycol, steanc acid, talc, ttanium dioxide crospovidone, FD&C blue#1 aluminum lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene acesuliame potassium, colloidal silicon dioxide, corn starch, croscarmellose sodium, povidone, pregelatinized starch, propylene glycol, stearic acid, talc, titanium dioxide crospovidone, flavor, magnesium stearate, microcrystaline cellulose, polyetnylene glycol, polyvinyl alcohol, acesulfame potassium, colloidal silicon dioxide, com starch, croscarmellose sodium, Cold Max Day Inactive ingredients

nund racts (continued)

Actual Size

 retain carton for complete product information and warnings ■ store between 20-25°C (68-77°F) in a dry place awallow whole; do not crush, chew, or dissolve ■ take 2 caplets every 4 hours ■ do not take more than a total of 10 caplets in 24 hours ■ do not take Day and Night caplets at the same time do not take more than directed (see overdose waming) (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice

any signs or symptoms.

Uther information

children under 12 years 12 years and over

adults and children

DIrections

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away Keep out of reach of children.

It bregnant or breast-feeding, ask a health professional before use.

■ ask a doctor

These could be signs of a serious condition.

 conditionnes pack of occurs with rash of headache that lasts ■ uew symptoms occur ■ tever gets worse or lasts more than 3 days ■ redness or swelling is present

■ pain, nasal congestion, or cough gets worse or lasts more than 7 days

Stop use and ask a doctor if a nervousness, dizziness, or sleeplessness occur

 pe careful when driving a motor vehicle or operating machinery avoid alcoholic drinks alcohol, sedatives, and tranquilizers may increase drowsiness marked drows in ess may occur excuspility may occur, especially in children

> In addition, when using Cold Max Night ■ do not exceed recommended dosage When using this product

 taking the blood thinning drug warfarin taking sedatives or tranquilizers Ask a doctor or pharmacist before use if you are

a breathing problem such as emphysema or chronic bronchits

 beisistent or drionic cough such as occurs with smoking, asthma, or emphysema trouble uninating due to an enlarged prostate gland

 condu that occurs with too much phiegm (mucus) a urgu noou presente a creacin moulin =

DO NOT USE IF BLISTER UNITS ARE TORN OR BROKEN



1 West First Av Conshohocken, P. Questions or Cor Distributed E AmerisourceBe www.mygnp.

ASK a doctor before use if you have

 if you have ever had an allergic reaction to this product or any of its ingredients do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression,

חגת*לו* **גיפור (co**uriuned)



F115002GNP

COLD MAX DAY AND NIGHT

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, and phenylephrine hydrochloride kit

Product Information

HUMAN OTC DRUG NDC:46122-411 **Product Type** Item Code (Source)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-411-62	1 in 1 CARTON	08/01/2018	

Ouantity of Parts

~	• -	
Part #	Package Quantity	Total Product Quantity
Part 1	1 BLISTER PACK	12
Part 2	1 BLISTER PACK	8

Part 1 of 2

ACETAMINOPHEN, DEXTROMETHORPHAN HYDROBROMIDE, PHENYLEPHRINE HYDROCHLORIDE

acetaminophen, dextromethorphan hydrobromide, and phenylephrine hydrochloride tablet, coated

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg		
DEXTROMETHORPHAN HYDROBROMIDE (LINII: 9D2RTI9KYH)	DEXTROMETHORPHAN			

(DEXTROMETHORPHAN - UNII:7355X3ROTS)	HYDROBROMIDE	TO HIG	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1W5297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients		
Ingredient Name	Strength	
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)		
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)		
CROSPOVIDONE (UNII: 2S7830E561)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)		
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)		
STARCH, PREGELATINIZED CORN (UNII: 08232NY3SJ)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
STEARIC ACID (UNII: 4ELV7Z65AP)		
TALC (UNII: 7SEV7J4R1U)		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		

Product Characteristics			
Color	white	Score	no score
Shape	OVAL (capsule-shaped)	Size	17mm
Flavor	MINT	Imprint Code	AAA;1138
Contains			

l	Pā	Packaging			
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341		

Part 2 of 2

ACETAMINOPHEN, CHLORPHENIRAMINE MALEATE, DEXTROMETHORPHAN HYDROBROMIDE, PHENYLEPHRINE HYDROCHLORIDE

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, and phenylephrine

Product Information

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg	
CHLORPHENIRAMINE MALEATE (UNII: V1Q0090J9Z) (CHLORPHENIRAMINE - UNII: 3U6I01965U)	CHLORPHENIRAMINE MALEATE	2 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1W5297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients	
Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics				
Color	blue	Score	no score	
Shape	OVAL (capsule-shaped)	Size	17mm	
Flavor	MINT	Imprint Code	AAA;1139	
Contains				

Pa	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1		8 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC monograph final	part341				
Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
OTC monograph final	part341	08/01/2018			

Labeler - AmeriSource Bergen (007914906)

Revised: 8/2022 AmeriSource Bergen