

**CHILDRENS PAIN AND FEVER- acetaminophen tablet, chewable**  
**Topco Associates, LLC**

-----  
**Topcare 44-450**

***Active ingredient (in each chewable tablet)***

Acetaminophen 160 mg

***Purpose***

Pain reliever/fever reducer

***Uses***

- temporarily relieves minor aches and pains due to:
  - headache
  - sore throat
  - flu
  - toothache
  - the common cold
- temporarily reduces fever

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this

product

**Ask a doctor before use if your child has**

liver disease.

**Ask a doctor or pharmacist before use if your child is**

taking the blood thinning drug warfarin.

**When using this product**

**do not exceed recommended dosage.**

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor

<b>Weight (lb)</b>	<b>Age (yr)</b>	<b>Dose (chewable tablets)*</b>
Under 24	Under 2	Ask a doctor
24-35	2-3	1 tablet
36-47	4-5	1 1/2 tablets
48-59	6-8	2 tablets
60-71	9-10	2 1/2 tablets
72-95	11	3 tablets

\*or as directed by a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid high humidity
- see end flap for expiration date and lot number

**Inactive ingredients**

citric acid, crospovidone, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrates hydrated, ethylcellulose, flavor, magnesium stearate, mannitol, polyethylene, stearic acid, sucralose

**Questions or comments?**

**1-888-423-0139**

**Principal display panel**

**+TopCare®**  
health

NDC 36800-450-08

**COMPARE TO CHILDREN'S  
TYLENOL® ACTIVE INGREDIENT†**

**children's  
Pain & Fever**

**ACETAMINOPHEN** 160 mg, CHEWABLE TABLETS  
PAIN RELIEVER • FEVER REDUCER

• Ibuprofen Free • Aspirin Free

Chew or crush tablets completely before swallowing.

BUBBLE  
GUM  
FLAVOR

**24** CHEWABLE TABLETS

For Ages 2 to 11 Years

actual size

**TAMPER EVIDENT: DO NOT USE IF  
IMPRINTED SAFETY SEAL UNDER  
CAP IS BROKEN OR MISSING**

**QUALITY GUARANTEED**

†This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Children's

Tylenol®. 50844 REV0522B45008

DISTRIBUTED BY TOPCO ASSOCIATES LLC  
ELK GROVE VILLAGE, IL 60007 ©TOPCO LNKA0521

QUESTIONS? 1-888-423-0139

topcare@topco.com www.topcarebrand.com

Visit here for more information: <http://topbrnds.com/48ZZSF>

no print / no varnish area  
lot no. & exp. date



QUALITY GUARANTEED

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING



NDC 36800-450-08  
COMPARE TO CHILDREN'S  
TYLENOL® ACTIVE INGREDIENT†

## children's Pain & Fever

ACETAMINOPHEN 160 mg, CHEWABLE TABLETS  
PAIN RELIEVER • FEVER REDUCER

• Ibuprofen Free • Aspirin Free



BUBBLE GUM FLAVOR

Chew or crush tablets completely before swallowing.

**24 CHEWABLE TABLETS** For Ages 2 to 11 Years actual size



## children's Pain & Fever

ACETAMINOPHEN 160 mg, CHEWABLE TABLETS  
PAIN RELIEVER • FEVER REDUCER

**24 CHEWABLE TABLETS**

**Drug Facts (continued)**

Weight (lb)	Age (yr)	Dose (chewable tablets)*
Under 24	Under 2	Ask a doctor
24-35	2-3	1 Tablet
36-47	4-5	1 1/2 tablets
48-59	6-8	2 tablets
60-71	9-10	2 1/2 tablets
72-95	11	3 tablets

\*or as directed by a doctor

**Other information**

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ avoid high humidity
- see end flap for expiration date and lot number

**Inactive ingredients** citric acid, crospovidone, D&C red #27 aluminum lake, D&C red #30 aluminum lake, dextrates hydrated, ethylcellulose, flavor, magnesium stearate, mannitol, polyethylene, stearic acid, sucralose

**Questions or comments? 1-888-423-0139**

†This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark, Children's Tylenol®. 50844 REV0522B45008

DISTRIBUTED BY TOPCO ASSOCIATES LLC  
ELK GROVE VILLAGE, IL 60007 ©TOPCO LNKA0521  
QUESTIONS? 1-888-423-0139  
topcare@topco.com www.topcarebrand.com  
Visit here for more information: <http://topbrnds.com/48ZZSF>

B-1910-450-08-R  
REV0522B45008

**Drug Facts (continued)**

- if your child is allergic to acetaminophen or any of the inactive ingredients in this product
- Ask a doctor before use if your child has liver disease.
- Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

When using this product do not exceed recommended dosage.

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ this product does not contain directions or complete warnings for adult use

- do not give more than directed
- find the right dose on chart below. If possible, use weight to dose; otherwise, use age.
- chew or crush tablets completely before swallowing; do not swallow tablets whole
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- do not give for more than 5 days unless directed by a doctor

Topcare 44-450

## CHILDRENS PAIN AND FEVER

acetaminophen tablet, chewable

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:36800-450
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
D&C RED NO. 27 ALUMINUM LAKE (UNII: ZK64F7XSTX)	
D&C RED NO. 30 (UNII: 2S42T2808B)	
DEXTROSE MONOHYDRATE (UNII: LX22YL083G)	
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MANNITOL (UNII: 3OWL53L36A)	
HIGH DENSITY POLYETHYLENE (UNII: UG00KM4WR7)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

## Product Characteristics

<b>Color</b>	pink	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	16mm
<b>Flavor</b>	BUBBLE GUM	<b>Imprint Code</b>	44;450
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:36800-450-08	1 in 1 CARTON	02/15/2019	
1		24 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013	02/15/2019	

**Labeler** - Topco Associates, LLC (006935977)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(36800-450)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(36800-450) , pack(36800-450)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(36800-450)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(36800-450)

Revised: 12/2023

Topco Associates, LLC