CVS NIGHTTIME COLD AND FLU RELIEF- acetaminophen, dextromethorphan hbr, doxylamine succinate solution CVS PHARMACY

CVS Nighttime Cold & Flu Relief Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Antihistamine

Uses

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- Skin reddening
- Blisters
- Rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 30 mL contains: sodium 16 mg
- store at room temperature and do not refrigerate

Inactive ingredients

anhydrous citric acid, D&C Yellow No. 10, disodium edetate, FD&C Green No. 3, FD & C Yellow No 6, flavors, propylene glycol, propyl gallate, purified water, sodium benzoate, sodium citrate, sorbitol liquid, sucralose.

Questions or comments?

1-866-467-2748

Package/Label Principal Display Panel

Compare to the active ingredients in VICKS® Nyquil™Cold & Flu*

NDC 51316-692-12

Nighttime Cold & Flue Relief

Acetaminophen, Dextromethorphan succinate,

Doxylamine HBr

Headache, Fever, Sore Throat

Minor Aches & Pains

Sneezing, Runny Nose, Cough

Original Flavor

12 FL OZ (354 mL)

Distributed by:

This product is not manufactured or distributed by Procter & Gamble, the distributor of Vicks[®] Nyquil™Cold & Flu

Compare to the active **CVS** Drug Facts (continued) Drug Facts TAMPER EVIDENT: DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN. ingredients in Vicks® Health. Active ingredients (in each 30 mL) When using this product • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle NyQuil™ Cold & Flu* Purnose Acetaminophen 650 mg..... Dextromethorphan HBr 30 mg... or operating machinery • alcohol, sedatives, and tranquilizers may incre of operand y mature in a mature, southers, and a templace is a five interest or in a five mature. On the five in a f **Multi-Symptom** ...Cough suppressant Doxylamine succinate 12.5 mg. Nighttime Uses temporarily relieves common cold/flu symptoms:

• cough due to minor throat and bronchial irritation • sore throat • headache If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poiso Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms. minor aches and pains • fever • runny nose and sneezing ACETAMINOPHEN-Pain reliever: Fever reducer Warnings
Liver warning. This product contains acetaminophen. Severe liver damage may occur if you take - more than 4 doses in 24 hours, which is the maximum daily amount for this product - with other drugs containing acetaminophen - 3 or more alcoholic drinks every day while using this product
Allerny alert. Acetamin polher may cause severe skin reactions. Symptoms may include: - shin reaction occurs, shop use and seek medical help right away
If a skin reaction occurs, shop use and seek medical help right away
Sore throat warning. If soe throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomitting, consult a doctor promotify. Directions • take only as directed • only use the dose cup provided DEXTROMETHORPHAN HBr-.5" Height do not exceed 4 doses per 24 hrs adults & children 12 yrs & over | 30 mL every 6 hrs DOXYLAMINE SUCCINATEchildren 4 to under 12 yrs ask a doctor the distributor of Vicks® NyQuil* Cold & Flu. Other information • each 30 mL contains: Relieves: Headache, Fever, Sore throat, Minor aches & pains, Sneezing, Runny nose, Cough sodium 16 mg • store at room temperature and do not refrigerate Inactive ingredients citric acid, D&C Yellow No. 10, disodium edetate, FD&C Green No. 3, FD&C Yellow No. 6, flavors propyl galate, progylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol liquid, sucralose. Do not use
"with any other drug containing acetaminophen (prescription or nonprescription).
If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
"A you are now king a prescription monoramine oxides inhibitor (MADI) (setting a drugs for depression, psychatric, or emotional conditions, or Parlinson's disease), or
for 2 verest after shopping the MADI (large, through or the product of the
contains an MADI, ask a doctor or pharmacist before taking this product. ### 1-800 Process

6920RG12CVSIRI Distributed by:
One CVS Pharmacy, Inc.
One CVS Process
O Questions or comments? 1-866-467-2748 COLD & FLU Ask a doctor before use if you have - liver disease • glaucoma • cough that occurs vitt too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema • trouble urinating du to an enlarged prostate gland Original Flavor 100% money back guaranteed. CVS.com/returnpolicy Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers
• taking the blood thinning drug warfarin 12 FL 0Z (354 mL)

7.75" Width

CVS NIGHTTIME COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate solution

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51316-692
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	650 mg in 30 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL	
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL	

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)		

EDETATE DISODIUM (UNII: 7FLD91C86K) FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S) FD&C YELLOW NO. 6 (UNII: H77VEI93A8) PROPYL GALLATE (UNII: 8D4SNN7V92) PROPYLENE GLYCOL (UNII: 6DC9Q167V3) WATER (UNII: 059QF0KO0R) SODIUM BENZOATE (UNII: 0J245FE5EU) SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR) SORBITOL (UNII: 506T60A25R) SUCRALOSE (UNII: 96K6U037D4)		
FD&C YELLOW NO. 6 (UNII: H77VEI93A8) PROPYL GALLATE (UNII: 8D4SNN7V92) PROPYLENE GLYCOL (UNII: 6DC9Q167V3) WATER (UNII: 059QF0KO0R) SODIUM BENZOATE (UNII: OJ245FE5EU) SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR) SORBITOL (UNII: 506T60A25R)	EDETATE DISODIUM (UNII: 7FLD91C86K)	
PROPYL GALLATE (UNII: 8D4SNN7V92) PROPYLENE GLYCOL (UNII: 6DC9Q167V3) WATER (UNII: 059QF0KO0R) SODIUM BENZOATE (UNII: OJ245FE5EU) SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR) SORBITOL (UNII: 506T60A25R)	FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3) WATER (UNII: 059QF0KO0R) SODIUM BENZOATE (UNII: 0J245FE5EU) SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR) SORBITOL (UNII: 506T60A25R)	FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
WATER (UNII: 059QF0KO0R) SODIUM BENZOATE (UNII: OJ245FE5EU) SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR) SORBITOL (UNII: 506T60A25R)	PROPYL GALLATE (UNII: 8D4SNN7V92)	
SODIUM BENZOATE (UNII: OJ245FE5EU) SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR) SORBITOL (UNII: 506T60A25R)	PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR) SORBITOL (UNII: 506T60A25R)	WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	SODIUM BENZOATE (UNII: OJ245FE5EU)	
	SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SUCRALOSE (UNII: 96K6UO3Z D4)	SORBITOL (UNII: 506T60A25R)	
	SUCRALOSE (UNII: 96K6UQ3ZD4)	

I	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:51316-692- 12	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/13/2023	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M010	02/13/2023	

Labeler - CVS PHARMACY (062312574)

Revised: 11/2023 CVS PHARMACY