

PEPTO BISMOL FAST MELTS WINTER FROST FLAVORED- bismuth subsalicylate tablet, chewable
The Procter & Gamble Manufacturing Company

Pepto Bismol Fast Melts Winter Frost

Drug Facts

Active ingredient (in each soft chewable tablet)

Bismuth subsalicylate 262 mg

Purpose

Upset stomach reliever and antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
 - heartburn
 - indigestion
 - nausea
 - gas
 - belching
 - fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use if

you have

- an ulcer
- a bleeding problem

- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- fully chew before swallowing
- adults and children 12 years and over:
 - 2 soft chewable tablets every 1/2 hour or 4 soft chewable tablets every hour as needed for diarrhea
 - 2 soft chewable tablets every 1/2 hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 soft chewable tablets) in 24 hours • use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each soft chewable tablet contains:** • calcium 55 mg • salicylate 100 mg • low sodium
- Store at no greater than 25°C. Avoid excessive heat (over 40°C).

Inactive ingredients

calcium carbonate, carrageenan, corn syrup, FD&C Blue No. 1, FD&C Red No. 40, flavor, hydrogenated coconut oil, soy lecithin, sucrose, water

Questions?**1-800-717-3786****TAMPER EVIDENT: DO NOT USE IF INNER PRINTED SEAL ON BOTTLE IS
BROKEN OR MISSING.****DIST. BY: PROCTER & GAMBLE,
CINCINNATI, OH 45202****PRINCIPAL DISPLAY PANEL - 24 SOFT CHEWABLE TABLETS****Pepto Bismol®**

Bismuth Subsalicylate

Upset Stomach Reliever/Antidiarrheal

FAST MELTS

COATS & COOLS

WINTER FROST FLAVORED

24 SOFT CHEWABLE TABLETS



DISCARD SEAL, EMPTY & REPLACE CAP
 how2recycle.info
PLASTIC BOTTLE
 DIST. BY: PROCTER & GAMBLE, CINCINNATI, OH 45202
 www.pg.com
 pepto-bismol.com
 Patents: www.pg.com/patents

Pepto Bismol

Bismuth Subsalicylate Upset Stomach Reliever/Antidiarrheal




TAMPER EVIDENT: DO NOT USE IF INNER PRINTED SEAL ON BOTTLE IS BROKEN OR MISSING.

Drug Facts	
Active ingredient (in each soft chewable tablet) Bismuth subsalicylate 262 mg.....	Purpose Upset stomach reliever and antidiarrheal
Uses relieves • travelers' diarrhea • diarrhea • upset stomach due to overindulgence in food and drink, including: • heartburn • indigestion • nausea • gas • belching • fullness	
Warnings Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Contains salicylate. Do not take if you are • allergic to salicylates (including aspirin) • taking other salicylate products Do not use if you have • an ulcer • a bleeding problem • bloody or black stool	

21028843

PEEL BACK FOR DRUG FACTS ▶

Drug Facts (continued)	
Ask a doctor before use if you have • fever • mucus in the stool	
Ask a doctor or pharmacist before use if you are taking any drug for • anticoagulation (thinning the blood) • diabetes • gout • arthritis	
When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur	
Stop use and ask a doctor if • symptoms get worse or last more than 2 days • ringing in the ears or loss of hearing occurs • diarrhea lasts more than 2 days	
If pregnant or breast feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions • fully chew before swallowing • adults and children 12 years and over: • 2 soft chewable tablets every ½ hour or 4 soft chewable tablets every hour as needed for diarrhea • 2 soft chewable tablets every ½ hour to 1 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) • do not exceed 8 doses (16 soft chewable tablets) in 24 hours • use until diarrhea stops but not more than 2 days • children under 12 years: ask a doctor • drink plenty of clear fluids to help prevent dehydration caused by diarrhea	
Other information • each soft chewable tablet contains: • calcium 55 mg • salicylate 110 mg • low sodium • Store at no greater than 25°C. Avoid excessive heat (over 40°C).	
Inactive ingredients calcium carbonate, carrageenan, corn syrup, FD&C Blue No. 1, FD&C Red No. 40, flavor, hydrogenated coconut oil, soy lecithin, sucrose, water	
Questions? 1-800-717-3786	



PEPTO BISMOL FAST MELTS WINTER FROST FLAVORED

bismuth subsalicylate tablet, chewable

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69423-832
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)		BISMUTH SUBSALICYLATE	262 mg

Inactive Ingredients

Ingredient Name	Strength
CALCIUM CARBONATE (UNII: H0G9379FGK)	
CARRAGEENAN (UNII: 5C69YCD2YJ)	
CORN SYRUP (UNII: 9G5L16BK6N)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HYDROGENATED COCONUT OIL (UNII: JY81OXM1OM)	
LECITHIN, SOYBEAN (UNII: 1DI56QDM62)	
SUCROSE (UNII: C151H8M554)	
WATER (UNII: 059QF0KOOR)	

Product Characteristics

Color	blue	Score	no score
Shape	ROUND	Size	22mm
Flavor	MINT	Imprint Code	C
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69423-832-24	24 in 1 BOTTLE; Type 0: Not a Combination Product	03/05/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	03/05/2024	

Labeler - The Procter & Gamble Manufacturing Company (004238200)

Revised: 2/2024

The Procter & Gamble Manufacturing Company