ENTERIC COATED ASPIRIN REGULAR STRENGTH- aspirin tablet, delayed release

Strategic Sourcing Services LLC

SMK - 1081B - 2019-0517

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

* nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Uses

- for the temporary relief of minor aches and pains due to:
 - headache
 - muscle pain
 - toothache
 - menstrual pain
 - colds
 - minor pain of arthritis
- or as recommended by your doctor

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance

is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for:

- diabetes
- gout
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- redness or swelling is present
- fever gets worse or lasts for more than 3 days
- new symptoms occur
- ringing in the ears or loss of hearing occurs

These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 to 2 tablets every 4 hours while

symptoms last; do not take more than 12 tablets in 24 hours unless directed by a doctor

• children under 12 years: ask a doctor

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

corn starch, croscarmellose sodium, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, methacrylic acid copolymer, microcrystalline cellulose, mineral oil, polysorbate 80, simethicone, sodium hydroxide, sodium lauryl sulfate, talc, titanium dioxide, triethyl citrate

PRINCIPAL DISPLAY PANEL

sunmark®

COMPARE TO ECOTIN® REGULAR STRENGTH ACTIVE INGREDIENT

NDC 70677-0071-1

enteric coated aspirin


Regular Strength

Pain Reliever (NSAID)

ASPIRIN REGIMEN**

SAFETY COATED

125 TABLETS - 325 MG EACH



ENTERIC COATED ASPIRIN REGULAR STRENGTH

aspirin tablet, delayed release

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70677-0071
Route of Administration	ORAL		

	Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength	
ı	ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg	
ı				

Inactive Ingredients		
	Ingredient Name	Strength

STARCH, CORN (UNII: O8232NY3SJ) CROSCARMELLOSE SODIUM (UNII: M280L1HH48) **D&C YELLOW NO. 10** (UNII: 35SW5USQ3G) **ALUMINUM OXIDE** (UNII: LMI26O6933) FD&C YELLOW NO. 6 (UNII: H77VEI93A8) HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO) METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13) CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) MINERAL OIL (UNII: T5L8T28FGP) POLYSORBATE 80 (UNII: 60ZP39ZG8H) **DIMETHICONE** (UNII: 92RU3N3Y10) **SODIUM HYDROXIDE** (UNII: 55X04QC32I) **SODIUM LAURYL SULFATE** (UNII: 368GB5141J) TALC (UNII: 7SEV7J4R1U) **TITANIUM DIOXIDE** (UNII: 15FIX9V2JP) TRIETHYL CITRATE (UNII: 8Z96QXD6UM)

Product Characteristics				
Color	orange	Score	no score	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	Т	
Contains				

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70677- 0071-1	1 in 1 CARTON	05/17/2019	02/28/2025
1		1 in 1 PACKAGE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	05/17/2019	02/28/2025

Labeler - Strategic Sourcing Services LLC (116956644)

Revised: 10/2023 Strategic Sourcing Services LLC