4364 FIRST AID KIT- 4364 first aid kit Honeywell Safety Products USA, INC

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

0498-4364: First Aid Kit (Triple, Burn Jel, PVP wipes, Sing Rel, BZK wipes, aypanal- SF00003201)

Burn Jel Active ingredient

Lidocaine HCI 2.0%

Burn Jel *Purpose*

External analgesic

Burn Jel *Uses*

• temporarily relieves pain due to minor burns

Burn Jel *Warnings*

For external use only

Do not use

• on large areas of the body, particularly over raw surfaces or blistered areas

When using this product

avoid contact with eyes

Stop use and ask a doctor if

- the condition gets worse
- symptoms persist for more than 7 days
- condition clears up and recurs within a few days

Keep out of reach of children

• If swallowed, get medical help or contact a Poison Control Center right away.

Burn Jel

Directions

- adults and children 2 years of age and older; apply to affected area not more than 3 to 4 times daily
- children under 2 years of age: ask a doctor
- you may report a serious reaction to this product to 800-430-5490

Burn Jel Other information

- store at room temperature
- do not use if opened or torn

Burn Jel Inactive ingredients

carbopol 940, carbopol 1342, diazolidinyl urea, glycerin, melaleuca alternifolia (tea tree) leaf oil, methylparaben, octoxynol-9, propylene glycol, propylparaben, trolamine, water ...

Burn Jel *Questions*

1-800-430-5490

Povidone Iodine Swab Active ingredient

Povidone-iodine solution USP, 10% (equivalent to 1% titratable iodine)

Povidone Iodine Swab *Purpose*

First aid antiseptic

Povidone Iodine Swab *Uses*

first aid to help prevent the risk of infection in minor cuts, scrapes, and burns

Povidone Iodine Swab *Warnings*

For external use only

Do not use

- over large areas of the body
- on individuals who are allergic or sensitive to iodine

Ask a doctor before use if you have

- deep or puncture wounds,
- animal bites
- serious burns

When using this product

do not use longer than one wek unless directed by a doctor

Stop use and ask a doctor if

- conditions persists or gets worse
- irritation and redness develops

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away

Povidone Iodine Swab Directions

Reverse cardboard sleeve, then crush at dot between thumb and forefinger. Allow solution to saturate tip and apply solution to injury.

- clean affected area
- apply to affected area 1 to 3 times daily
- may be covered with a sterile bandage
- discard swab after single use

Povidone Iodine Swab Other information

- store at room temperature away from light
- keep from freezing or excessive heat
- do not use if package is torn or open

Povidone Iodine Swab Inactive ingredients

citric acid, disodium phosphate, nonoxynol-9, sodium hydroxide, water

Povidone Iodine Swab Questions and comments

1-800-430-5490

Aypanal

Active ingredient

Acetaminophen 325 mg

Aypanal *Purpose*

Pain reliever/ fever reducer

Aypanaly

Uses

 temporarily relieves minor aches and pains due to the common cold and headache temporarily reduces fever

Aypanal *Warnings*

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- child takes more than 5 doses in 24 hours, which is the maximum daily amount
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product:

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- If a skin rash occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription).
- If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if

you are taking the blood thinning drug warfarin

Stop using and ask a doctor if

- pain gets worse or lasts more than 10 days in adults
- pain gets worse or lasts more than 5 days in children under 12 years
- fever gets worse or lasts more than 3 days

- redness or swelling is present
- new symptoms occur

If pregnant or breast-feeding

If pregnant or breast-feeding, ask a health professional before use.

Overdose warning

- In case of accidental overdose, get medical help or contact a Poison Control Center right away.
- Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Keep out of reach of children.

Keep out of reach of children.

Aypanal Directions

do not take more than directed (see overdose warning)

adults and children 12 years of age or older

- take two tablets every 4-6 hours while symptoms last
- do not take more than 12 tablets in 24 hours

children 6 to under 12 years of age

- take 1 tablet every 4-6 hours while symptoms last
- do not take more than 5 tablets in 24 hours

children under 6 years consult a doctor

Aypanal

Other information

- store at room temperature 15 0 to 30 0 C (59 0 86 0 F)
- TAMPER EVIDENT PACKETS- DO NOT USE IF OPEN OR TORN

Aypanal Inactive ingredients

corn starch, microcrystalline cellulose, povidone, sodium starch glycolate, stearic acid

Aypanal Questions

1-800-430-5490

Triple

Active ingredients (eeach gram contains)

Bacitracin zinc 400 units

Neomycin sulfate (5 mg equivalent to 3.5 mg Neomycin base)

Polymyxin B sulfate 5000 units

Triple *Purpose*

First aid antibiotic

First aid antibiotic

First aid antibiotic

Triple

Uses

- first aid to help prevent infection in
- minor cuts
- scrapes
- burns

Triple Warnings

For external use only

Allergy alert do not use if you are allergic to any of the ingredients

Do not use

- in the eyes
- over large areas of the body

Ask a doctor before use if you have

- a deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- the condition persists or gets worse
- a rash or other allergic reaction develops
- you need to use longer than 1 week

Keep out of the reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Triple

Directions

- clean the affected area
- apply a small amount of the product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily
- may be covered with a sterile bandage

Triple

Other information

- store at 15 0 to 25 0 C (59 0 to 77 0 F)
- tamper evident sealed packets
- do not use if packet is torn or opened

Triple

Inactive ingredient

petrolatum

Triple *Questions?*

1-800-430-5490

Sting Relief Active ingredient (in each wipe)

Ethyl alcohol 50.0%

Lidocaine HCl 2.0%

Sting Relief Purpose

Antiseptic

Topical pain relief

Sting Relief

Uses

• prevent infection in minor scrapes, and temporary relief of itching of insect bites

Sting Relief Warnings

For external use only

Flammable, keep away from open fire or flame

Do not use

- over large areas of the body
- in eyes
- over raw or blistered areas

Stop use and ask a doctor

 if conditions worsen or persist for more than 7 days or clear up and occur again within a few days

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Sting Relief

Directions

- adults and children 2 years and older: Apply to cleaned affected area not more than 3 times daily.
- children under 2 years of age: consult a doctor.

Sting Relief Inactive ingredients

benzalkonium chloride, menthol, and purified water

Sting Relief Questions or Comments?

1-800-430-5490

BZK Active ingredient

Benzalkonium chloride 0.13% w/v

BZK *Purpose*

First aid antiseptic

BZK Uses

Antiseptic cleansing of face, hands, and body without soap and water

BZK

Warnings

For external use only

Do not use

- in the eyes or over large areas of the body
- on mucous membranes
- on irritated skin
- in case of deep puncture wounds, animal bites or serious burns, consult a doctor
- longer than 1 week unless directed by a doctor

Stop use and ask a doctor if

- if irritation, redness or other symptoms develop
- the condition persists or gets worse

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away

BZK

Directions

tear open packet and use as a washcloth

BZK

Other information

- store at room temperature 15 0 to 30 0 C (59 0 86 0 F)
- do not reuse towelette

BZK

Inactive ingredients

water

BZK Ouestions

1-800-430-5490


4364 SF00003201 Kit Contents

- 1 TRIPLE ANTIBIOTIC 10 PER
- 1 EYE DRESS PKT W/4 ADH STRIPS
- 2 TRIANGULAR BDG, NON-STERILE
- 1 WIRE SPLINT 1 PER


- 1 GAUZE PADS, 3" X 3", 4 PER
- 1 ADH TAPE, .5" X 2.5 YD, 2 PER
- 1 AYPANAL 26 PER
- 1 GAUZE BANDAGE, 2" X 6 YD,2 PER
- 1 INSTANT COLD PACK 4" X 6"
- 1 ADHESIVE BDG, PLSTIC, 1"X3"16PER
- 1 FINGERTIP BANDAGE, 10 PER
- 1 BURN JEL 1/8 OZ, 6 PER
- 1 PVP IODINE WIPES 10 PER
- 1 STING RELIEF WIPES 10 PER BOX
- 1 ANTIMCRBL ANTSPTC TWLETTS
- 1 FIRST AID GUIDE ASHI
- 1 SCISSOR BDGE 4" RED PLS HDL
- 1 KIT TWEEZER 3 1/2" SLANTED
- 2 PR LRG NITRILE GLVES ZIP BAG
- 1 KIT STL 24 UN WHITE 01

Burn Jel Principal Display Panel






Principal Display Panel



Aypanal Principal Display Panel



Triple Principal Display Panel




Sting Relief Principal Display Panel



BZK Principal Display Panel

well	
02-16-3	5MD
Antiseptic Towele	ettes
Benzalkonium c First aid am	
Six-Saturated Tow	velettes
ty nc. 2917	



4364 Kit Label SF00003201





Distributed by Honeywell Safety Products USA, Inc. Smithfield, RI 02917

4364 FIRST AID KIT

4364 first aid kit kit

D			
Uroa	IIICT I	INTORM STIAN	1
PIUU		Information	

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0498-4364

	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
ı		NDC 0400 4364	1 : 1 KT T		

NDC:0498-4364- 1 in 1 KIT; Type 0: Not a Combination

Quant	Quantity of Parts				
Part # Package Quantity		Total Product Quantity			
Part 1	1 PACKET	1.4 mL			
Part 2	10 POUCH	4 mL			
Part 3	10 PACKET	9 g			
Part 4	10 POUCH	3 mL			
Part 5	6 PACKET	21 g			
Part 6	26 PACKET	52			

Part 1 of 6

ANTISEPTIC TOWELETTE

Product

benzalkonium chloride liquid

Product Information

Item Code (Source)	NDC:0498-0501
Route of Administration	TOPICAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)	BENZ ALKONIUM CHLORIDE	1.3 mg in 1 mL	

Inactive Ingredients		
	Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)		

l	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
		NDC:0498-0501- 00	1.4 mL in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information			
Marketing Application Number or Monograph Category Citation		Marketing Start Date	Marketing End Date
unapproved drug other		12/22/2017	

Part 2 of 6

STING RELIEF PAD

ethyl alcohol, lidocaine swab

Product Information

Item Code (Source) NDC:0498-0733

Route of Administration TOPICAL

Active Ingredient/Active Moiety Ingredient Name Basis of Strength ALCOHOL (UNII: 3K9958V90M) (ALCOHOL - UNII:3K9958V90M) LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNIDOCAINE HYDROCHLORIDE ANHYDROUS LIDOCAINE HYDROCHLORIDE in 1 mL

Inactive Ingredients			
Ingredient Name	Strength		
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)			
MENTHOL (UNII: L7T10EIP3A)			
WATER (UNII: 059QF0KO0R)			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0498-0733- 00	0.4 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		12/23/2017	

Part 3 of 6

TRIPLE ANTIBIOTIC

bacitracin zinc, polymyxin b sulfate, neomycin sulfate ointment

Product Information		
Item Code (Source)	NDC:0498-0750	
Route of Administration	TOPICAL	

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RW052I)	BACITRACIN	400 [iU] in 1 g
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII: J2VZ 07J96K)	POLYMYXIN B	5000 [iU] in 1 g
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:116QD7X297)	NEOMYCIN	3.5 mg in 1 g

Inactive Ingredients	
Ingredient Name	Strength
PETROLATUM (UNII: 4T6H12BN9U)	

Product Characteristics			
Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

l	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:0498-0750- 35	0.9 g in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/19/2018	

Part 4 of 6

PVP IODINE WIPE

povidone-iodine 10% swab

Product Information

Item Code (Source)	NDC:0498-0121
Route of Administration	TOPICAL

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
POVIDONE-IODINE (UNII: 85H0HZU99M) (IODINE - UNII:9679TC07X4)	IODINE	10 mg in 1 mL

Inactive Ingredients		
Ingredient Name	Strength	
NONOXYNOL-9 (UNII: 48Q180SH9T)		
WATER (UNII: 059QF0KO0R)		

	Packaging			
7	# Item Code	Package Description	Marketing Start Date	Marketing End Date
:	NDC:0498-0121-	0.3 mL in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2018	

Part 5 of 6

BURN JEL

gel for burns gel

Product Information		
Item Code (Source)	NDC:0498-0203	
Route of Administration	TOPICAL	

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
LIDOCAINE HYDROCHLORIDE (UNII: V13007Z41A) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	2 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
TROLAMINE (UNII: 903K93S3TK)	
CARBOMER HOMOPOLYMER TYPE C (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: 4Q93RCW27E)	
CARBOMER HOMOPOLYMER TYPE B (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: HHT01ZNK31)	
DIAZOLIDINYL UREA (UNII: H5RIZ3MPW4)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
OCTOXYNOL-9 (UNII: 7JPC6Y25QS)	
DIPROPYLENE GLYCOL (UNII: E107L85C40)	
TEA TREE OIL (UNII: VIF565UC2G)	
GLYCERIN (UNII: PDC6A3C0OX)	
METHYLPARABEN (UNII: A2I8C7HI9T)	

ı	P	ackaging			
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:0498-0203- 00	3.5 g in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information				
Marketing Category	Marketing End Date			
unapproved drug other		09/19/2018		

Part 6 of 6

AYPANAL NON-ASPIRIN

acetaminophen tablet

Product Information		
Item Code (Source)	NDC:0498-2001	
Route of Administration	ORAL	

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg		

ı	Inactive Ingredients	
l	Ingredient Name	Strength

STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
POVIDONE (UNII: FZ 989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	

Product Characteristics				
Color	white	Score	2 pieces	
Shape	ROUND	Size	10mm	
Flavor		Imprint Code	circle;U	
Contains				

Pac	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1		2 in 1 PACKET; Type 0: Not a Combination Product			

Marketing Information				
Marketing Application Number or Monograph Category Citation		Marketing Start Date	Marketing End Date	
unapproved drug other		04/10/2012		

Marketing Information				
Marketing Application Number or Monograph Category Citation		Marketing Start Date	Marketing End Date	
unapproved drug other		10/18/2018		

Labeler - Honeywell Safety Products USA, INC (118768815)

Revised: 1/2024 Honeywell Safety Products USA, INC