

**IBUPROFEN- ibuprofen capsule, liquid filled**  
**L.N.K. International, Inc.**

-----

**Convenience Valet 44-754**

**Active ingredient (in each capsule)**

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)\*  
(present as the free acid and potassium salt)

\*nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
  - minor pain of arthritis
  - headache
  - the common cold
  - backache
  - menstrual cramps
  - toothache
  - muscular aches
- temporarily reduces fever

**Warnings**

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- facial swelling
- blisters
- shock
- hives
- skin reddening
- asthma (wheezing)
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding.

The chance is higher if you

- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product
- are age 60 or older
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers

- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke

**Ask a doctor or pharmacist before use if you are**

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - have bloody or black stools
  - feel faint
  - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - leg swelling
  - slurred speech
  - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

**Other information**

- **each capsule contains:** potassium 20 mg
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F). Protect from light.

**Inactive ingredients**

FD&C blue #1, gelatin, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitan, sorbitol

**Questions or comments?**

**1-877-507-6516** (M-F, 8AM – 4:30PM CST)

**Principal display panel**

24/7 life  
by 7-Eleven™

**Ibuprofen**  
**Solubilized Ibuprofen Capsules,**  
200 mg  
Pain Reliever/Fever Reducer **(NSAID)**

compare to **Advil® Liqui-Gels®**  
active ingredient\*\*

**20**  
LIQUID-FILLED  
CAPSULES

Actual Size

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED  
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

\*\*This product is not manufactured or distributed  
by GlaxoSmithKline Consumer Healthcare  
Holdings (US) LLC, owner of the registered  
trademark Advil® Liqui-Gels®.  
(Liqui-Gels® is a registered trademark of Catalent  
Pharma Solutions, Inc.)

50844 REV0123A75409  
Distributed by: 7-Eleven, Inc.  
Irving, TX 75063 www.7-ELEVEN.com

**Satisfaction Guaranteed 1-800-255-0711**



**Convenience Valet 44-754**

<b>IBUPROFEN</b>		
ibuprofen capsule, liquid filled		
<b>Product Information</b>		
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b> NDC:50844-754
<b>Route of Administration</b>	ORAL	
<b>Active Ingredient/Active Moiety</b>		
Ingredient Name	Basis of Strength	Strength
<b>IBUPROFEN</b> (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg
<b>Inactive Ingredients</b>		
Ingredient Name	Strength	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)		
<b>GELATIN, UNSPECIFIED</b> (UNII: 2G86QN327L)		
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)		
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)		
<b>WATER</b> (UNII: 059QF0K00R)		
<b>SORBITAN</b> (UNII: 6O92ICV9RU)		
<b>SORBITOL</b> (UNII: 506T60A25R)		

**Product Characteristics**

<b>Color</b>	blue	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	IB1
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50844-754-09	1 in 1 CARTON	04/08/2020	
1		20 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078682	04/08/2020	

**Labeler** - L.N.K. International, Inc. (038154464)**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(50844-754)

**Establishment**

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(50844-754)

Revised: 2/2024

L.N.K. International, Inc.