

ACETAMINOPHEN- acetaminophen tablet
Southern Sales & Service, Inc.

Acetaminophen

Drug Facts

Active Ingredient (in each Tablet)

Acetaminophen 325 mg

Purpose

Pain reliever/fever reducer

Uses

for the temporary relief of minor aches and pains due to:

Headache

Muscular aches

Backache

Toothache

Premenstrual and menstrual cramps

Temporarily reduces fever.

• Liver warning *Warnings*

this product contains Acetaminophen. Severe liver damage may occur if

- adult takes more than 12 tablets in 24 hours, which is the maximum daily amount
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription)
- you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if the user has

liver disease

Ask a doctor or pharmacist before use if the user is

taking the blood thinning drug warfarin

Stop use and ask a doctor if

Pain gets worse or lasts more than 10 days in adults.

Pain gets worse or lasts more than 5 days in children under 12 years

Fever gets worse or last more than 3 days

new symptoms occur

Redness or swelling is present. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a doctor before use.

Keep out of the reach of children.

Overdose warning:

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

- Do not take more than directed (see overdose warning)

Directions

Directions

Adults and Children 12 years and older	Take 2 tablets every 4 to 6 hours while symptoms last Do not take more than 12 tablets in 24 hours
Children 6 to 11 years	Take 1 tablet every 4 to 6 hours while symptoms last Do not take more than 5 tablets in 24 hours
Children under 6 years	Do not use regular Strength products in children under 6 years of age; this will provide more than the recommended dose (overdose) of acetaminophen and may cause liver damage.

Other information

store at room temperature 20°-25°C (68-77°F)

Inactive Ingredients

Povidone, Pregelatinized Starch, Sodium Starch Glycolate, Stearic Acid

Acetaminophen 100 tablet

<p>Continuation of the Drug Facts</p> <p>Stop use and ask a doctor if: • Pain gets worse or lasts more than 10 days in adults. • Pain gets worse or lasts more than 5 days in children under 12 years • Fever gets worse or lasts more than 3 days • New symptoms occur • Redness or swelling is present. These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a doctor before use.</p> <p>Keep out of the reach of children.</p> <p>Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. • Do not take more than directed (see overdose warning)</p> <p>Directions</p> <p>Adults and Children 12 years and older</p> <ul style="list-style-type: none"> • Take 2 tablets every 4 to 6 hours while symptoms last • Do not take more than 12 tablets in 24 hours <p>Children 6 to 11 years</p> <ul style="list-style-type: none"> • Take 1 tablet every 4 to 6 hours while symptoms last • Do not take more than 5 tablets in 24 hours <p>Children under 6 years</p> <ul style="list-style-type: none"> • Do not use regular Strength products in children under 6 years of age; this will provide more than the recommended dose (overdose) of acetaminophen and may cause liver damage. <p>Other information: store at room temperature 20°-25° C (68°-77° F)</p> <p>Inactive ingredients: Povidone, Pregelatinized Starch, Sodium Starch Glycolate, Stearic Acid</p>		<p>Distributed by LEON MEDICAL CENTERS, Inc. Miami, FL 33166</p> <p>8 50011 55723 5</p> <p>Pain reliever • Fever Reducer 100 Tablets</p> <p>ACETAMINOPHEN 325 mg</p> <p>LEON MEDICAL CENTERS</p> <p>NDC: 69822-692-10</p> <p>Drug Facts</p> <p>Active ingredient (in each Tablet) Purpose Acetaminophen 325 mg... Pain reliever/fever reducer</p> <p>Uses for the temporary relief of minor aches and pains due to: • Headache • Muscular aches • Backache • Toothache • Premenstrual and menstrual cramps • Temporarily reduces fever.</p> <p>Warnings • Liver warning this product contains Acetaminophen. Severe liver damage may occur if • adult takes more than 12 tablets in 24 hours, which is the maximum daily amount • child takes more than 5 doses in 24 hours • taken with other drugs containing acetaminophen • adult has 3 or more alcoholic drinks every day while using this product</p> <p>Do not use • with any other drug containing acetaminophen (prescription or nonprescription) • if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist • if you are allergic to acetaminophen or any of the inactive ingredients in this product</p> <p>Ask a doctor before use if the user has liver disease Ask a doctor or pharmacist before use if the user is taking the blood thinning drug warfarin</p> <p>Lowest Used by:</p> <p>DRUG FACTS CONTINUED ON SIDE PANEL</p>
---	--	---

ACETAMINOPHEN

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69822-692(NDC:0536-1327)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE (UNII: FZ989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	PH020
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69822-692-10	325 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/01/2019	

Labeler - Southern Sales & Service, Inc. (013114906)**Registrant** - Southern Sales & Service, Inc. (013114906)**Establishment**

Name	Address	ID/FEI	Business Operations
Southern Sales & Service, Inc.		013114906	repack(69822-692)

Revised: 11/2023

Southern Sales & Service, Inc.