

**BISMUTH- bismuth subsalicylate suspension
MWI/VetOne**

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click [here](#).

VETone® Bismuth

**NDC 13985-103-21
VET one
Bismuth
Suspension
Anti-Diarrheal Liquid**

For animal use only.

Keep out of reach of children.

V1 510045

Net Contents: 1 Gallon (3.785L)

CONTAINS:

Bismuth Subsalicylate ... 1.75%

In a palatable aqueous suspension. Flavoring and coloring added.

INDICATIONS:

A palatable oral suspension for use as an aid in controlling simple diarrhea in cattle, horses, dogs and cats

CAUTION:

This product contains salicylate; do not administer with other salicylate-containing products, such as aspirin.

DOSAGE & ADMINISTRATION:

Shake well before using. Administer orally.

Dogs and Cats: 1 to 3 tablespoonfuls every 1-3 hours

Calves and Foals: 3 to 4 oz. every 2-3 hours

Cattle and Horses: 6 to 10 oz. every 2-3 hours

WARNING:

If diarrhea persists after using this product for 2 days, contact a veterinarian.

FOR ORAL USE ONLY. SHAKE WELL BEFORE USING.

STORAGE:

Store in a dry place at room temperature not above 37°C (98.6°F). Protect from freezing.

TAKE TIME OBSERVE LABEL DIRECTIONS

MADE IN USA Rev. 07/12


Distributed by: MWI

Boise, ID 83705

(888) 694-8381

www.vetone.net

NDC 13985-103-21
Gallon



Bismuth

Suspension

CONTAINS:
Bismuth Subsalicylate 1.75%
In a palatable aqueous suspension. Flavoring and coloring added.

INDICATIONS: A palatable oral suspension for use as an aid in controlling simple diarrhea in cattle, horses, dogs and cats.

CAUTION: This product contains salicylate; do not administer with other salicylate-containing products, such as aspirin.

DOSAGE & ADMINISTRATION: Shake well before using. Administer orally.

Dogs and Cats: 1-3 tablespoonfuls every 1-3 hours
Calves and Foals: 3-4 oz. every 2-3 hours
Cattle and Horses: 6-10 oz. every 2-3 hours

Anti-Diarrheal Liquid

For animal use only.
Keep out of reach of children.



V1 510045

Net Contents: 1 Gallon (3.785 L)

WARNING: If diarrhea persists after using this product for 2 days, contact a veterinarian.

FOR ORAL USE ONLY. SHAKE WELL BEFORE USING.

STORAGE: Store in a dry place at room temperature not above 37°C (98.6°F). Protect from freezing.

TAKE TIME  OBSERVE LABEL DIRECTIONS

MADE IN USA  Rev. 07/12



3 13985 00998 3

Distributed by: MWI
Boise, ID 83705
(888) 694-8381
www.vetone.net

Bottle Label
Bottle Label

BISMUTH

bismuth subsalicylate suspension

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:13985-103
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE)	BISMUTH SUBSALICYLATE	18 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GUAR GUM (UNII: E89I1637KE)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	

SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
METHYL SALICYLATE (UNII: LAV5U5022Y)	
WATER (UNII: 059QF0K00R)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:13985-103-21	3785 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
Unapproved drug other		11/17/2010	

Labeler - MWI/VetOne (019926120)

Revised: 9/2013

MWI/VetOne