

**MINI NICOTINE- nicotine polacrilex lozenge**  
**Rite Aid Corporation**

-----  
**Drug Facts**

**Active ingredient (in each lozenge)**

Nicotine polacrilex 4 mg

**Purpose**

Stop smoking aid

**Use**

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**

**If you are pregnant or breast-feeding,**

**only use this medicine on the advice of your health care provider.** Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth problems occur
- persistent indigestion or severe sore throat occurs
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness,

diarrhea, weakness and rapid heartbeat

- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

### **Keep out of reach of children and pets.**

Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### **Directions (4 mg lozenge)**

- **if you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the lozenge on your quit day
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine lozenge
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

- **nicotine lozenge is a medicine and must be used a certain way to get the best results**
- place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. **Do not chew or swallow lozenge.**
- you may feel a warm or tingling sensation
- occasionally move the lozenge from one side of your mouth to the other until completely dissolved
- do not eat or drink 15 minutes before using or while the lozenge is in your mouth
- to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks
- do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects
- **do not use more than 5 lozenges in 6 hours. Do not use more than 20 lozenges per day.**
- It is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

### **Other information**

- Phenylketonurics: Contains Phenylalanine 2.8 mg per lozenge
- store at 20-25°C (68-77°F)
- keep bottle tightly closed and protect from light

## ***Inactive ingredients***

aspartame, calcium polycarbophil, flavor, maltodextrin, mannitol, sodium alginate, sodium bicarbonate, sodium stearyl fumarate, talc, xanthan gum

## ***Questions or comments?***

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

## **Principal Display Panel (4 mg lozenge)**

Compare to the active ingredient in Nicorette® mini Lozenge\*

FREE | SUGAR

FROM | FREE

## **Mini Nicotine Lozenge**

Nicotine Polacrilex 4 mg

Stop Smoking Aid

Mint Flavor

## **FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.**

If you smoke your first cigarette **MORE THAN 30 MINUTES** after waking up, use Nicotine Polacrilex 2 mg Lozenge.

**LOZENGES**, 4 mg Each (Vials)

## **TO INCREASE YOUR SUCCESS IN QUITTING:**

- 1.** You must be motivated to quit.
- 2. Use Enough** - Use **at least 9** Nicotine mini lozenges per day during the first six weeks.
- 3. Use Long Enough** - Use Nicotine mini lozenges for the full 12 weeks.
- 4. Use With a Support Program** as directed in the enclosed User's Guide.
  - **not for sale to those under 18 years of age**
  - **proof of age required**
  - **not for sale in vending machines or from any source where proof of age cannot be verified**

\*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® *mini* Lozenge.

**See inside for more details or visit [www.nitthehabit.com](http://www.nitthehabit.com)**

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**RETAIN OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT**

# INFORMATION.

## DISTRIBUTED BY:

RITE AID, 30 HUNTER LANE,  
CAMP HILL, PA 17011

[www.riteaid.com](http://www.riteaid.com)

## Package Label (4 mg lozenge)

**Drug Facts**

**Active ingredient (in each lozenge)** Nicotine polacrilex 4 mg. **Purpose** Stop smoking aid

**Use**  
reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**  
If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have:**  
heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.  
high blood pressure not controlled with medication. Nicotine can increase your blood pressure.  
stomach ulcer or diabetes  
history of seizures

**Ask a doctor or pharmacist before use if you are:**  
using a non-nicotine stop smoking drug  
taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if:**  
mouth problems occur  
persistent indigestion or severe sore throat occurs  
irregular heartbeat or palpitations occur  
you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness and rapid heartbeat  
you have symptoms of an allergic reaction (such as difficulty breathing or rash)

**Keep out of reach of children and pets.** Nicotine lozenges may have enough nicotine to make children and pets sick. If you need to remove the lozenge, wrap it in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**  
If you are under 18 years of age, ask a doctor before use. No studies have been done to show if this product will work for you.  
before using this product, read the enclosed User's Guide for complete directions and other important information  
begin using the lozenge on your quit day  
if you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine lozenge  
if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine lozenge according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 lozenge every 1 to 2 hours	1 lozenge every 2 to 4 hours	1 lozenge every 4 to 8 hours

nicotine lozenge is a medicine and must be used a certain way to get the best results  
place the lozenge in your mouth and allow the lozenge to slowly dissolve. Minimize swallowing. Do not chew or swallow lozenge.  
you may feel a warm or tingling sensation

**Drug Facts (continued)**

occasionally move the lozenge from one side of your mouth to the other until completely dissolved  
do not eat or drink 15 minutes before using or while the lozenge is in your mouth  
to improve your chances of quitting, use at least 9 lozenges per day for the first 6 weeks  
do not use more than one lozenge at a time or continuously use one lozenge after another since this may cause you hiccups, heartburn, nausea or other side effects  
do not use more than 9 lozenges in 6 hours. Do not use more than 20 lozenges per day.  
it is important to complete treatment. If you feel you need to use the lozenge for a longer period to keep from smoking, talk to your health care provider.

**Other information**  
Phenylethanolamine: Contains Phenylethanolamine 2.8 mg per lozenge  
store at 20° to 25°C (68° to 77°F)  
keep bottle tightly closed and protect from light

**Inactive ingredients** aspartame, calcium polycarbophil, flavor, malto-dextrin, mannitol, sodium alginate, sodium bicarbonate, sodium stearoyl fumarate, talc, xanthan gum

**Questions or comments?**  
Call 1-877-733-3838 Monday-Friday 9AM-5PM EST

\*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® mini Lozenges.

**CAUTION: DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING.**

**RETAIN OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

not for sale to those under 18 years of age  
proof of age required  
not for sale in vending machines or from any source where proof of age cannot be verified

FREE FROM SUGAR FREE

NDC 11822-3622-5

Compare to the active ingredient in Nicorette® mini Lozenge\*

**OUR PROMISE:**  
Sugar Free

**Opening Directions:**  
Push down on cap and turn to open.

**TO INCREASE YOUR SUCCESS IN QUITTING:**  
1. You must be motivated to quit.  
2. Use Enough - Use at least 9 Nicotine mini Lozenges per day during the first six weeks.  
3. Use Long Enough - Use Nicotine mini Lozenges for the full 12 weeks.  
4. Use With a Support Program as directed in the enclosed User's Guide.

**mini Nicotine Lozenges**  
Nicotine Polacrilex 4 mg Stop Smoking Aid

**4 mg**

**FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP.**  
If you smoke your first cigarette MORE THAN 30 MINUTES after waking up, use Nicotine Polacrilex 2 mg Lozenge.

Includes User's Guide

**81**  
ACTUAL SIZE  
MINT FLAVOR  
LOZENGES, 4 mg EACH  
3 BOTTLES OF 27

**DISTRIBUTED BY:**  
RITE AID, 30 HUNTER LANE,  
CAMP HILL, PA 17011  
[www.riteaid.com](http://www.riteaid.com)

**SATISFACTION GUARANTEE**  
If you're not satisfied, we'll happily refund your money.

PLD-C215E F0005514

0 11822 33153 1

Lot No:  
Exp Date:

## RITE AID Mini Nicotine Lozenges Mint Flavor

## MINI NICOTINE

nicotine polacrilex lozenge

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-3622
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>NICOTINE</b> (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	4 mg

## Inactive Ingredients

Ingredient Name	Strength
ASPARTAME (UNII: Z0H242BBR1)	
CALCIUM POLYCARBOPHIL (UNII: 8F049NKY49)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MANNITOL (UNII: 3OWL53L36A)	
SODIUM ALGINATE (UNII: C269C4G2ZQ)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)	
TALC (UNII: 7SEV7J4R1U)	
XANTHAN GUM (UNII: TTV12P4NEE)	

## Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	10mm
Flavor	MINT	Imprint Code	131
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-3622-5	3 in 1 BOX	12/27/2019	
1		27 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA207868	12/27/2019	

**Labeler** - Rite Aid Corporation (014578892)