

# **3M AVAGARD- chlorhexidine gluconate and alcohol lotion**

## **Solventum US OpCo LLC**

-----

### **Drug Facts**

#### **Active ingredients**

Chlorhexidine gluconate 1% w/w

Ethyl alcohol 61% w/w

#### **Purpose**

Antiseptic

Antiseptic

#### **Uses**

- **surgical hand antiseptic:** significantly reduces the number of micro-organisms on the hands and forearms prior to surgery or patient care
- **healthcare personnel hand antiseptic:** hand rub to help reduce bacteria that potentially can cause disease
- recommended for repeated use

#### **Warnings**

**For external use only. Flammable, keep away from fire or flames.**

**Allergy alert:** This product may cause a severe allergic reaction. Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use** if you are allergic to chlorhexidine gluconate or any other ingredient in this product

#### **When using this product**

- do not touch the eye with hands that have been treated with this preparation
- keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye.
- if contact occurs, rinse with cold water right away
- do not use routinely if you have wounds which involve more than the superficial layers of the skin

**Stop use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

### **Directions**

- use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

### **Surgical hand antiseptic**

- apply to clean, dry hands and nails. For the first use of each day, clean under nails with a nail stick.
- dispense one pump (2 mL) into the palm of one hand
- dip the fingertips of the opposite hand into the lotion and work it under the nails
- spread the remaining lotion over the hand and up to just above the elbow
- using another 2 mL of lotion, repeat with the other hand
- dispense another 2 mL of lotion into either hand; reapply to all aspects of both hands up to the wrists
- allow to dry before donning gloves

### **Healthcare personnel hand antiseptic**

- apply to clean, dry hands and nails
- dispense one pump (2 mL) into the palm of one hand
- paying particular attention to the spaces between the fingers and under fingernails, apply the lotion evenly to cover both hands up to the wrists
- allow to dry without wiping

### **Other information**

store at 20-25°C (68-77°F)

### **Inactive ingredients**

beheneth 10, behenyl alcohol, C20-40 pareth-24, cetyl palmitate, diisopropyl dimer dilinoleate, dimethicone, glycerin, polyethylene glycol, squalane, water

### **Questions?**

Call 1-800-228-3957

### **Principle Display Panel - 500mL Bottle Label**

**3M**

NDC 17518-051-01

**Avagard™**

**(Chlorhexidine Gluconate 1% Solution and Ethyl Alcohol 61% w/w)**  
**Surgical and Healthcare Personnel Hand Antiseptic with Moisturizers**

Peel to read Drug Facts Panel before use

Made in U.S.A. for **3M Health Care**

2510 Conway Ave., St. Paul, MN 55144

1-800-228-3957

[www.3M.com/infectionprevention](http://www.3M.com/infectionprevention)

© 2017, 3M. All rights reserved. The shape and colors of the bottle and wall bracket are trademarks of 3M.

Patent: [3M.com/Patents](http://3M.com/Patents)

MAL10110039XC

Flammable, keep away from fire or flame, heat, sparks and sources of static discharge.


LOT 0000000000

EXP 0000-00-00

**16 fl oz • 500 mL**

**REF 9200**





## 3M AVAGARD

chlorhexidine gluconate and alcohol lotion

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:17518-051
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Chlorhexidine Gluconate</b> (UNII: MOR84MUD8E) (Chlorhexidine - UNII:R4KO0DY52L)	Chlorhexidine Gluconate	8.3 mg in 1 mL
<b>Alcohol</b> (UNII: 3K9958V90M) (Alcohol - UNII:3K9958V90M)	Alcohol	506.3 mg in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>Beheneth-10</b> (UNII: 313S43DM16)	
<b>Docosanol</b> (UNII: 9G10E216XY)	
<b>Cetyl Palmitate</b> (UNII: 5ZA2S6B08X)	
<b>Dimethicone</b> (UNII: 92RU3N3Y10)	
<b>Glycerin</b> (UNII: PDC6A3C0OX)	

polyethylene glycol, unspecified (UNII: 3WJQ0SDW1A)				
Squalane (UNII: GW89575KF9)				
Water (UNII: 059QF0KO0R)				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17518-051-01	8 in 1 CASE	06/14/2001	
1		500 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:17518-051-04	4 in 1 CASE	06/14/2001	
2		1200 mL in 1 BOTTLE; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA		NDA021074	06/14/2001	

**Labeler -** Solventum US OpCo LLC (006173082)

<b>Establishment</b>			
Name	Address	ID/FEI	Business Operations
Grain Processing Corporation		198484180	API MANUFACTURE(17518-051)

Revised: 4/2024

Solventum US OpCo LLC