

**ULTRA-SOFT BARRIER- bronopol solution solution**  
**AgroChem Inc**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----  
**Ultra-Soft Barrier**

**ULTRA-SOFT BARRIER**

**ULTRA-SOFT BARRIER**

**BRONOPOL TEAT DIP**

**WARNING**

**CAUSES SERIOUS EYE IRRITATION**

**FOR FARM AND INDUSTRIAL USE ONLY.**

**KEEP OUT OF REACH OF CHILDREN.**

**KEEP OUT OF DIRECT SUNLIGHT.**

**AVOID FREEZING - PRODUCT WILL BECOME THICKER WITH COLD.**

**PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.**

**FIRST AID / RESPONSE:**

**IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.**

**STORAGE: Store locked up.**

**DISPOSAL: Dispose of contents/ container to an approved waste disposal plant**

**EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494**

**INGREDIENTS:**

**Active Ingredients:**

**Bonopol – 0.25%**

**Emollients – 14% (Lanolin & Glycerine)**

**Use Ultra-Soft Barrier Dip effectively to reduce the spread of mastitis causing organisms. The specially selected emollients in Ultra-Soft Barrier Dip soften teat skin and protect the surface of the teat by keeping it moistened and smooth. Ultra-Soft Barrier Dip is suitable for use as a pre or post dip. Directions:**

**General Use**

**Ready to use. Do not dilute. Do not mix product with water, any other teat dips, or other product.**

**Use a teat dip cup to apply. Clean dip cup just prior to use. If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip. If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears**

the proper container labeling for this product. If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container. Teat Sprays can also be foamed or sprayed using recommended equipment.

### Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis. Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels. Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds. Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

### Post Dipping Use

Immediately after each milking, use product. Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

### Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.

Dipping should be resumed at least twice daily one week before calving.

# ULTRA-SOFT BARRIER

BRONOPOL  
TEAT DIP

## WARNING

AVOID FREEZING.  
HARMFUL IF SWALLOWED.  
KEEP OUT OF DIRECT SUNLIGHT.  
CAUSES SERIOUS EYE IRRITATION.  
KEEP OUT OF REACH OF CHILDREN.  
FOR FARM AND INDUSTRIAL USE ONLY.

PREVENTION: KEEP FROM FREEZING.  
If frozen, let thaw and shake thoroughly before using. Do NOT get in eyes or on clothing. Do NOT take internally. Do NOT allow to contaminate food. Wash thoroughly after handling.

FIRST AID / RESPONSE:  
IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.  
INGESTION: Drink large amounts of milk, egg whites, or water. Avoid alcohol. Contact a physician. Do not give anything by mouth to an unconscious person. Note to Physician: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

STORAGE: Keep container tightly closed when not in use. Store in a cool, dry, well-ventilated area out of direct sunlight.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

INGREDIENTS:  
Active Ingredients:  
Bronopol – 0.25%  
Emollients – 14% (Lanolin & Glycerine)

NET CONTENTS:

LOT#:

EXP:



Use Ultra-Soft Barrier Dip effectively to reduce the spread of mastitis causing organisms. The specially selected emollients in Ultra-Soft Barrier Dip soften teat skin and protect the surface of the teat by keeping it moistened and smooth. Ultra-Soft Barrier Dip is suitable for use as a pre or post dip.

#### Directions:

**General Use:**  
Ready to use. Do not dilute. Do not mix product with water, any other teat dips, or other product. Use a teat dip cup to apply. Clean dip cup just prior to use. If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip. If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product. If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container. Teat Sprays can also be foamed or sprayed using recommended equipment.

#### Pre Dipping Use:

Perform forestripping and inspect teats for clinical mastitis. Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels. Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds. Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

#### Post Dipping Use:

Immediately after each milking, use product. Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

#### Extended Use:

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.



**AgroChem**

26 Freedom Way • Saaboga Springs, NY 12866 • (518)226-4850

# ULTRA-SOFT BARRIER

bronopol solution solution

## Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:17307-3530
<b>Route of Administration</b>	TOPICAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BRONOPOL (UNII: 6PU1E16C9W) (BRONOPOL - UNII:6PU1E16C9W)	BRONOPOL	0.0025 kg in 1 kg

## Inactive Ingredients

Ingredient Name	Strength
C10-16 PARETH-1 (UNII: 3Y76363WPB)	.00445 kg in 1 kg
C9-11 PARETH-3 (UNII: 8O4974OMUO)	.00445 kg in 1 kg

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17307-3530-1	19 kg in 1 CONTAINER		
2	NDC:17307-3530-2	57 kg in 1 DRUM		
3	NDC:17307-3530-3	208 kg in 1 DRUM		
4	NDC:17307-3530-4	1040 kg in 1 CONTAINER		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		03/11/2020	

**Labeler** - AgroChem Inc (805427700)

**Registrant** - AgroChem Inc (805427700)

## Establishment

Name	Address	ID/FEI	Business Operations
AgroChem Inc		805427700	api manufacture, manufacture

Revised: 3/2020

AgroChem Inc