

# EPINEPHRINE PROFESSIONAL- epinephrine and isopropyl alcohol

## Focus Health Group

### Epinephrine Professional Convenience Kit

For treating anaphylaxis:

**Adults and Children 30 kg (66 lbs) or more:** 0.3 to 0.5 mg (0.3 to 0.5 ml) intramuscularly or subcutaneously into anterolateral aspect of the thigh every 5 to 10 minutes as necessary.

**Children 30 kg (66 lbs) or less:** 0.01 mg/kg (0.01 ml/kg), up to 0.3 mg (0.3 ml), intramuscularly or subcutaneously into anterolateral aspect of the thigh every 5 to 10 minutes as necessary.

### Principal Display Panel

**EPINEPHRINE PROFESSIONAL**  
EPINEPHRINE CONVENIENCE KIT with SAFETY SEAL

**For use by licensed health care professionals authorized to administer epinephrine intramuscularly or subcutaneously.**

**EPINEPHRINE 1mg/mL (1:1000)**  
See back of box for proper dosage and administration.

**Contents:**

- (1) Epinephrine Injection, 1 mg/mL (1:1000) VIAL
- (3) Syringes (25G x 1")
- (4) Alcohol Prep Pads

**Storage:** Store at 20 to 25 °C (68 to 77 °F). Epinephrine is light sensitive; protect from light and freezing.

**Contraindications:** Hypertension, Diabetes, Coronary artery disease, Myocardial infarction, Atherosclerosis, Hyperthyroidism, Glaucoma, Asthma, Pregnancy, Lactation, and use of MAO inhibitors.

**Warnings:** Do not use if the solution is cloudy or if it contains particulate matter. Do not inject into the buttock, digit, hand, or foot. Do not repeat repeated injections at the same site, as the resulting vasoconstriction may cause tissue necrosis.

**Directions:** Follow your provider's protocol or plan for dosage and administration. For treating anaphylaxis, intramuscularly into anterolateral aspect of the thigh every 5 to 15 minutes as necessary. For treating asthma, intramuscularly into anterolateral aspect of the thigh every 5 to 15 minutes as necessary. For treating hypertension, intramuscularly into anterolateral aspect of the thigh every 5 to 15 minutes as necessary.

**Caution:** Epinephrine may increase heart rate and blood pressure. Use with caution in patients with underlying heart disease. Administer with caution when used intravenously or subcutaneously. Epinephrine may increase heart rate and blood pressure. Use with caution in patients with underlying heart disease. Administer with caution when used intravenously or subcutaneously.

**Adverse Reactions:** Flushing, dizziness, headache, palpitations, tachycardia, tremor, and anxiety. May cause hypotension and/or hypoxemia. May cause hyperkalemia and/or hypocalcemia. May cause hypotension and/or hypoxemia. May cause hyperkalemia and/or hypocalcemia.

**Other information:** This product is not for use in children under 2 years of age. For more information, visit [www.focushealthgroup.com](http://www.focushealthgroup.com).

**Focus Health Group**  
NDC 24357-011-13 RX ONLY

# EPINEPHRINE PROFESSIONAL

epinephrine and isopropyl alcohol kit

## Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:24357-011
---------------------	-------------------------	---------------------------	---------------

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:24357-011-13	1 in 1 CARTON; Type 0: Not a Combination Product	07/01/2019	

## Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 VIAL	1 mL
Part 2	4 PACKET	4 mL

## Part 1 of 2

### ADRENALIN

epinephrine injection

## Product Information

<b>Item Code (Source)</b>	NDC:42023-159
<b>Route of Administration</b>	INTRAMUSCULAR, SUBCUTANEOUS, INTRAVENOUS

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
EPINEPHRINE (UNII: YKH834O4BH) (EPINEPHRINE - UNII:YKH834O4BH)	EPINEPHRINE	1 mg in 1 mL

## Inactive Ingredients

Ingredient Name	Strength
TARTARIC ACID (UNII: W4888I119H)	2.25 mg in 1 mL
HYDROCHLORIC ACID (UNII: QTT17582CB)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	1 mg in 1 mL
WATER (UNII: 059QF0KO0R)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	0.2 mg in 1 mL
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	0.457 mg in 1 mL
SODIUM CHLORIDE (UNII: 451W47IQ8X)	7.3 mg in 1 mL

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42023-159-25	1 mL in 1 VIAL; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA204200	07/01/2013	

## Part 2 of 2

### MCKESSON ALCOHOL PREP PAD

isopropyl alcohol swab

## Product Information

Item Code (Source)	NDC:68599-5804
Route of Administration	TOPICAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ISOPROPYL ALCOHOL (UNII: ND2M416302) (ISOPROPYL ALCOHOL - UNII:ND2M416302)	ISOPROPYL ALCOHOL	0.7 mL in 1 mL

## Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68599-5804-1	1 mL in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC			

OTC monograph not final	part333A	04/09/2010	
<b>Marketing Information</b>			
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
NDA	NDA204200	07/01/2019	

**Labeler** - Focus Health Group (826939949)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Par Sterile Products LLC		808402890	manufacture(42023-159)

<b>Establishment</b>			
<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
Focus Health Group		826939949	manufacture(24357-011)

Revised: 10/2022

Focus Health Group