

HONGO KILLER- tolnaftate powder
Efficient Laboratories Inc

Drug Facts

| Active Ingredients: | Purpose |
|-------------------------------|-------------------|
| Tolnaftate 1% W/W..... | Antifungal |

Uses

For effective treatment of:

- athlete's foot (tinea pedis)
- jock itch (Tinea Cruris)
- ringworm (tinea corporis)

For the relief of itching, burning feet.

For the prevention of athlete's foot with daily use

Purpose

Antifungal

Warnings

For external use only. Avoid contact with eyes.

Do not use on children under 2 years of age

Stop use and ask a doctor if irritation occurs or if there's no improvement within 4 weeks (for athlete's foot or ringworm) or within 2 weeks for jock itch.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. If pregnant or breast-feeding consult a health professional before use.

Directions

- clean the affected area and dry thoroughly
- Sprinkle Hongo Killer powder liberally over the affected area twice daily (morning and night) or as direct by doctor
- For athlete's foot: pay special attention to spaces between the toes. Wear well-fitting, ventilated shoes and change shoes and socks at least once daily
- For athlete's foot and ringworm, use daily for 4 weeks; for jock itch, use daily for 2 weeks. If condition persists longer, consult a doctor. This product is not effective on scalp or nails.
- Supervise children in the use of Hongo Killer

Questions or comments?

www.efficientlabs.com

Please Call 305-805-3456 (M-F 9:00-5:00 pm EST)

Inactive ingredients Corn (Zea Mays) Starch, Methylparaben, Propylparaben, Sodium


Bicarbonate, Talc and Zinc Oxide.


| Drug Facts | |
|---|----------------|
| Active Ingredients (%) | Purpose |
| Tolnaftate 1% w/w | Antifungal |
| Uses For effective treatment of: • athlete's foot (Tinea Pedis) • jock itch (Tinea Cruris) • ringworm (Tinea Corporis). For the relief of itching, burning feet • For the prevention of athlete's foot with daily use. | |
| Warnings For external use only. Avoid contact with eyes. | |
| Do not use • on children under 2 years of age. | |
| Stop use and ask a doctor if • irritation occurs or if there is no improvement within 4 weeks (for athlete's foot or ringworm) or within 2 weeks for (jock itch). | |
| Keep Out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. If pregnant or breast-feeding consult a health professional before use. | |
| Directions • Clean the affected area and dry thoroughly • Sprinkle Hongo Killer® powder liberally over the affected area twice daily (morning and night) or as directed by a doctor. For athlete's foot: Pay special attention to spaces between the toes, wear well-fitting ventilated shoes and change shoes and socks at least once daily. • For athlete's foot and ringworm, use daily for 4 weeks; for jock itch, use daily for 2 weeks. If condition persists longer, consult a doctor. This product is not effective on the scalp or nails. • Supervise children in the use of Hongo Killer®. | |
| Other Information • Store between 15° - 30°C (59° - 86° F) • Hongo Killer® is also available in spray, antifungal solution, cream, nail formula and spray powder. | |
| Inactive ingredients • Corn (Zea Mays) Starch, Methylparaben, Propylparaben, Sodium Bicarbonate, Talc, and Zinc Oxide. | |
| Questions or Comments? www.efficientlabs.com - 305-805-3456 M-F 9am-5pm EST | |

Distributed by Efficient Laboratories, Inc.
Miami, FL 33166
Manufactured in Dominican Republic

REV: 11/15







Hongo Killer®

Tolnaftate ANTIFUNGAL


Powder



**Cures
Athlete's
Foot-Jock
Itch &
Ringworm**

**Alivio
Efectivo
del Picor
y Ardor**

NET WT 90 g (3.17 oz)



110 mm

Drug Facts

| Active ingredient (%) | Purpose |
|-----------------------|------------|
| Tolnaftate 1% w/w | Antifungal |

Uses For effective treatment of: • athlete's foot (Tinea Pedis) • jock itch (Tinea Cruris) • ringworm (Tinea Corporis). For the relief of itching, burning feet • For the prevention of athlete's foot with daily use.

Warnings For external use only. Avoid contact with eyes. Do not use • on children under 2 years of age, unless directed by a doctor.

Stop use and ask a doctor if • irritation occurs (for the prevention of athlete's foot) or if irritation occurs or if there is no improvement within 4 weeks (for athlete's foot or ringworm) or within 2 weeks for (jock itch)

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. If pregnant or breast-feeding, consult a health professional before use.

Directions • Clean the affected area and dry thoroughly. • Sprinkle a thin layer of Hongo Killer® powder over affected area twice daily (morning and night) or as directed by a doctor. • Supervise children in the use of Hongo Killer®. For athlete's foot: Pay special attention to spaces between the toes, wear well-fitting, ventilated shoes, and change shoes and socks at least once daily. • For athlete's foot and ringworm, use daily for 4 weeks; for jock itch, use daily for 2 weeks. If condition persists longer, consult a doctor. This product is not effective on the scalp or nails. • To prevent athlete's foot, wash the feet and dry thoroughly. Apply a thin layer of Hongo Killer® powder to the feet once or twice daily (morning and/or night).

Other information • Store between 15°- 30° C (59°- 86° F)

Inactive ingredients • Corn (Zea Mays) Starch, Fragrance, Methylparaben, Propylparaben, Sodium Bicarbonate, Talc, and Zinc Oxide.

Questions or Comments? www.efficientlabs.com - 305-805-3456 M-F 9am-5pm EST

• Hongo Killer® is also available in spray, antifungal solution, cream, nail formula and spray powder.

REV: 10/16 Distributed by:
EFFICIENT LABORATORIES
Miami, FL 33166
Made in the Dominican Republic

| HONGO KILLER | | | |
|--|--|---------------------------|-----------------|
| tolnaftate powder | | | |
| Product Information | | | |
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:58593-288 |
| Route of Administration | TOPICAL | | |
| Active Ingredient/Active Moiety | | | |
| | Ingredient Name | Basis of Strength | Strength |
| | TOLNAFTATE (UNII: 06KB629TKV) (TOLNAFTATE - UNII:06KB629TKV) | TOLNAFTATE | 1 g in 100 g |
| Inactive Ingredients | | | |
| | Ingredient Name | | Strength |

| | |
|--|--|
| STARCH, CORN (UNII: O8232NY3SJ) | |
| METHYLPARABEN (UNII: A2I8C7HI9T) | |
| PROPYLPARABEN (UNII: Z8IX2SC1OH) | |
| SODIUM BICARBONATE (UNII: 8MDF5V39QO) | |
| TALC (UNII: 7SEV7J4R1U) | |
| ZINC OXIDE (UNII: SOI2LOH54Z) | |

| Packaging | | | | |
|------------------|------------------|--|----------------------|--------------------|
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | NDC:58593-288-04 | 90 g in 1 BOTTLE; Type 0: Not a Combination Product | 01/01/2000 | 10/15/2015 |
| 2 | NDC:58593-288-07 | 200 g in 1 BOTTLE; Type 0: Not a Combination Product | 05/02/2016 | |

| Marketing Information | | | |
|------------------------------|--|----------------------|--------------------|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| OTC Monograph Drug | M005 | 01/01/2000 | |

Labeler - Efficient Laboratories Inc (969044932)

| Establishment | | | |
|--------------------------------|---------|-----------|------------------------|
| Name | Address | ID/FEI | Business Operations |
| Laboratorios Dr. Collado, S.A. | | 817468001 | manufacture(58593-288) |

Revised: 5/2024

Efficient Laboratories Inc