

**CLEARLAX- polyethylene glycol 3350 powder, for solution**  
**Meijer, Inc.**

-----  
**Meijer Distribution, Inc. Clearlax® Drug Facts**

**Active ingredient (in each dose)**

Polyethylene Glycol 3350, 17 g (cap filled to line)

**Purpose**

Osmotic Laxative

**Use**

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

**Warnings**

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use**

if you have kidney disease, except under the advice and supervision of a doctor

**Ask a doctor before use if you have**

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug

**When using this product**

you may have loose, watery, more frequent stools

**Stop use and ask a doctor if**

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

**If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
  - use once a day
  - fill to top of white section in cap which is marked to indicate the correct dose (17 g)
  - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
- do not combine with starch-based thickeners used for difficulty swallowing
- ensure that the powder is fully dissolved before drinking
- do not drink if there are any clumps
- do not use more than 7 days
- children 16 years of age or under: ask a doctor

### **Other information**

- store at 20°-25°C (68°-77°F)
- tamper -evident: do not use if printed foil seal under cap is missing, open or broken

### **Inactive ingredients**

natural and artificial orange flavor, sucralose

### **Questions or comments?**

**1-800-719-9260**

### **Package/Label Principal Display Panel**

meijer®

ClearLax®

COMPARE TO MIRALAX® ACTIVE INGREDIENT

Polyethylene Glycol 3350

Powder for Solution

Osmotic Laxative

Grit Free Powder

RELIEVES OCCASIONAL CONSTIPATION/IRREGULARITY SOFTENS STOOL

ORANGE FLAVOR

14 ONCE-DAILY DOSES

Sugar Free

Dye Free

NET WT 8.3 OZ (238 g)

meijer.

NDC 79481-0090-0

# ClearLax<sup>®</sup>

COMPARE TO  
MIRALAX<sup>®</sup>  
ACTIVE INGREDIENT\*

Polyethylene Glycol 3350  
Powder for Solution  
Osmotic Laxative

RELIEVES OCCASIONAL  
CONSTIPATION/IRREGULARITY  
SOFTENS STOOL

Grit Free  
Powder

ORANGE FLAVOR



Sugar Free  
Dye Free

14  
ONCE-DAILY  
DOSES

NET WT 8.3 OZ (238 g)

: 7U9E1 6E F1

Peel Here

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool.  
Dissolves in Any Beverage

### Drug Facts

**Active ingredient (in each dose)** Purpose  
Polyethylene Glycol 3350,  
17 g (cap filled to line).....Osmotic Laxative

### Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

### Warnings

**Allergy alert:** Do not use if you are allergic to polyethylene glycol

**Do not use** if you have kidney disease, except under the advice and supervision of a doctor

### Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

**Ask a doctor or pharmacist before use if you are taking a prescription drug**

**When using this product** you may have loose, watery, more frequent stools

### Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week



3

PID 5339681

: 7U9E1 6E B1

### Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### Directions

- do not take more than directed unless advised by your doctor
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
  - use once a day
  - fill to top of white section in cap which is marked to indicate the correct dose (17 g)
  - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
  - do not combine with starch-based thickeners used for difficulty swallowing
  - ensure that the powder is fully dissolved before drinking
  - do not drink if there are any clumps
  - do not use more than 7 days
- children 16 years of age or under: ask a doctor

### Other information

- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

**Inactive ingredients** natural and artificial orange flavor, sucralose

**Questions or comments?** 1-800-719-9260

\*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark MiraLAX<sup>®</sup>.

Pat. No. 11,793,876

DIST. BY MEIJER  
DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com

ADHESIVE AREA • NO COPY

## CLEARLAX

polyethylene glycol 3350 powder, for solution

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:79481-0090
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>POLYETHYLENE GLYCOL 3350</b> (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g in 17 g

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

## Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	ORANGE	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:79481-0090-3	510 g in 1 BOTTLE; Type 0: Not a Combination Product	01/13/2023	
2	NDC:79481-0090-0	238 g in 1 BOTTLE; Type 0: Not a Combination Product	01/30/2024	

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA090685	01/13/2023	

**Labeler** - Meijer, Inc. (006959555)