

MUCINEX INSTASOOTHE SORE THROAT PLUS COUGH RELIEF-
dextromethorphan hydrobromide and hexylresorcinol lozenge
RB Health (US) LLC

MUCINEX® INSTASOOTHE™
SORE THROAT PLUS COUGH RELIEF

Drug Facts

<i>Active ingredients (in each lozenge)</i>	<i>Purposes</i>
Dextromethorphan HBr 5.0 mg	Cough suppressant
Hexylresorcinol 2.0 mg	Oral pain reliever

Uses

temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold
- occasional irritation and pain due to a sore mouth or sore throat
- pain associated with canker sores

Warnings

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- in children under 6 years of age

When using this product do not exceed recommended dosage

Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor or dentist if

- sore mouth symptoms do not improve in 7 days
- irritation, pain, or redness persists or worsens
- swelling develops
- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent

headache. These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

dose as follows or as directed by a doctor or dentist

adults and children 12 years of age and older	take 2 lozenges (one immediately after the other) and allow each lozenge to dissolve slowly in the mouth; may be repeated every 4 hours as needed, not to exceed 12 lozenges in any 24-hour period
children 6 to under 12 years of age	take 1 lozenge and allow product to dissolve slowly in the mouth; may be repeated every 4 hours as needed, not to exceed 6 lozenges in any 24-hour period
children under 6 years of age	do not use

Other information

- store at 20-25°C (68-77°F)
- protect contents from humidity

Inactive ingredients

corn syrup, D&C yellow no. 10, ethyl lactate, FD&C blue no. 1, flavor, propylene glycol, purified water, sucrose.

Questions?

1-866-MUCINEX (1-866-682-4639)

Dist. by: RB Health (US)

**Parsippany,
NJ 07054-0224**

Made in Thailand

PRINCIPAL DISPLAY PANEL - 20 Drop Pouch Bag

NDC 72854-263-20

Mucinex®
INSTASOOTHE™

SORE THROAT
+ COUGH RELIEF

Dextromethorphan HBr 5.0 mg | Cough Suppressant
Hexylresorcinol 2.0 mg | Oral Pain Reliever

3 IN 1 FORMULA

- ✓ Up to 4 Hours Cough Relief
- ✓ Cooling Comfort
- ✓ Fast Acting

See back
panel for
actual size

Winter Mint Flavor

20 MEDICATED DROPS



SORE THROAT + COUGH RELIEF

Dextromethorphan HBr 5.0 mg | Cough Suppressant
Hexylresorcinol 2.0 mg | Oral Pain Reliever

3 IN 1 FORMULA

- ✓ Up to 4 Hours Cough Relief
- ✓ Cooling Comfort
- ✓ Fast Acting



See back
panel for
actual size

WINTER MINT
FLAVOR

20 MEDICATED DROPS

Choose from our range of Mucinex® INSTASOOTHE™ products to treat your symptoms.

POWERFUL RELIEF FOR SORE THROAT, COUGH OR BOTH

3-IN-1 FORMULAS

SOOTHING COMFORT | PAIN RELIEF | COUGH RELIEF

Drug Facts		Drug Facts (continued)							
Active ingredients (in each lozenge)		Purposes							
Dextromethorphan HBr 5.0 mg	Cough suppressant	If pregnant or breast-feeding, ask a health professional before use.							
Hexylresorcinol 2.0 mg	Oral pain reliever	Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.							
Uses temporarily relieves		Directions ■ dose as follows or as directed by a doctor or dentist							
<ul style="list-style-type: none"> cough due to minor throat and bronchial irritation as may occur with the common cold occasional irritation and pain due to a sore mouth or sore throat pain associated with canker sores 		<table border="1"> <tr> <td>adults and children 12 years of age and older</td> <td>take 2 lozenges (one immediately after the other) and allow each lozenge to dissolve slowly in the mouth; may be repeated every 4 hours as needed, not to exceed 12 lozenges in any 24-hour period</td> </tr> <tr> <td>children 6 to under 12 years of age</td> <td>take 1 lozenge and allow product to dissolve slowly in the mouth; may be repeated every 4 hours as needed, not to exceed 6 lozenges in any 24-hour period</td> </tr> <tr> <td>children under 6 years of age</td> <td>do not use</td> </tr> </table>		adults and children 12 years of age and older	take 2 lozenges (one immediately after the other) and allow each lozenge to dissolve slowly in the mouth; may be repeated every 4 hours as needed, not to exceed 12 lozenges in any 24-hour period	children 6 to under 12 years of age	take 1 lozenge and allow product to dissolve slowly in the mouth; may be repeated every 4 hours as needed, not to exceed 6 lozenges in any 24-hour period	children under 6 years of age	do not use
adults and children 12 years of age and older	take 2 lozenges (one immediately after the other) and allow each lozenge to dissolve slowly in the mouth; may be repeated every 4 hours as needed, not to exceed 12 lozenges in any 24-hour period								
children 6 to under 12 years of age	take 1 lozenge and allow product to dissolve slowly in the mouth; may be repeated every 4 hours as needed, not to exceed 6 lozenges in any 24-hour period								
children under 6 years of age	do not use								
Warnings		Other information							
<p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use</p> <ul style="list-style-type: none"> if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. in children under 6 years of age <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> persistent or chronic cough such as occurs with smoking, asthma, or emphysema cough that occurs with too much phlegm (mucus) 		<ul style="list-style-type: none"> store at 20-25°C (68-77°F) protect contents from humidity keep lozenge in outer pack until use. Reseal outer pack after use. 							
When using this product do not exceed recommended dosage		Inactive ingredients corn syrup, D&C yellow no. 10, ethyl lactate, FD&C blue no. 1, flavor, propylene glycol, purified water, sucrose							
<p>Stop use and ask a doctor or dentist if</p> <ul style="list-style-type: none"> sore mouth symptoms do not improve in 7 days irritation, pain, or redness persists or worsens swelling develops cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition. 		Questions? 1-866-MUCINEX (1-866-682-4639)							



PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

ACTUAL SIZE





3 63824 21572 3



50044453

Coding Area - Copy free

Dist. by: RB Health (US)
Parsippany, NJ
07054-0224
Made in Thailand
©2023 RB Health
Patents:
www.reckitt.com/patents
www.mucine x.com
091323

Keep outer pack for full information.

MUCINEX INSTASOOTHE SORE THROAT PLUS COUGH RELIEF

dextromethorphan hydrobromide and hexylresorcinol lozenge

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72854-263
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg
HEXYLRESORCINOL (UNII: R9QTB5E82N) (HEXYLRESORCINOL - UNII:R9QTB5E82N)	HEXYLRESORCINOL	2 mg

Inactive Ingredients

Ingredient Name	Strength
CORN SYRUP (UNII: 9G5L16BK6N)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
ETHYL LACTATE (UNII: F3P750VW8I)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SUCROSE (UNII: C151H8M554)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	turquoise	Score	no score
Shape	ROUND	Size	18mm
Flavor	MINT	Imprint Code	S
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72854-263-20	20 in 1 BAG	07/01/2021	
1		1 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:72854-263-40	40 in 1 BAG	07/01/2021	
2		1 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	07/01/2021	

Labeler - RB Health (US) LLC (081049410)