ADAPALENE- adapalene gel AMERISOURCE BERGEN

Adapalene

Drug Facts

Active ingredient

Adapalene USP, 0.1% (retinoid) ¹

1 read consumer information leaflet

Purpose

Acne treatment

Use

For the treatment of acne

Warnings

For external use only

Do not use

- on damaged skin (cuts, abrasions, eczema, sunburn)
- if you are allergic to adapalene or any of the ingredients in this product.

If pregnant or breast-feeding, ask a doctor before use.

When using this product

- limit sun exposure, including light from tanning beds, and use sunscreen when going outdoors
- do not wax to remove hair in areas where the product has been applied
- during the early weeks of use, your acne may appear to worsen before it improves (this is normal); continue using as directed, unless you get irritation that becomes severe
- irritation (redness, itching, dryness, burning) is more likely to occur:
 - in the first few weeks of use
 - if using more than one topical acne medication at a time
 - but irritation usually lessens with continued use of this product
- it may take up to 3 months of once daily use to see results
- avoid product contact with eyes, lips, and mouth. If contact occurs, immediately flush the area with water.
- wash hands after use

Stop use and ask doctor if

- you become pregnant, or are planning to become pregnant, while using the product
- you have symptoms of an allergic reaction (such as itching, rash, hives, swelling of the lips, evelids, and shortness of breath)
- irritation becomes severe
- you see no improvement after 3 months of once daily use

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 12 years of age and older:

- use **once**daily
- clean the skin gently and pat dry before applying the product
- cover the entire affected area with a thin layer. For example, if your acne is on the face, apply the product to the entire face.
- do not use more than one time a day. Applying more than directed will not provide faster or better results, but may worsen skin irritation.

Children under 12 years of age:ask a doctor

Other information

- TAMPER EVIDENT: DO NOT USE IF THE SEAL ON THE TUBE IS PUNCTURED OR NOT VISIBLE.
- store at 20° to 25°C (68° to 77°F)[see USP Controlled Room Temperature].
- protect from freezing

Inactive ingredients

carbomer homopolymer type C, edetate disodium, phenoxyethanol, poloxamer 182, propylene glycol, purified water, sodium hydroxide.

Questions?

1-866-923-4914

Distributed By AmerisourceBergen 1 West First Ave., Conshohocken PA 19428

PRINCIPAL DISPLAY PANEL - 15 g Tube Carton

NDC 46122-768-36

Compare to the Differin [®]Gel [†] active ingredient

GOOD NEIGHBOR PHARMACY ®

PARABEN FREE

Adapalene Gel USP, 0.1% Acne Treatment

- Previously available only by prescription
- FDA approved
- Dermatologist developed
- Once Daily Topical Retinoid*
- * Read consumer information leaflet before use

NET WT 0.5 oz (15 g)



391.61 B294.0 1324

AND EXP DATE PRINT **# TOJ ROJ 9AJ 3 SIHT** NO COPY / NO COLOR **DA ON \HSINFAV ON**

0224 2248188

OIL FREE

FRAGRANCE FREE PARABEN FREE

DERMATOLOGIST DEVELOPED AND TESTED

First FDA-approved over-the-counter topical retinoid* for acne

*Read carton and enclosed consumer information leaflet before using this product. Keep this carton and consumer information leaflet. They contain important information.

[†]All trademarks are property of their respective owners. This product is not affiliated with the makers/owners of Differin® Gel.



Distributed By AmerisourceBerger at Ave., Conshohocki PA 19428 1 West First A Questions or Concerns?

www.mygnp.com Made in Canada

Rev: 02/2024



NDC 46122-768-36

Compare to the Differin®Gel† active ingredient



PARABEN FREE!

Adapalene Gel USP, 0.1%

Acne Treatment

- Previously available only by prescription
- FDA approved
- Dermatologist developed
- Once Daily Topical Retinoid*
- * Read consumer information leaflet before use

NET WT 0.5 oz (15 g)

Drug Facts

Active ingredient Purpose

Adapalene USP, 0.1% *read consumer information leaflet

For the treatment of acne

Warnings For external use only

Do not use

- on damaged skin (cuts, abrasions,
- eczema, sunburn) if you are allergic to adapalene or any of the ingredients in

If pregnant or breast-feeding, ask a doctor before use.

When using this product

- limit sun exposure, including light from tanning beds, and use sunscreen when going outdoors
- do not wax to remove hair in areas where the product has been applied during the early weeks of use, your
- acne may appear to worsen before it improves (this is normal); continue using as directed, unless you get irritation that becomes
- irritation (redness, itching, dryne burning) is more likely to occur.
- in the first few weeks of use
- if using more than one topical
- acne medication at a time but irritation usually lessen with continued use of
- it may take up to 3 months of once daily use to see results

Drug Facts (continued)

- avoid product contact with eyes, lips, and mouth. If contact occurs, immediately flush the area with water wash hands after use

Stop use and ask doctor if

- you become pregnant, or are planning to become pregnant, while using the product
- you have symptoms of an allergic reaction (such as itching, rash, hives, swelling of the lips, eyelids, and shortness of breath)
- irritation becomes severe
- you see no improvement after 3 months of once

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center

Directions

Adults and children 12 years of age and older:

- use once daily clean the skin gently and pat dry before applying
- the product cover the entire affected area with a thin layer. For
- example, if your acne is on the face, apply the product to the entire face. do not use more than one time a day. Applying more than directed will not provide faster or better results,
- but may worsen skin irritation. Children under 12 years of age: ask a doctor

Other information

- TAMPER EVIDENT: DO NOT USE IF THE SEAL ON THE TUBE IS PUNCTURED OR NOT VISIBLE. store at 20° to 25°C (68° to 77°F) [see USP
- Controlled Room Temperature).
- protect from freezing

Inactive ingredients

carbomer homopolymer type C, edetate disodium, phenoxyethanol, poloxamer 182, propylene glycol, purified water, sodium hydroxide

Questions? 1-866-923-4914



ADAPALENE

adapalene gel

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:46122-768

K COATING FREE AREA **Route of Administration**

TOPICAL

Active Ingredient/Active Moiety

, , , , , , , , , , , , , , , , , , ,		
Ingredient Name	Basis of Strength	Strength
ADAPALENE (UNII: 1L4806J2QF) (ADAPALENE - UNII:1L4806J2QF)	ADAPALENE	1 mg in 1 g

Inactive Ingredients		
Ingredient Name	Strength	
CARBOMER HOMOPOLYMER TYPE C (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: 4Q93RCW27E)		
EDETATE DISODIUM (UNII: 7FLD91C86K)		
PHENOXYETHANOL (UNII: HIE492ZZ3T)		
POLOXAMER 182 (UNII: JX0HIX6OAG)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SODIUM HYDROXIDE (UNII: 55X04QC32I)		

F	Packaging				
#	t Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:46122-768- 36	1 in 1 CARTON	02/26/2024		
1		15 g in 1 TUBE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215940	02/26/2024	

Labeler - AMERISOURCE BERGEN (007914906)

Registrant - Taro Pharmaceuticals U.S.A., Inc. (145186370)

Establishment				
Name	Address	ID/FEI	Business Operations	
Taro Pharmaceuticals Inc.		206263295	manufacture(46122-768)	

Revised: 3/2024 AMERISOURCE BERGEN