

ORIS CHX- chlorhexidine gluconate rinse DENTSPLY Professional


Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

ORIS Chlorhexidine Gluconate Oral Rinse, 0.12%

Rx Only

DESCRIPTION

ORIS Chlorhexidine Gluconate Oral Rinse, 0.12% is an oral rinse containing 0.12% chlorhexidine gluconate (1, 1'-hexamethylene bis[5-(p-chlorophenyl) biguanide] di-D-gluconate) in a base containing 11.6% v/v alcohol, FD&C Blue No. 1, glycerin, natural and artificial peppermint flavor, PEG-40 sorbitan diisostearate, saccharin sodium, and purified water. Chlorhexidine Gluconate Oral Rinse, 0.12% is a near-neutral solution (pH range 5-7). Chlorhexidine gluconate is a salt of chlorhexidine and gluconic acid. Its structure is:



CLINICAL PHARMACOLOGY

Chlorhexidine Gluconate Oral Rinse, 0.12% provides antimicrobial activity during oral rinsing. The clinical significance of 0.12% chlorhexidine gluconate oral rinse's anti-microbial activities is not clear. Microbiological sampling of plaque has shown a general reduction of counts of certain assayed bacteria, both aerobic and anaerobic, ranging from 54-97% through six months' use.

Use of chlorhexidine gluconate oral rinse in a six month clinical study did not result in any significant changes in bacterial resistance, overgrowth of potentially opportunistic organisms or other adverse changes in the oral microbial ecosystem. Three months after chlorhexidine gluconate use was discontinued, the number of bacteria in plaque had returned to baseline levels and resistance of plaque bacteria to chlorhexidine gluconate was equal to that at baseline.

Pharmacokinetics

Pharmacokinetic studies with 0.12% chlorhexidine gluconate oral rinse indicate approximately 30% of the active ingredient, chlorhexidine gluconate, is retained in the oral cavity following rinsing. This retained drug is slowly released into the oral fluids. Studies conducted on human subjects and animals demonstrate chlorhexidine gluconate is poorly absorbed from the gastrointestinal tract. The mean plasma level of chlorhexidine gluconate reached a peak of 0.206 µg/g in humans 30 minutes after they ingested a 300 mg dose of the drug. Detectable levels of chlorhexidine gluconate were not present in the plasma of these subjects 12 hours after the compound was administered. Excretion of chlorhexidine gluconate occurred primarily through the feces (~90%). Less than 1% of the chlorhexidine gluconate ingested by these subjects was excreted in the urine.

INDICATIONS

Chlorhexidine Gluconate Oral Rinse, 0.12% is indicated for use between dental visits as part of a professional program for the treatment of gingivitis as characterized by redness and swelling of the gingivae, including gingival bleeding upon probing. Chlorhexidine Gluconate Oral Rinse, 0.12% has not been tested among patients with acute necrotizing ulcerative gingivitis (ANUG). For patients having coexisting gingivitis and periodontitis, see **PRECAUTIONS**.

CONTRAINDICATIONS

Chlorhexidine Gluconate Oral Rinse, 0.12% should not be used by persons who are known to be hypersensitive to chlorhexidine gluconate or other formula ingredients.

WARNINGS

The effect of Chlorhexidine Gluconate Oral Rinse, 0.12% on periodontitis has not been determined. An increase in supragingival calculus was noted in clinical testing in chlorhexidine gluconate oral rinse users compared with control users. It is not known if chlorhexidine gluconate use results in an increase in subgingival calculus. Calculus deposits should be removed by a dental prophylaxis at intervals not greater than six months. Anaphylaxis, as well as serious allergic reactions, have been reported during postmarketing use with dental products containing chlorhexidine. (See **CONTRAINDICATIONS**).

PRECAUTIONS

General

1. For patients having coexisting gingivitis and periodontitis, the presence or absence of gingival inflammation following treatment with Chlorhexidine Gluconate Oral Rinse, 0.12% should not be used as a major indicator of underlying periodontitis.
2. Chlorhexidine Gluconate Oral Rinse, 0.12% can cause staining of oral surfaces, such as tooth surfaces, restorations, and the dorsum of the tongue. Not all patients will experience a visually significant increase in tooth staining. In clinical testing, 56% of chlorhexidine gluconate oral rinse users exhibited a measurable increase in facial anterior stain, compared to 35% of control users after six months; 15% of chlorhexidine gluconate users developed what was judged to be heavy stain, compared to 1% of control users after six months. Stain will be more pronounced in patients who have heavier accumulations of unremoved plaque. Stain resulting from the use of Chlorhexidine Gluconate Oral Rinse, 0.12% does not adversely affect health of the gingivae or other oral tissues. Stain can be removed from most tooth surfaces by conventional professional prophylactic techniques. Additional time may be required to complete the prophylaxis. Discretion should be used when prescribing to patients with anterior facial restorations with rough surfaces or margins. If natural stain cannot be removed from these surfaces by a dental prophylaxis, patients should be excluded from Chlorhexidine Gluconate Oral Rinse, 0.12% treatment if permanent discoloration is unacceptable. Stain in these areas may be difficult to remove by dental prophylaxis and on rare occasions may necessitate replacement of these restorations.
3. Some patients may experience an alteration in taste perception while undergoing treatment with chlorhexidine gluconate oral rinse. Rare instances of permanent taste alteration following chlorhexidine gluconate oral rinse have been reported via postmarketing surveillance.

Pregnancy

Teratogenic Effects

Pregnancy Category B

Reproduction studies have been performed in rats and rabbits at chlorhexidine gluconate doses up to 300 mg/kg/day and 40 mg/kg/day, respectively, and have not revealed evidence of harm to the fetus.

However, adequate and well-controlled studies in pregnant women have not been done. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when Chlorhexidine Gluconate Oral Rinse, 0.12% is administered to a nursing women. In parturition and lactation studies with rats, no evidence of impaired parturition or of toxic effects to suckling pups was observed when chlorhexidine gluconate was administered to dams at doses that were over 100 times greater than that which would result from a person's ingesting 30 mL (2 capfuls) of chlorhexidine gluconate oral rinse, 0.12% per day.

Pediatric Use

Clinical effectiveness and safety of Chlorhexidine Gluconate Oral Rinse, 0.12% have not been established in children under the age of 18.

Carcinogenesis, Mutagenesis, and Impairment of Fertility

In a drinking water study in rats, carcinogenic effects were not observed at doses up to 38 mg/kg/day. Mutagenic effects were not observed in two mammalian *in vivo* mutagenesis studies with chlorhexidine gluconate. The highest doses of chlorhexidine used in a mouse dominant-lethal assay and a hamster cytogenetics test were 1000 mg/kg/day and 250 mg/kg/day, respectively. No evidence of impaired fertility was observed in rats at doses up to 100 mg/kg/day.

ADVERSE REACTIONS

The most common side effects associated with chlorhexidine gluconate oral rinses are 1) an increase in staining of teeth and other oral surfaces; 2) an increase in calculus formation; and 3) an alteration in taste perception, see **WARNINGS** and **PRECAUTIONS**. Oral irritation and local allergy-type symptoms have been spontaneously reported as side effects associated with use of chlorhexidine gluconate rinse. The following oral mucosal side effects were reported during placebo-controlled adult clinical trials: aphthous ulcer, grossly obvious gingivitis, trauma, ulceration, erythema, desquamation, coated tongue, keratinization, geographic tongue, mucocele, and short frenum. Each occurred at a frequency of less than 1.0%. Among post-marketing reports, the most frequently reported oral mucosal symptoms associated with chlorhexidine gluconate oral rinse, 0.12% are stomatitis, gingivitis, glossitis, ulcer, dry mouth, hypesthesia, glossal edema, and paresthesia. Minor irritation and superficial desquamation of the oral mucosa have been noted in patients using chlorhexidine gluconate oral rinse, 0.12%. There have been cases of parotid gland swelling and inflammation of the salivary glands (sialadenitis) reported in patients using chlorhexidine gluconate oral rinse.

To report SUSPECTED ADVERSE REACTIONS, contact DENTSPLY Professional, at 1-800-989-8825 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

OVERDOSAGE

Ingestion of 1 to 2 ounces of Chlorhexidine Gluconate Oral Rinse, 0.12% by a small child (~10 kg body weight) might result in gastric distress, including nausea, or signs of alcohol intoxication. Medical attention should be sought if more than 4 ounces of Chlorhexidine Gluconate Oral Rinse, 0.12% is ingested by a small child or if signs of alcohol intoxication develop.

DOSAGE AND ADMINISTRATION

Chlorhexidine Gluconate Oral Rinse, 0.12% therapy should be initiated directly following a dental prophylaxis. Patients using Chlorhexidine Gluconate Oral Rinse, 0.12% should be reevaluated and

given a thorough prophylaxis at intervals no longer than six months. Recommended use is twice daily oral rinsing for 30 seconds, morning and evening after toothbrushing. Usual dosage is 1/2 fl oz (marked in cup) of undiluted Chlorhexidine Gluconate Oral Rinse, 0.12%. Patients should be instructed to not rinse with water or other mouthwashes, brush teeth, or eat immediately after using Chlorhexidine Gluconate Oral Rinse, 0.12%. Chlorhexidine Gluconate Oral Rinse, 0.12% is not intended for ingestion and should be expectorated after rinsing.

HOW SUPPLIED

Chlorhexidine Gluconate Oral Rinse, 0.12% is a blue, peppermint flavored liquid in:

A 16 fl oz (473 mL) amber plastic bottle with a child-resistant closure and dosage cup for consumer use.

It should be dispensed in original container or in amber glass.

Store above freezing 0°C (32°F)

Rx Only

Keep out of reach of children

Manufactured for
DENTSPLY Professional
DENTSPLY International
1301 Smile Way
York, PA 17404 USA

Measuring cup is shrink wrapped to container for your protection. Remove shrink wrap and dosage cup. Press cap down while turning counterclockwise to open container. To reseal, turn cap clockwise until tightly locked.

DIRECTIONS FOR USE: Fill measuring cup to the fill line (1/2 ounce). Swish in mouth undiluted for 30 seconds twice daily, after breakfast and before bedtime, expel remainder. Or, use as prescribed.

NOTE: To minimize medicinal taste, do not rinse with water immediately after use.

INGREDIENTS: 0.12% chlorhexidine gluconate in a base containing 11.6% alcohol, FD&C Blue No. 1, glycerin, natural and artificial peppermint flavor, PEG-40 sorbitan diisostearate, saccharin sodium and purified water.

Rx Only

What to expect when using Oris[®] Chlorhexidine Gluconate Oral Rinse, 0.12%

Your dentist has prescribed Oris Chlorhexidine Gluconate Oral Rinse, 0.12% to treat your gingivitis, to help reduce the redness and swelling of your gums, and also to help you control any gum bleeding. Use Oris Chlorhexidine Gluconate Oral Rinse, 0.12% regularly, as directed by your dentist, in addition to daily brushing. Spit out after use. Oris Chlorhexidine Gluconate Oral Rinse should not be swallowed.

If you develop allergic symptoms such as skin rash, itch, generalized swelling, breathing difficulties, light headedness, rapid heart rate, upset stomach or diarrhea, seek medical attention immediately. Oris Chlorhexidine Gluconate Oral Rinse, 0.12% should not be used by persons who have a sensitivity to it or its components.

Oris Chlorhexidine Gluconate Oral Rinse, 0.12% may cause some tooth discoloration, or increases in tartar (calculus) formation, particularly in areas where stain and tartar usually form. It is important to see your dentist for removal of any stain or tartar at least every six months, or more frequently if your dentist advises.

- Both stain and tartar can be removed by your dentist or hygienist. Oris Chlorhexidine Gluconate Oral Rinse, 0.12% may cause permanent discoloration of some front- tooth fillings.

- To minimize discoloration, you should brush and floss daily, emphasizing areas which begin to discolor.
- Oris Chlorhexidine Gluconate Oral Rinse, 0.12% may taste bitter to some patients and can affect how foods and beverages taste. This will become less noticeable in most cases with continued use of Oris Chlorhexidine Gluconate Oral Rinse, 0.12%.
- To avoid taste interference, rinse with Oris Chlorhexidine Gluconate Oral Rinse, 0.12% *after* meals. Do not rinse with water or other mouthwashes immediately after rinsing with Oris Chlorhexidine Gluconate Oral Rinse, 0.12%

If you have any questions or comments about Oris Chlorhexidine Gluconate Oral Rinse, 0.12%, contact your dentist or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to DENTSPLY Professional, at 1-800-989-8825 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Manufactured for:
DENTSPLY Professional
DENTSPLY International
1301 Smile Way
York, PA 17404 USA

P/N 130030-2 Rev-13 09/14
Rev. 720:05 03/14

PRINCIPAL DISPLAY PANEL - 473 mL Bottle Label

Lift Here
DENTSPLY
PROFESSIONAL

ORIS®

CHLORHEXIDINE
GLUCONATE
ORAL RINSE, 0.12%

PLACE PHARMACY
LABEL HERE.

Dispense in original container
or in amber glass.

PEPPERMINT
FLAVOR

Rx Only
STORE ABOVE FREEZING 0°C (32°F).
KEEP OUT OF REACH OF CHILDREN
16 fl oz (473 mL)

Reorder No. 130030

ADVERSE REACTIONS: The most common side effects associated with chlorhexidine gluconate oral rinses are 1) an increase in staining of teeth and other oral surfaces; 2) an increase in calculus formation; and 3) an alteration in taste perception, see **WARNINGS** and **PRECAUTIONS**. Oral irritation and local allergy-type symptoms have been spontaneously reported as side effects associated with use of chlorhexidine gluconate rinse. The following oral mucosal side effects were reported during placebo-controlled adult clinical trials: aphthous ulcer, grossly obvious gingivitis, trauma, ulceration, erythema, desquamation, coated tongue, keratinization, geographic tongue, mucocele, and short frenum. Each occurred at a frequency of less than 1.0%. Among post-marketing reports, the most frequently reported oral mucosal symptoms associated with chlorhexidine gluconate oral rinse, 0.12% are stomatitis, gingivitis, glossitis, ulcer, dry mouth, hypesthesia, glossal edema, and paresthesia. Minor irritation and superficial desquamation of the oral mucosa have been noted in patients using chlorhexidine gluconate oral rinse, 0.12%. There have been cases of parotid gland swelling and inflammation of the salivary glands (sialadenitis) reported in patients using chlorhexidine gluconate oral rinse.

To report SUSPECTED ADVERSE REACTIONS, contact DENTSPLY Professional, at 1-800-989-8825 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

OVERDOSAGE: Ingestion of 1 to 2 ounces of Chlorhexidine Gluconate Oral Rinse, 0.12% by a small child (~10 kg body weight) might result in gastric distress, including nausea, or signs of alcohol intoxication. Medical attention should be sought if more than 4 ounces of Chlorhexidine Gluconate Oral Rinse, 0.12% is ingested by a small child or if signs of alcohol intoxication develop.

DOSAGE AND ADMINISTRATION: Chlorhexidine Gluconate Oral Rinse, 0.12% therapy should be initiated directly following a dental prophylaxis. Patients using Chlorhexidine Gluconate Oral Rinse, 0.12% should be reevaluated and given a thorough prophylaxis at intervals no longer than six months. Recommended use is twice daily oral rinsing for 30 seconds, morning and evening after toothbrushing. Usual dosage is 1/2 fl oz (marked in cup) of undiluted Chlorhexidine Gluconate Oral Rinse, 0.12%. Patients should be instructed to not rinse with water or other mouthwashes, brush teeth, or eat immediately after using Chlorhexidine Gluconate Oral Rinse, 0.12%. Chlorhexidine Gluconate Oral Rinse, 0.12% is not intended for ingestion and should be expectorated after rinsing.

HOW SUPPLIED: Chlorhexidine Gluconate Oral Rinse, 0.12% is a blue, peppermint flavored liquid in: A 16 fl oz (473 mL) amber plastic bottle with a child-resistant closure and dosage cup for consumer use. It should be dispensed in original container or in amber glass.

Store above freezing 0°C (32°F)

Rx Only
Keep out of reach of children

Manufactured for
DENTSPLY Professional
DENTSPLY International
1301 Smile Way
York, PA 17404 USA

ORIS CHX

chlorhexidine gluconate rinse

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:65222-720
Route of Administration	DENTAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Chlorhexidine Gluconate (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	Chlorhexidine Gluconate	1.2 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
Water (UNII: 059QF0KO0R)	
Alcohol (UNII: 3K9958V90M)	
Glycerin (UNII: PDC6A3C0OX)	
PEG-40 Sorbitan Diisostearate (UNII: JL4CCU7IIG)	
Saccharin Sodium (UNII: SB8ZUX40TY)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

Product Characteristics

Color	BLUE	Score	
Shape		Size	
Flavor	PEPPERMINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65222-720-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
UNAPPROVED DRUG OTHER		01/01/1999	

Labeler - DENT SPLY Professional (144140845)

Revised: 5/2015

DENT SPLY Professional