

IBUPROFEN- ibuprofen tablet, film coated
CVS Pharmacy

CVS 44-291

Active ingredient (in each brown tablet)

Ibuprofen USP, 200 mg (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - the common cold
 - toothache
 - muscular aches
 - backache
 - minor pain of arthritis
 - menstrual cramps
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- rash
- skin reddening
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take more or for a longer time than directed
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- take a blood thinning (anticoagulant) or steroid drug
- have 3 or more alcoholic drinks every day while using this product

Do not use

- right before or after heart surgery
- if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over; take 1 tablet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 tablet, 2 tablets may be used
 - do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years; ask a doctor

Other information

- store between 20°-25°C (68°-77°F)

- avoid excessive heat 40°C (104°F)
- use by expiration date on package

Inactive ingredients

carnauba wax, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments?

Call 1-800-426-9391 8:30AM-4:00PM ET, Monday-Friday

Principal Display Panel

CVS Health™

Compare to the active ingredient in Advil® Tablets†

Coated Tablets

NDC 59779-291-03

TAMPER EVIDENT: USE ONLY IF THIS BLISTER IS INTACT

Ibuprofen

IBUPROFEN TABLETS USP, 200 MG

Pain reliever,
Fever reducer (**NSAID**)

**10 COATED
TABLETS**

Actual Size

†This product is not manufactured or distributed by Wyeth Consumer Healthcare, owner of the registered trademark Advil® Tablets.

50844 ORG061629103

Distributed by: CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895

© 2016 CVS/pharmacy CVS.com®

1-800-SHOP CVS V-11112

√ **CVS Quality**

Money Back Guarantee



Compare to the active ingredient in Advil® Tablets†

Coated Tablets

Ibuprofen

IBUPROFEN TABLETS USP, 200 mg

Pain reliever, Fever reducer (NSAID)

10 COATED TABLETS

Actual Size

NO COAT AREA

RETAIN VIAL CARD FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredient (in each brown tablet) Purpose
 Ibuprofen USP, 200 mg (NSAID)* Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - menstrual cramps
 - backache
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- rash
- skin reddening
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- right before or after heart surgery
- if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk

Drug Facts (continued)

if stomach upset occurs or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 tablet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 tablet, 2 tablets may be used
- do not exceed 6 tablets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)
- avoid excessive heat 40°C (104°F)
- use by expiration date on package

Inactive ingredients carnauba wax, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

This product is not manufactured or distributed by Wyeth Consumer Healthcare, owner of the registered trademark Advil® Tablets. ©2015 CVS Pharmacy, Inc. Distributed by: CVS Pharmacy, Inc. One CVS Drive, Woonsocket, RI 02899. ©2015 CVS Pharmacy. CVS.com 1-800-THOP CVS. V-11132

CVS 44-291

IBUPROFEN

ibuprofen tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:59779-291
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	BROWN	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;291
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59779-291-98	2 in 1 PACKAGE	05/24/1988	
1		10 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:59779-291-15	1 in 1 CARTON	05/24/1988	
2		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC:59779-291-29	1 in 1 CARTON	05/24/1988	
3		150 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
4	NDC:59779-291-12	1 in 1 CARTON	05/24/1988	
4		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
5	NDC:59779-291-03	10 in 1 VIAL; Type 0: Not a Combination Product	05/24/1988	
6	NDC:59779-291-05	400 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	05/24/1988	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075010	05/24/1988	

Labeler - CVS Pharmacy (062312574)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(59779-291)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(59779-291)