

## **PREMIER VALUE FIBER LAXATIVE- calcium polycarbophil tablet, film coated** **Chain Drug Consortium**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **Premier Value Fiber Caplets**

#### **Active ingredient (in each caplet )**

Calcium polycarbophil 625 mg equivalent to 500 mg polycarbophil

#### **Purpose**

Bulk-forming laxative

#### **Uses**

- relieves occasional constipation to help restore and maintain regularity
- this product generally produces bowel movement in 12 to 72 hours.

#### **Warnings**

Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

#### **Ask your doctor before use if you have**

- abdominal pain, nausea or vomiting
- a sudden change in bowel habits that persists over a period of 2 weeks

#### **Ask your doctor or pharmacist before use if**

you are taking any other drug. Take this product 2 or more hours before or after other drugs. All laxatives may affect how other drugs work.

#### **When using this product**

- do not use for more than 7 days unless directed by a doctor
- do not take more than 8 caplets in a 24 hour period unless directed by a doctor

#### **Stop use and ask a doctor if**

rectal bleeding occurs or if you fail to have a bowel movement after use of this or any other laxative. These could be signs of a serious condition.

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

## Directions

- take each dose of this product with at least 8 ounces (a full glass) of water or other fluid. Taking this product without enough liquid may cause choking. See choking warning.
- This product works naturally so continued use for one to three days is normally required to provide full benefit. Dosage may vary according to diet, exercise, previous laxative use or severity of constipation

age	recommended dose	daily maximum
adults & children 12 years of age and over	2 caplets once a day	up to 4 times a day
children under 12 years	consult a physician	

## Other information

- **each caplet contains:** calcium 125 mg
- do not use if printed seal under cap is torn or missing
- store at 20-25°C (68-77°F)
- protect contents from moisture

## Inactive ingredients

calcium carbonate, caramel, crospovidone, hypromellose, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium lauryl sulfate

## Principal Display Panel

NDC 68016-259-00

Premier Value

Fiber Caplets

FIBER LAXATIVE

Calcium Polycarbophil 625 mg

- Relieves constipation
- Restores regularity
- Daily source of fiber

COMPARE TO FIBERCON® ACTIVE INGREDIENT  
90 CAPLETS

This product is not manufactured or distributed by Pfizer Consumer Healthcare, the distributor of FIBERCON®

Distributed by:

Chain Drug Consortium LLC.

3301 N.W. Boca Raton Blvd.

Suite 101, Boca Raton FL 33431

Fiber Laxative Label

NDC 68016-259-00

COMPARE TO FIBERCON®  
ACTIVE INGREDIENT\*



# Fiber Caplets

**FIBER LAXATIVE**  
Calcium Polycarbophil 625 mg

- Relieves constipation
- Restores regularity
- Daily source of fiber

90 CAPLETS



**Drug Facts**

**Active ingredient (in each caplet)** **Purpose**  
Calcium polycarbophil 625 mg  
equivalent to 500 mg polycarbophil.....Bulk-forming laxative

**Uses** ■ relieves occasional constipation to help restore and maintain regularity ■ this product generally produces bowel movement in 12 to 72 hours.

**Warnings**  
**Choking:** Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

**Ask your doctor before use if you have** ■ abdominal pain, nausea or vomiting ■ a sudden change in bowel habits that persists over a period of 2 weeks

**Ask your doctor or pharmacist before use if you are taking any other drug.** Take this product 2 or more hours before or after other drugs. All laxatives may affect how other drugs work.

**When using this product** ■ do not use for more than 7 days unless directed by a doctor ■ do not take more than 8 caplets in a 24 hour period unless directed by a doctor

**Stop use and ask a doctor if** rectal bleeding occurs or if you fail to have a bowel movement after use of this or any other laxative. These could be signs of a serious condition. ▶

\*This product is not manufactured or distributed by Pfizer Consumer Healthcare, the distributor of FIBERCON®

Distributed by:  
Chain Drug Consortium, LLC.  
3301 N.W. Boca Raton Blvd.,  
Suite 101, Boca Raton, FL 33431



8 40986 01538 0

12090PVL8

LOT: \_\_\_\_\_

EXP: \_\_\_\_\_

**PEEL BACK HERE**

READ ENTIRE LABEL INCLUDING DRUG FACTS

**Drug Facts (continued)**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

**Directions** ■ take each dose of this product with at least 8 ounces (a full glass) of water or other fluid. Taking this product without enough liquid may cause choking. See choking warning.  
■ this product works naturally so continued use for one to three days is normally required to provide full benefit. Dosage may vary according to diet, exercise, previous laxative use or severity of constipation

age	recommended dose	daily maximum
adults & children 12 years of age and over	2 caplets once a day	up to 4 times a day
children under 12 years	consult a physician	

**Other information** ■ each caplet contains: calcium 125 mg ■ do not use if printed seal under cap is torn or missing ■ store at 20-25°C (68-77°F) ■ protect contents from moisture

**Inactive ingredients** calcium carbonate, caramel, crospovidone, hypromellose, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium lauryl sulfate.

**STOP PEELING HERE**

# PREMIER VALUE FIBER LAXATIVE

calcium polycarbophil tablet, film coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68016-259
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>CALCIUM POLYCARBOPHIL</b> (UNII: 8F049NKY49) (POLYCARBOPHIL - UNII:W25LM17A4W)	CALCIUM POLYCARBOPHIL	625 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)	
<b>CARMEL</b> (UNII: T9D99G2B1R)	
<b>CROSPROVIDONE (15 MPA.S AT 5%)</b> (UNII: 68401960MK)	
<b>HYPROMELLOSE 2208 (100 MPA.S)</b> (UNII: B1QE5P712K)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MALTODEXTRIN</b> (UNII: 7CVR7L4A2D)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POLYETHYLENE GLYCOL 800</b> (UNII: UH6KR4953D)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	

## Product Characteristics

<b>Color</b>	BROWN (light)	<b>Score</b>	2 pieces
<b>Shape</b>	CAPSULE (caplet)	<b>Size</b>	19mm
<b>Flavor</b>		<b>Imprint Code</b>	RP120
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:68016-259-00	90 in 1 BOTTLE; Type 0: Not a Combination Product	10/28/2015	

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph not final	part334	10/28/2015	

