

POLYETHYLENE GLYCOL 3350- polyethylene glycol 3350 powder, for solution
Discount Drug Mart

Drug Mart 44-742

Active ingredient (in each dose)

Polyethylene Glycol 3350,
17 grams (cap filled to the indicated "17 GRAMS" line)

Purpose

Laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use

if you have kidney disease, except under the advice and supervision of a doctor.

Ask a doctor before use if you have

- nausea, vomiting, or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are

taking a prescription drug.

When using this product

you may have loose, watery, more frequent stools.

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you need to use a laxative for longer than 1 week
- you get diarrhea

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle cap is a measuring cup marked to contain 17 grams of powder when filled to the indicated "17 GRAMS" line
- adults and children 17 years of age and older:
 - fill to indicated "17 GRAMS" line in cap which is marked to indicate the correct dose 17 grams
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - use once a day
 - use no more than 7 days
- children 16 years of age or younger: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if foil seal under cap printed with "SEALED for YOUR PROTECTION" is missing, open or broken.

Inactive ingredients

none

Questions or comments?

1-800-426-9391

Principal display panel

Discount

Drug

Mart

Food Fair

NDC 53943-742-90

ORIGINAL PRESCRIPTION STRENGTH

Polyethylene

Glycol 3350

Powder For Solution **Osmotic Laxative**

Relieves occasional constipation (irregularity)
and softens stool

- Sugar Free
- Dissolves in any beverage

*Compare to the active ingredient in MiraLAX®

UNFLAVORED POWDER

NET WT 17.9 OZ (510 g) **30 ONCE-DAILY DOSES**

SATISFACTION GUARANTEED IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE

TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART 211 COMMERCE DRIVE, MEDINA, OHIO 44256

DISTRIBUTED BY: DRUG MART - FOOD FAIR
211 COMMERCE DRIVE
MEDINA, OH 44256
www.discount-drugmart.com

Polyethylene Glycol 3350 increases the frequency of bowel movements and softens the stool

TAMPER-EVIDENT: DO NOT USE IF PRINTED FOIL SEAL UNDER CAP IS MISSING, OPEN OR BROKEN

*This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark MiraLAX®.

50844 REV0719A74290

7116-0510-Q8-BL

Rev 06/21

 <p>NDC 53943-742-90</p> <p>ORIGINAL PRESCRIPTION STRENGTH</p> <h1>Polyethylene Glycol 3350</h1> <p>Powder For Solution Osmotic Laxative</p> <p>Relieves occasional constipation (irregularity) and softens stool</p> <p>◆ Sugar Free ◆ Dissolves in any beverage</p> <p>*Compare to the active ingredient in MiraLAX®</p> <p>UNFLAVORED POWDER</p> <p>NET WT 17.9 OZ (510g) 30 ONCE-DAILY DOSES</p>	<p>SATISFACTION GUARANTEED IF DISSATISFIED, RETURN UNUSED PORTION AND PACKAGE TO STORE WHERE PURCHASED. IF UNABLE TO RETURN TO STORE, SEND REASON FOR DISSATISFACTION, NAME, ADDRESS AND EMPTY PACKAGE TO: DISCOUNT DRUG MART 211 COMMERCE DRIVE, MEDINA, OHIO 44256</p> <p>DISTRIBUTED BY: DRUG MART - FOOD FAIR 211 COMMERCE DRIVE MEDINA, OH 44256 www.discount-drugmart.com</p>  <p>0 93351 02939 4</p> <p>Exp. Date: _____</p> <p>Lot No.: _____</p> <p>Imprint/No. Varnish Area</p> <p>7116-0510-Q8-BL Rev 06/21</p>	<p>Polyethylene Glycol 3350 increases the frequency of bowel movements and softens the stool</p> <p>TAMPER-EVIDENT: DO NOT USE IF PRINTED FOIL SEAL UNDER CAP IS MISSING, OPEN OR BROKEN</p> <table border="1"> <tr> <th>Drug Facts</th> <th>Purpose</th> </tr> <tr> <td> Active ingredient (in each dose) Polyethylene Glycol 3350, 17 grams (cap filled to the indicated "17 GRAMS" line) </td> <td>Osmotic Laxative</td> </tr> <tr> <td colspan="2"> Use ■ relieves occasional constipation (irregularity) ■ generally produces a bowel movement in 1 to 3 days </td> </tr> <tr> <td colspan="2"> Warnings Allergy alert: Do not use if you are allergic to polyethylene glycol Do not use if you have kidney disease, except under the advice and supervision of a doctor. Ask a doctor before use if you have ■ nausea, vomiting, or abdominal pain ■ a sudden change in bowel habits that lasts over 2 weeks ■ irritable bowel syndrome Ask a doctor or pharmacist before use if you are taking a prescription drug. When using this product you may have loose, watery, more frequent stools. Stop use and ask a doctor if ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. ■ you need to use a laxative for longer than 1 week ■ you get diarrhea </td> </tr> </table> <table border="1"> <tr> <th>Drug Facts (continued)</th> </tr> <tr> <td> Directions ■ do not take more than directed unless advised by your doctor ■ the bottle cap is a measuring cup marked to contain 17 grams of powder when filled to the indicated "17 GRAMS" line ■ adults and children 17 years of age and older: ■ fill to indicated "17 GRAMS" line in cap which is marked to indicate the correct dose 17 grams ■ stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink ■ ensure that the powder is fully dissolved before drinking ■ do not drink if there are any clumps ■ use once a day ■ use no more than 7 days ■ children 16 years of age or younger: ask a doctor </td> </tr> <tr> <td> Other information ■ store at 20°-25°C (68°-77°F) ■ tamper-evident: do not use if foil seal under cap printed with "SEALED FOR YOUR PROTECTION" is missing, open or broken. </td> </tr> <tr> <td> Inactive ingredients none </td> </tr> <tr> <td> Questions or comments? 1-800-426-8391 </td> </tr> </table>	Drug Facts	Purpose	Active ingredient (in each dose) Polyethylene Glycol 3350, 17 grams (cap filled to the indicated "17 GRAMS" line)	Osmotic Laxative	Use ■ relieves occasional constipation (irregularity) ■ generally produces a bowel movement in 1 to 3 days		Warnings Allergy alert: Do not use if you are allergic to polyethylene glycol Do not use if you have kidney disease, except under the advice and supervision of a doctor. Ask a doctor before use if you have ■ nausea, vomiting, or abdominal pain ■ a sudden change in bowel habits that lasts over 2 weeks ■ irritable bowel syndrome Ask a doctor or pharmacist before use if you are taking a prescription drug. When using this product you may have loose, watery, more frequent stools. Stop use and ask a doctor if ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. ■ you need to use a laxative for longer than 1 week ■ you get diarrhea		Drug Facts (continued)	Directions ■ do not take more than directed unless advised by your doctor ■ the bottle cap is a measuring cup marked to contain 17 grams of powder when filled to the indicated "17 GRAMS" line ■ adults and children 17 years of age and older: ■ fill to indicated "17 GRAMS" line in cap which is marked to indicate the correct dose 17 grams ■ stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink ■ ensure that the powder is fully dissolved before drinking ■ do not drink if there are any clumps ■ use once a day ■ use no more than 7 days ■ children 16 years of age or younger: ask a doctor	Other information ■ store at 20°-25°C (68°-77°F) ■ tamper-evident: do not use if foil seal under cap printed with "SEALED FOR YOUR PROTECTION" is missing, open or broken.	Inactive ingredients none	Questions or comments? 1-800-426-8391
Drug Facts	Purpose														
Active ingredient (in each dose) Polyethylene Glycol 3350, 17 grams (cap filled to the indicated "17 GRAMS" line)	Osmotic Laxative														
Use ■ relieves occasional constipation (irregularity) ■ generally produces a bowel movement in 1 to 3 days															
Warnings Allergy alert: Do not use if you are allergic to polyethylene glycol Do not use if you have kidney disease, except under the advice and supervision of a doctor. Ask a doctor before use if you have ■ nausea, vomiting, or abdominal pain ■ a sudden change in bowel habits that lasts over 2 weeks ■ irritable bowel syndrome Ask a doctor or pharmacist before use if you are taking a prescription drug. When using this product you may have loose, watery, more frequent stools. Stop use and ask a doctor if ■ you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition. ■ you need to use a laxative for longer than 1 week ■ you get diarrhea															
Drug Facts (continued)															
Directions ■ do not take more than directed unless advised by your doctor ■ the bottle cap is a measuring cup marked to contain 17 grams of powder when filled to the indicated "17 GRAMS" line ■ adults and children 17 years of age and older: ■ fill to indicated "17 GRAMS" line in cap which is marked to indicate the correct dose 17 grams ■ stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink ■ ensure that the powder is fully dissolved before drinking ■ do not drink if there are any clumps ■ use once a day ■ use no more than 7 days ■ children 16 years of age or younger: ask a doctor															
Other information ■ store at 20°-25°C (68°-77°F) ■ tamper-evident: do not use if foil seal under cap printed with "SEALED FOR YOUR PROTECTION" is missing, open or broken.															
Inactive ingredients none															
Questions or comments? 1-800-426-8391															

Drug Mart 44-742

POLYETHYLENE GLYCOL 3350

polyethylene glycol 3350 powder, for solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:53943-742
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g in 17 g

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53943-742-90	510 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/21/2019	
2	NDC:53943-742-49	238 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/21/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090812	08/21/2019	

Labeler - Discount Drug Mart (047741335)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	pack(53943-742)

Revised: 12/2023

Discount Drug Mart