

**IBUPROFEN CA- ibuprofen tablet, coated**  
**TARGET CORPORATION**

-----  
**TARGET IBUPROFEN 200 MG CAPLETS (Brown)**

***Active ingredient (in each caplet)***

Ibuprofen 200 mg (NSAID)\*

\* nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever / fever reducer

***Uses***

temporarily relieves minor aches and pains due to:

- headache
- toothache
- backache
- menstrual cramps
- the common cold
- muscular aches
- minor pain of arthritis

temporarily reduces fever

***Warnings***

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

hives

- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

**Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

**Ask a doctor or pharmacist before use if you are**

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

**When using this product**

- take with food or milk if stomach upset occurs

**Stop use and ask a doctor if**

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better

you have symptoms of heart problems or stroke:

- chest pain
- trouble breathing
- weakness in one part or side of body
- slurred speech
- leg swelling

pain gets worse or lasts more than 10 days

fever gets worse or lasts more than 3 days

redness or swelling is present in the painful area

any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

### Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

### Directions

- **do not take more than directed**
- **the smallest effective dose should be used**

adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist

if pain or fever does not respond to 1 caplet, 2 caplets may be used  
do not exceed 6 caplets in 24 hours, unless directed by a doctor

children under 12 years: ask a doctor

**Inactive ingredients** colloidal silicon dioxide, croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

**Questions or comments?** call **1-800-910-6874**

**Ibuprofen**  
200 mg Ibuprofen Caplets  
Pain Reliever / Fever Reducer  
(NSAID)

**up&up**  
200 CAPLETS\*\*\* (\*\*CAPSULE-SHAPED TABLETS)

NDC 11673-990-20

**TAMPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOIL SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING**

**RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION**

**Uses** ■ temporarily relieves minor aches and pains due to: headache, toothache, backache, menstrual cramps, the common cold, muscular aches, minor pain of arthritis ■ temporarily reduces fever

**Warnings**  
**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: hives, facial swelling, asthma (wheezing), shock, skin reddening, rash, blisters if an allergic reaction occurs, stop use and seek medical help right away.  
**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:  
■ are age 60 or older ■ have had stomach ulcers or bleeding problems  
■ take a blood-thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
■ have 3 or more alcoholic drinks every day while using this product  
■ take more or for a longer time than directed

**Hear attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.  
**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions** ■ **do not take more than directed**  
■ **the smallest effective dose should be used**  
■ adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist  
■ if pain or fever does not respond to 1 caplet, 2 caplets may be used  
■ do not exceed 6 caplets in 24 hours, unless directed by a doctor  
■ children under 12 years: ask a doctor

**Other information** ■ read all warnings and directions before use  
■ store at 20-25°C (68-77°F)  
**Questions or comments?** Call **1-800-910-6874**

094010228 R00  
C-002262-01-034-0000  
Distributed by Target Corporation  
Minneapolis, MN 55403  
**Made in India**  
TM & © 2024 Target Brands, Inc.  
R00R\_1023  
4015003 Code No.: 60/0RUGS/515  
Lot No.:  
Exp. Date:

**Varnish Omit Area**

### Drug Facts

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION

**Active ingredient (in each caplet)** Purpose  
Ibuprofen 200 mg (NSAID)\* .....Pain reliever/Fever reducer  
\*nonsteroidal anti-inflammatory drug

#### Uses

- temporarily relieves minor aches and pains due to:
  - headache
  - toothache
  - backache
  - menstrual cramps
  - the common cold
  - muscular aches
  - minor pain of arthritis
- temporarily reduces fever

#### Warnings

**Allergy alert:** Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

### Drug Facts (continued)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Heart attack and stroke warning:** NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

#### Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

### Drug Facts (continued)

#### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic

#### Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

#### When using this product

- take with food or milk if stomach upset occurs

#### Stop use and ask a doctor if

- you experience any of the following signs of stomach

### Drug Facts (continued)

#### Bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
  - chest pain
  - trouble breathing
  - weakness in one part or side of body
  - slurred speech
  - leg swelling
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Drug Facts (continued)

(1-800-222-1222).

#### Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 caplet, 2 caplets may be used
- do not exceed 6 caplets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

#### Other information

- read all warnings and directions before use
- store at 20-25°C (68-77°F)

**Inactive ingredients** colloidal silicon dioxide,

### Drug Facts (continued)

croscarmellose sodium, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, pregelatinized starch, talc, titanium dioxide

**Questions or comments?** Call 1-800-910-6874

600R 1023

**Satisfaction guaranteed – Love it or your money back.**

Distributed by Target Corporation  
Minneapolis, MN 55403  
Made in India

TM & ©2024 Target Brands, Inc.

\*This product is not manufactured or distributed by Haleon group of companies, owner of the registered trademark Advil® Caplets.

**ibuprofen**  
200 mg Ibuprofen Caplets (NSAID)  
Pain Reliever / Fever Reducer



Code No.: 600R055/615

TAMPER EVIDENCE: DO NOT USE THIS PRODUCT IF THE IMPRINTED SEAL OVER THE MOUTH OF THE BOTTLE IS CUT, TORN, BROKEN OR MISSING



Lot No.:  
Exp. Date:

Varnish/Print  
Omit Area

Compare to active ingredient in Advil® Ibuprofen Caplets\*

NDC 11673-990-20

# Ibuprofen



**200 mg Ibuprofen Caplets**  
Pain Reliever / Fever Reducer (NSAID)

**200 CAPLETS\*\***  
(\*\*CAPSULE-SHAPED TABLETS)



Actual Size  
200 Caplets



Omit Area



Omit Area

Varnish/Print Omit Area



## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11673-990
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>IBUPROFEN</b> (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	brown	<b>Score</b>	no score
<b>Shape</b>	CAPSULE ((CAPLET))	<b>Size</b>	15mm
<b>Flavor</b>		<b>Imprint Code</b>	117
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:11673-990-42	24 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	
2	NDC:11673-990-20	200 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
ANDA	ANDA091239	01/01/2020	

**Labeler** - TARGET CORPORATION (006961700)

**Registrant** - TIME CAP LABORATORIES, INC (037005209)

**Establishment**

<b>Name</b>	<b>Address</b>	<b>ID/FEI</b>	<b>Business Operations</b>
MARKSANS PHARMA LTD		925822975	manufacture(11673-990)

Revised: 1/2024

TARGET CORPORATION