

ANTI DIARRHEAL ANTI GAS- loperamide hydrochloride, simethicone tablet
Rite Aid Corporation

Rite Aid Corporation Anti-Diarrheal Anti-Gas Drug Facts

Active ingredients (in each caplet)

Loperamide HCl 2 mg

Simethicone 125 mg

Purposes

Anti-diarrheal

Anti-gas

Uses

relieves symptoms of diarrhea plus bloating, pressure and cramps, commonly referred to as gas

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

- if you have bloody or black stool
- if you have difficulty swallowing

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.
- you have difficulty swallowing the caplet

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- take only on an empty stomach (1 hour before or 2 hours after a meal)
- take with a full (8 oz.) glass of water
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; ½ caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- **each caplet contains:** calcium 165 mg, sodium 3 mg
- store between 20°-25°C (68°-77°F). Protect from light.
- **do not use if printed blister unit is broken or torn**

Inactive ingredients

acesulfame potassium, croscarmellose sodium, dibasic calcium phosphate, maltodextrin, microcrystalline cellulose, natural and artificial flavor, propylene glycol, stearic acid

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to the active ingredients of Imodium[®] Multi-Symptom Relief

ANTI-DIARRHEAL/ANTI-GAS

LOPERAMIDE HYDROCHLORIDE & SIMETHICONE TABLETS, 2 mg/125 mg

ANTI-DIARRHEAL/ANTI-GAS

Multi-symptom relief caplet

Relieves symptoms of diarrhea plus:

cramps & pressure

bloating/gas

ACTUAL SIZE

24 CAPLETS*

*CAPSULE-SHAPED TABLETS



Drug Facts (continued)
Inactive ingredients: asarilium epotilium, crocarnellon sodium, dibasic calcium phosphate, malic acid, microcrystalline cellulose, methylcellulose, polyethylene glycol, stearic acid.
Questions or comments? 1-800-771-9928

Drug Facts
Warnings: Do not use if you have bloody or black stool, if you have difficulty swallowing, or if you have a history of abnormal heart rhythm. Ask a doctor before use if you have liver or kidney disease, or if you are taking a prescription drug. Loperamide may interact with certain prescription drugs. With use, dizziness, drowsiness, or weakness may occur. Be careful when driving or operating machinery. Stop use and ask a doctor if diarrhea lasts for more than 2 days, symptoms get worse, or you have difficulty swallowing or a serious condition. You may have difficulty swallowing if you take this medicine with a prescription drug.
Directions: If you are taking this medicine for the first time, ask your doctor for the right dose. If you are taking it again, ask your doctor for the right dose. Do not take more than 4 capsules in 24 hours. If you are taking it for the first time, ask your doctor for the right dose. If you are taking it again, ask your doctor for the right dose. Do not take more than 4 capsules in 24 hours. If you are taking it for the first time, ask your doctor for the right dose. If you are taking it again, ask your doctor for the right dose. Do not take more than 4 capsules in 24 hours. If you are taking it for the first time, ask your doctor for the right dose. If you are taking it again, ask your doctor for the right dose. Do not take more than 4 capsules in 24 hours.
Other information: Each caplet contains calcium 65 mg, sodium 3 mg. Do not use if you have difficulty swallowing or a serious condition. You may have difficulty swallowing if you take this medicine with a prescription drug.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

CODE AREA
 OR162 83 C1

DISTRIBUTED BY:
 RITE AID, 30 HUNTER LANE,
 CMP HILL, PA 17011
 www.riteaid.com

SATISFACTION GUARANTEE
 If you're not satisfied, we'll happily refund your money.

* This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium® Multi-Symptom Relief.

ANTI-DIARRHEAL / ANTI-GAS
 LOPERAMIDE HYDROCHLORIDE & SIMETHICONE TABLETS, 2 mg/125 mg
 ANTI-DIARRHEAL/ANTI-GAS

NDC 11822-1003-3

Compare to the active ingredients of Imodium® Multi-Symptom Relief*

ANTI-DIARRHEAL / ANTI-GAS
 LOPERAMIDE HYDROCHLORIDE & SIMETHICONE TABLETS, 2 mg/125 mg
 ANTI-DIARRHEAL/ANTI-GAS

Multi-symptom relief caplet
 Relieves symptoms of diarrhea *plus*:
 cramps & pressure bloating/gas

24 CAPLETS**
 **CAPSULE-SHAPED TABLETS

RITE AID PHARMACY



ANTI DIARRHEAL ANTI GAS

loperamide hydrochloride, simethicone tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-1003
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	125 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	OVAL	Size	17mm
Flavor		Imprint Code	L0R1;2;125
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-1003-2	12 in 1 CARTON	01/03/2019	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11822-1003-1	1 in 1 CARTON	01/03/2019	05/31/2020

2		42 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:11822-1003-3	24 in 1 CARTON	02/24/2020	
3		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA209837	01/03/2019	

Labeler - Rite Aid Corporation (014578892)

Revised: 12/2023

Rite Aid Corporation