

ACETAMINOPHEN- acetaminophen tablet, coated
TARGET CORPORATION

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

341R-TGT APAP 500 MG 11673-838

Active Ingredient (in each tablet)

acetaminophen 500 mg

hypromellose, mineral oil, povidone, pregelatinized starch, sodium starch glycolate* stearic acid, titanium dioxide

*may contain this ingredient

PAIN RELIEVER - FEVER REDUCER

Uses: temporarily relieves minor aches and pains due to: the common cold, headache, backache, minor pain of arthritis, toothache, muscular aches, premenstrual and menstrual cramps, temporarily reduces fever.

Directions: Do not take more than directed (see overdose warning)

adults and children 12 years and over: take 2 caplets every 4 to 6 hours while symptoms last

do not take more than 8 caplets in 24 hours

do not take for more than 10 days unless directed by a doctor

children under 12 years: do not use this adult extra strength product in children under 12 years of age; this will provide more than the recommended dose (overdose) of acetaminophen and may cause liver damage.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

NDC 11679-838-01

**extra strength
acetaminophen
caplets, 500 mg**

pain reliever/fever reducer
for adults



100 CAPLETS** 500 mg each
(**CAPSULE-SHAPED TABLETS)

KEEP THIS CARTON FOR COMPLETE PRODUCT INFORMATION.

Use

Use only to relieve minor aches and pains (e.g.,

headaches, toothaches, muscle aches, and

toothaches). Do not use to relieve

fever. Do not use to relieve

joint pain, back pain, or

other types of pain.

Do not use to relieve

any type of pain that

is worse than the

pain you are trying

to relieve.

Do not use to relieve

any type of pain

that is worse than

the pain you are

trying to relieve.

Do not use to relieve

any type of pain

that is worse than

the pain you are

trying to relieve.

Do not use to relieve

any type of pain

that is worse than

the pain you are

trying to relieve.

Do not use to relieve

any type of pain

Coating Omit Area

NDC 11674-839-24

**extra strength
acetaminophen
caplets, 500 mg**

pain reliever/fever reducer

for adults



225 CAPLETS**
500 mg each
CAPSULE-SHAPED TABLETS

PLEASE READ THE FOLLOWING INFORMATION CAREFULLY BEFORE YOU TAKE THIS MEDICINE. IT MAY BE IMPORTANT FOR YOU TO KNOW OTHER THINGS THAT MAY AFFECT THE WAY THIS MEDICINE WORKS FOR YOU. THIS INFORMATION DOES NOT TAKE THE PLACE OF INFORMATION YOU RECEIVE FROM YOUR HEALTH CARE PROVIDER AND DOES NOT TAKE THE PLACE OF INFORMATION YOU RECEIVE FROM YOUR PHARMACEUTICAL PROVIDER.

Drug Facts

Active Ingredient (in each caplet)

acetaminophen

Warnings

Do not take if you have ever had liver disease or if you have a history of liver problems. Do not take if you are taking other medicines that may increase your risk of liver damage.

Directions

Take 1 caplet by mouth every 6 hours as needed for pain or fever. Do not take more than 6 caplets in 24 hours. If you are taking this medicine for pain, do not take it for more than 3 days. If you are taking it for fever, do not take it for more than 3 days. Do not take it for more than 10 days if you are taking it for pain or fever.

Other information

This medicine contains acetaminophen. Do not take other medicines that contain acetaminophen while you are taking this medicine. Do not take this medicine if you are taking any of the following medicines: aspirin, ibuprofen, naproxen, or other nonsteroidal anti-inflammatory drugs (NSAIDs).

Contains

225 caplets

Keep out of reach of children.

See the inside of this container for important information about this medicine.

Other information

Use by expiration date on package

Other information

225 caplets

Keep out of reach of children.

See the inside of this container for important information about this medicine.

Other information

Use by expiration date on package

Other information

225 caplets

Keep out of reach of children.

See the inside of this container for important information about this medicine.

Other information

Use by expiration date on package

Other information

225 caplets

Keep out of reach of children.

See the inside of this container for important information about this medicine.

Other information


Use by expiration date on package

NDC 11673-838-50

Compare to active ingredient in Extra Strength Tylenol® Caplets†

extra strength acetaminophen caplets, 500 mg

pain reliever/fever reducer
for adults



500 CAPLETS

500 CAPLETS** - 500 mg each
(**CAPSULE-SHAPED TABLETS)

ACTUAL SIZE

DAZPER EVIDENT: DO NOT USE THIS PRODUCT IF THE IMPRINTED FOLIO SEAL OVER THE MOUTH OF THE BOTTLE IS GUT, TORN, BROKEN OR MISSING

<p>Drug Facts</p> <p>Active ingredient (in each caplet) Purpose Acetaminophen 500 mg Pain reliever/fever reducer</p> <p>Uses</p> <ul style="list-style-type: none"> temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> headaches the common cold sore throat muscle aches menstrual cramps temporarily reduces fever <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: <ul style="list-style-type: none"> more than 8 caplets in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen or more alcoholic drinks every day while using this product. <p>Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> skin redness hives itch If a skin reaction occurs, stop use and seek medical help right away.</p> </p>	<p>Drug Facts (continued)</p> <p>Do not use <ul style="list-style-type: none"> with any other drug containing acetaminophen (paracetamol or propacetamol) If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist If you are allergic to acetaminophen or any of the inactive ingredients in this product for more than 10 days for pain unless directed by a doctor for more than 3 days for fever unless directed by a doctor <p>Ask a doctor before use if you have liver disease</p> <p>Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin</p> <p>Stop using and ask a doctor if <ul style="list-style-type: none"> symptoms do not improve new symptoms occur pain or fever persists or gets worse <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> </p></p>	<p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none"> do not take more than directed adults and children 12 years and over: <ul style="list-style-type: none"> take 2 caplets (1,000 mg) every 6 hours while symptoms last do not take more than 6 caplets (3,000 mg) in 24 hours, unless directed by a doctor do not use for more than 10 days unless directed by a doctor children under 12 years: ask a doctor <p>Other information</p> <ul style="list-style-type: none"> STORAGE: <ul style="list-style-type: none"> store at 20°-25° (68°-77°) excursions permitted between 15°-30° (59°-86°) use by expiration date on package Inactive ingredients: hypromellose, mannitol, polydextrose, pregelatinized starch, sodium starch glycolate, croscarmellose, titanium dioxide. *may contain this ingredient <p>Dist. by: Target Corp., Mpls., MN 55403 TM & © 2020 Target Brands, Inc.</p>
---	---	--

†This product is not interchangeable with other brands. Consumer features, cover of the original packaging may vary. © 2019

894.01.0218 R00 C-001227-01-4331

80% LUPC

FOR POSITION

194.8 34.150

Vanish Omit Area

ACETAMINOPHEN

acetaminophen tablet, coated

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:11673-838

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
MINERAL OIL (UNII: T5L8T28FGP)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	TCL341
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-838-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	
2	NDC:11673-838-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	
3	NDC:11673-838-50	500 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	
4	NDC:11673-838-26	225 in 1 BOTTLE; Type 0: Not a Combination Product	01/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	01/01/2020	

Labeler - TARGET CORPORATION (006961700)

Registrant - TIME CAP LABORATORIES, INC (037052099)

Establishment

Name	Address	ID/FEI	Business Operations
TIME CAP LABORATORIES, INC.		037052099	manufacture(11673-838)

