

0.5% IODINE TEAT DIP- iodine solution

Metz Sales

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).

0.5% Iodine Teat Dip

DIRECTIONS

Do not dilute this product. Use at full strength. Discard dip solution when it becomes visibly dirty or if sediment is present. If teats are chapped or sore, consult your veterinarian before beginning or continuing to use this product.

AS A PRE-DIP:

Remove all visible dirt by washing the teats and the base of the udder, using a minimal amount of an approved udder wash solution and a clean single service towel. Dry the teats and the udder completely with a clean single service paper towel. Dip or spray the teats. Allow 30 seconds contact time. Before attaching milker, remove the teat dip by **DRYING THOROUGHLY WITH A CLEAN SINGLE SERVICE PAPER TOWEL.**

AS A POST-DIP:

Immediately after milker is removed, dip or spray the teats with undiluted product. If the cow will be returned to a below freezing temperature environment allow the teat dip to dry before discharging her from the milking area. Immediately before the next milking wash and/or pre-dip the teats and udder.

ACTIVE INGREDIENTS

a-(p-Nonylphenyl) - omegahydroxypoly (oxyethylene) iodine complex.
(Provides 0.5% titratable iodine. Equivalent to 5,000 parts per million titratable iodine.)

CAUTION:

AVOID CONTACT WITH FOOD

KEEP OUT OF REACH OF CHILDREN!

FIRST AID

IF SPLASHED IN EYES: Rinse immediately with plenty of water, also under the eyelids, for at least 15 minutes. Get medical attention if irritation occurs.

IN CASE OF CONTACT: Wash off immediately with plenty of water for at least 15 minutes. Get medical attention if symptoms occur.

IF INHALED: Move to fresh air. If symptoms persist, call a physician.

IF SWALLOWED: Drink plenty of water. Do NOT induce vomiting. Never give anything by

**CONTAINS
5% GLYCERIN**

METZ
One Half Percent
IODINE TEAT TIP

**MINIMUM
pH 4.0**

An aid in controlling the spread of bacteria that may cause mastitis.

DIRECTIONS

Do not dilute this product. Use at full strength. Discard dip solution when it becomes visibly dirty or if sediment is present. If teats are chapped or sore, consult your veterinarian before beginning or continuing to use this product.

AS A PRE-DIP:

Remove all visible dirt by washing the teats and the base of the udder, using a minimal amount of an approved udder wash solution and a clean single service towel. Dry the teats and the udder completely with a clean single service paper towel. Dip or spray the teats. Allow 30 seconds contact time. Before attaching milker, remove the teat dip by **DRYING THOROUGHLY WITH A CLEAN SINGLE SERVICE PAPER TOWEL.**

AS A POST-DIP:

Immediately after milker is removed, dip or spray the teats with undiluted product. If the cow will be returned to a below freezing temperature environment allow the teat dip to dry before discharging her from the milking area. Immediately before the next milking wash and/or pre-dip the teats and udder.

ACTIVE INGREDIENTS

a-(p-Nonylphenyl) - omegahydroxypoly (oxyethylene) iodine complex. (Provides 0.5% titratable iodine. Equivalent to 5,000 parts per million titratable iodine.)

CAUTION:

AVOID CONTACT WITH FOOD

KEEP OUT OF REACH OF CHILDREN!

FIRST AID

IF SPLASHED IN EYES: Rinse immediately with plenty of water, also under the eyelids, for at least 15 minutes. Get medical attention if irritation occurs.

IN CASE OF CONTACT: Wash off immediately with plenty of water for at least 15 minutes. Get medical attention if symptoms occur.

IF INHALED: Move to fresh air. If symptoms persist, call a physician.

IF SWALLOWED: Drink plenty of water. Do NOT induce vomiting. Never give anything by mouth to an unconscious person. Consult a physician if necessary

Manufactured By:

METZ SALES, LLC.
Williamsburg, PA 16693

24 HOUR EMERGENCY SERVICE ASSISTANCE CALL:

CHEMTREC AT 1-800-424-9300

LOT NUMBER EXPIRES NET CONTENTS

0.5% IODINE TEAT DIP

iodine solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:50748-0018
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	0.002 g in 1 g

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	0.8901 g in 1 g
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	0.003 g in 1 g
NONOXYNOL-12 (UNII: 6NR43D77O6)	0.03 g in 1 g
GLYCERIN (UNII: PDC6A3C0OX)	0.0487 g in 1 g
SODIUM HYDROXIDE (UNII: 55X04QC32I)	0.0262 g in 1 g

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50748-0018-2	4 in 1 CASE		
1	NDC:50748-0018-1	1 g in 1 JUG		
2	NDC:50748-0018-3	5 g in 1 PAIL		
3	NDC:50748-0018-4	15 g in 1 DRUM		
4	NDC:50748-0018-5	30 g in 1 DRUM		
5	NDC:50748-0018-6	55 g in 1 DRUM		
6	NDC:50748-0018-7	250 g in 1 CONTAINER		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		06/14/2016	

Labeler - Metz Sales (054824487)

Establishment

Name	Address	ID/FEI	Business Operations
Metz Sales		054824487	manufacture, api manufacture

Revised: 6/2016

Metz Sales