

**ASSURED ASPIRIN- aspirin tablet, delayed release**  
**SPIRIT PHARMACEUTICALS LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**ASSURED™ ASPIRIN**

**Drug Facts**

**Active ingredient (in each tablet)**

Aspirin 81 mg (NSAID) <sup>1</sup>

---

1 nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Uses**

- for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches, fever or other symptoms needing immediate relief.**
- ask your doctor about other uses for Aspirin 81 mg

**Warnings**

**Reye's syndrome**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning**

This product contains an NSAID, which may cause stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use** if you are allergic to aspirin or any other pain reliever/fever reducer

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are** taking a prescription drug for:

- gout
- diabetes
- arthritis

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding**, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### **Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- **each tablet contains:** sodium 0.36 mg
- save carton for full directions and warnings
- store at room temperature

**Inactive ingredients**

acrycoat, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10, lactose, microcrystalline cellulose, polysorbate 80, sodium hydroxide, talc, titanium dioxide, triethyl citrate, yellow oxide of iron

**Questions or comments?**

**1-888-333-9792**

**DISTRIBUTED BY  
GREENBRIER INTERNATIONAL, INC.  
500 VOLVO PARKWAY, CHESAPEAKE, VA 23320**

**PRINCIPAL DISPLAY PANEL**

ASSURED™

COMPARE TO ACTIVE INGREDIENT OF  
BAYER® LOW DOSE ASPIRIN\*

LOW DOSE

Aspirin

Aspirin (NSAID) 81 mg

Pain Reliever

Actual Size



**Drug Facts (continued)**

Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)  
Have 3 or more alcoholic drinks every day while using this product  
Do not use if you are allergic to aspirin or any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for:** ■ gout ■ diabetes ■ arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or a loss of hearing occurs

**SAVE BOX FOR COMPLETE DRUG FACTS**

**Drug Facts**

**Active ingredient (in each tablet)**  
Aspirin 81 mg (NSAID)\*  
\*nonsteroidal anti-inflammatory drug

**Uses** ■ for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches, fever or other symptoms needing immediate relief.  
■ ask your doctor about other uses for Aspirin 81 mg

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:  
■ hives ■ facial swelling  
■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause stomach bleeding. The chance is higher if you:  
■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug

**ASSURED**

**LOW DOSE**

**Aspirin**

• Aspirin (NSAID) 81 mg  
Pain Reliever

COMPARE TO ACTIVE INGREDIENT OF  
BAYER® LOW DOSE ASPIRIN\*



Actual Size

**60 coated tablets**

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- each tablet contains: sodium 0.36 mg
- save carton for full directions and warnings
- store at room temperature

**Inactive ingredients**

acryoxol, colloidal silicon dioxide, croscarmellose sodium, D&C yellow #10, lactose, microcrystalline cellulose, polyphosphate 80, sodium hydroxide, talc, titanium dioxide, triethyl citrate, yellow oxide of iron

**Questions or comments? 1-888-333-9792**

\*This product is not manufactured or distributed by Bayer Corporation Consumer Care division, owner of the registered trademark Bayer® Low Dose Aspirin Regimen

LOT:  
EXP:

6 39277 34336 9



TAMPEN BUI DIEN: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

DISTRIBUTED BY  
GREENBRIER INTERNATIONAL, INC.  
500 VOLVO PARKWAY, CHESAPEAKE, VA 23320

ORIG 01/16



**ASSURED™**

COMPARE TO ACTIVE INGREDIENT OF  
BAYER® LOW DOSE ASPIRIN\*

**LOW DOSE**

**Aspirin**

• Aspirin (NSAID) 81 mg  
Pain Reliever



Actual Size

**40 coated tablets**

## ASSURED ASPIRIN

aspirin tablet, delayed release

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68210-0015
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

### Inactive Ingredients

Ingredient Name	Strength
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
LACTOSE, UNSPECIFIED FORM (UNII: J2B2A4N98G)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
WATER (UNII: 059QF0KO0R)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

### Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	8mm

<b>Flavor</b>		<b>Imprint Code</b>	1
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68210-0015-4	1 in 1 CARTON	04/06/2020	01/12/2022
1		40 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	07/01/2016	

**Labeler** - SPIRIT PHARMACEUTICALS LLC (179621011)

Revised: 11/2022

SPIRIT PHARMACEUTICALS LLC