

**MIGRAINE FORMULA- acetaminophen, aspirin, caffeine tablet, film coated**  
**Meijer Distribution Inc**

-----  
**Meijer Distribution, Inc. Migraine Formula Drug Facts**

**Active ingredients (in each caplet)**

Acetaminophen 250 mg

Aspirin 250 mg (NSAID\*)

Caffeine 65 mg

\*nonsteroidal anti-inflammatory drug

**Purposes**

Pain reliever

Pain reliever aid

**Use**

- treats migraine

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Medication overuse headache warning:** Headaches may worsen if this product is used for 10 or more days per month.

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

### **Ask a doctor before use if**

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever

- reducers
- you have vomiting with your migraine headache

### **Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for:
- diabetes
- gout
- arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- your migraine is not relieved or worsens after first dose
- new or unexpected symptoms occur
- ringing in the ears or loss of hearing occurs

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- do not use more than directed
- adults: take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

### **Other information**

- store at 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

### **Inactive ingredients**

carnauba wax, crospovidone, FD&C blue #1 aluminum lake, hypromellose, microcrystalline cellulose, povidone, pregelatinized starch, propylene glycol, saccharin sodium, stearic acid, titanium dioxide

### **Questions or comments?**

**1-800-719-9260**

### **Principal Display Panel**

meijer®

Compare to Excedrin® Migraine active ingredients

migraine formula

Acetaminophen | Aspirin (NSAID) | Caffeine Tablets

Pain Reliever | Pain Reliever Aid

100 Coated Caplets\*\*

\*\*capsule shaped tablets

actual size



**Drug Facts (continued)**

- Stop use and ask a doctor if:
  - an allergic reaction occurs. Seek medical help right away.
  - you experience any of the following signs or stomach bleeding:
    - feel faint, ■ vomit blood, ■ have bloody or black stools
    - have stomach pain that does not get better
  - your migraine is not relieved or worsens after 1st dose
  - new or unexpected symptoms occur
  - ringing in the ears or loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless directed by a doctor to do so for a reason (may cause problems in the unborn child or complications during delivery).  
 Keep out of reach of children.  
 In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not use more than directed
- adults: take 2 caplets with a glass of water
- if symptoms persist or worsen, ask your doctor
- do not take more than 2 caplets in 24 hours, unless directed by a doctor
- under 18 years of age: ask a doctor

**Other information**

- store at 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

**Inactive ingredients**

camalua wax, croscopolone, FD&C blue #1 aluminum lake, hypromellose, microcrystalline cellulose, polydine, pregelatinized starch, polyethylene glycol, saccharin sodium, stearic acid, titanium dioxide

Questions or comments? 1-800-719-9260



NDC 41250-374-78  
 Compare to  
 Excedrin® Migraine  
 active ingredients\*\*

# migraine formula

Acetaminophen | Aspirin (NSAID) | Caffeine Tablets



Pain Reliever | Pain Reliever Aid

100 Coated Caplets\*\*

\*\*capsule shaped tablets



actual size





## Drug Facts

### Active ingredients (in each caplet) Purposes

Acetaminophen 250 mg.....	Pain reliever
Aspirin 250 mg (NSAID)*.....	Pain reliever
Caffeine 65 mg.....	Pain reliever aid

\*nonsteroidal anti-inflammatory drug

**Use** ■ treats migraine

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening ■ blisters ■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 2 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

## Drug Facts (continued)

**Medication overuse headache warning:** Headaches may worsen if this product is used for 10 or more days per month.

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

### Do not use


- if you have ever had an allergic reaction to acetaminophen, aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

### Ask a doctor before use if

- you have never had migraines diagnosed by a health professional
- you have a headache that is different from your usual migraines
- you have the worst headache of your life
- you have fever and stiff neck
- you have headaches beginning after or caused by head injury, exertion, coughing or bending
- you experienced your first headache after the age of 50
- you have daily headaches
- you have a migraine so severe as to require bed rest
- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have vomiting with your migraine headache

### Ask a doctor or pharmacist before use if you are

- taking a prescription drug for:
  - diabetes ■ gout ■ arthritis
- under a doctor's care for any serious condition
- taking any other drug
- taking any other product that contains aspirin, acetaminophen, or any other pain reliever/fever reducer



## MIGRAINE FORMULA

acetaminophen, aspirin, caffeine tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-374
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
<b>CAFFEINE</b> (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>CROSPVIDONE, UNSPECIFIED</b> (UNII: 2S7830E561)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

### Product Characteristics

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	L374
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-374-71	1 in 1 CARTON	04/30/2002	
1		50 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41250-374-78	1 in 1 CARTON	04/30/2002	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:41250-374-85	250 in 1 BOTTLE; Type 0: Not a Combination Product	04/30/2002	06/28/2014
4	NDC:41250-374-82	1 in 1 CARTON	04/30/2002	
4		200 in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075794	04/30/2002	

**Labeler** - Meijer Distribution Inc (006959555)

Revised: 5/2023

Meijer Distribution Inc