

ANTI-DIARRHEAL- loperamide hcl tablet
H E B

HEB 44-375

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.

Heart alert: Taking more than directed can cause serious heart problems or death.

Do not use

if you have bloody or black stool.

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product,

drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store between 20°-25°C (68°-77°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

Compare to Imodium® A-D active ingredient*

NDC 37808-375-08

H-E-B®

Anti-Diarrheal

Loperamide Hydrochloride Tablets, 2 mg
Anti-Diarrheal

Diarrhea Relief

Controls Symptoms of Diarrhea,
Including Travelers' Diarrhea

24 CAPLETS

Each caplet contains 2 mg loperamide hydrochloride

actual
size

1. To open, tear at perforations and remove a section.
2. Use ✂ to cut through plastic for caplet.

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark Imodium® A-D.

50844 REV0619B37508

MADE WITH PRIDE AND CARE FOR H-E-B®, SAN ANTONIO, TX 78204

H-E-B®

100%
GUARANTEE
promise

If you aren't completely pleased with this product, we'll be happy to replace it or refund your money. You have our word on it.

Compare to **Imodium® A-D** active ingredient*

NDC 37808-375-08



Anti-Diarrheal

Loperamide Hydrochloride Tablets, 2 mg
Anti-Diarrheal

Drug Facts
Active ingredient (in each caplet) Loperamide HCl 2 mg
Purpose Anti-diarrheal
Use controls symptoms of diarrhea, including Traveler's Diarrhea

Drug Facts (continued)
Warnings
 Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.
 Heart alert: Taking more than directed can cause serious heart problems or death.
 Do not use if you have bloody or black stool.
 Ask a doctor before use if you have a fever, a history of abnormal heart rhythm, mucus in the stool, or a history of liver disease.
 Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.
 When using this product, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.
 Stop use and ask a doctor if symptoms get worse, diarrhea lasts for more than 2 days, you get abdominal swelling or bloating. These may be signs of a serious condition.
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
 Drink plenty of clear fluids to help prevent dehydration caused by diarrhea.
 Find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information
 TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BUSTER IS TORN OR BROKEN

Drug Facts (continued)
 store between 20°-25°C (68°-77°F) ■ see end flap for expiration date and lot number
Inactive ingredients D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, D&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide
Questions or comments? Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

50844 REV0619837508 43202-2404



MADE WITH PRIDE AND CARE FOR
 H-E-B®, SAN ANTONIO, TX 78204

100% GUARANTEE
 If you aren't completely pleased with this product, we'll be happy to replace it or refund your money. You have our word on it.
Pharmacia

B-0712-375-08-RR
 REV0619837508

*This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark Imodium® A-D.



1. To open, tear at perforations and remove a section.



2. Use scissors to cut through plastic for caplet.

no print / no varnish area
 lot no. & Exp. date

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BUSTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING



HEB 44-375

ANTI-DIARRHEAL

loperamide hcl tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37808-375
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	green	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	44;375
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-375-08	4 in 1 CARTON	05/03/2005	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination		

1		Product		
2	NDC:37808-375-02	2 in 1 CARTON	05/03/2005	02/18/2022
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076497	05/03/2005	

Labeler - H E B (007924756)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(37808-375) , pack(37808-375)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(37808-375)

Revised: 7/2024

H E B