

**ACETAMINOPHEN- acetaminophen tablet, film coated, extended release
PHARMACY VALUE ALLIANCE, LLC**

Arthritis Pain

Acetaminophen

Extended-release Tablets USP, 650 mg

Pain Reliever / Fever Reducer

For Temporary Relief of Minor Arthritis Pain

Contains No Aspirin

Active ingredient

(in each caplet)

Acetaminophen USP, 650 mg

Purpose

Pain reliever/fever reducer

Uses

(For Muscle Aches & Pain)

- temporarily relieves minor aches and pains due to:
 - muscular aches
 - backache
- minor pain of arthritis
- toothache
- premenstrual and menstrual cramps
- headache
- the common cold
- temporarily reduces fever

Uses

(For Arthritis Pain)

- temporarily relieves minor aches and pains due to:
 - minor pain of arthritis
 - muscular aches
 - backache
- premenstrual and menstrual cramps
- the common cold
- headache
- toothache
- temporarily reduces fever

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding

ask a health professional before use.

Keep out of reach of children

Keep out of reach of children

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions**(For Muscle Aches & Pain)**

- do not take more than directed (see overdose warning)
- adults and children 12 years of age and over
- take 2 caplets every 8 hours with water
 - swallow whole; do not crush, chew, split or dissolve
 - do not take more than 6 caplets in 24 hours
 - do not use for more than 10 days unless directed by a doctor
- children under 12 years of age
- do not use

Directions**(For Arthritis Pain)**

- do not take more than directed (see overdose warning)
- adults
- take 2 caplets every 8 hours with water
 - swallow whole; do not crush, chew, split or dissolve
 - do not take more than 6 caplets in 24 hours
 - do not use for more than 10 days unless directed by a doctor
- under 18 years of age
- ask a doctor

Other information

- store between 20-25°C (68-77°F)
- do not use if foil inner seal is broken or missing

Inactive ingredients

hydroxy ethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate, stearic acid.

Questions or comments?

Contact **1-877-770-3183** Mon-Fri 8:00 AM EST to 5:00 PM PST.

For Muscle Aches & Pain

end user :GRANDLASS
 Item :5816 3275 120720VWVW SCFF GRU
 code # :
 date :14/7/16 N 14/7/16 X 31/35/16
 ref # :PPI18130A view :FR
 material :.016 883 date :01

R-00
 529000028
 300000485

Drug Facts (continued)
Ingredients in this product
 Ask a doctor before use if you have liver disease
 Ask a doctor or pharmacist before use if you are taking the listed therapy drug warfarin
 Stop use and ask a doctor if
 in pain gets worse or lasts more than 10 days
 in fever gets worse or lasts more than 3 days
 in new symptoms occur
 in redness or swelling is present
 These could be signs of a serious condition.
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children.
 Overdose warnings: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
 Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions
 Do not take more than directed (see overdose warning)
 Adults and children 12 years of age and over
 in take 2 capsules every 8 hours with water
 in swallow whole; do not crush, chew, split or disperse
 in do not take more than 6 capsules in 24 hours
 in do not use for more than 10 days unless directed by a doctor
 Children under 12 years
 in do not use
Other Information
 in store between 20-25°C (68-77°F)
 in do not use if foil liner seal is broken or missing
Active Ingredients
 hydroxy ethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, p-

Drug Facts (continued)
 povidone, pregelatinized starch, sodium starch glycolate, stearic acid.
Questions or comments?
 contact 1-877-776-5883 Mon-Fri 8:00 AM EST to 5:00 PM PST.
 *This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® 8 HR Muscle Aches & Pain.
 Distributed by:
 Pharmacy Value Alliance, LLC
 407 East Lancaster Avenue,
 Wayne, PA 19087
 If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.

Muscle Aches & Pain
Acetaminophen
 8 HR
 Extended-release Tablets, USP, 650 mg
 Pain Reducer/Fever Reducer
 Not to be taken later of than shown below & this Contains No Aspirin

Muscle Aches & Pain
Acetaminophen
 8 HR
 Extended-release Tablets USP, 650 mg
 Pain Reducer/Fever Reducer
 Not to be taken later of than shown below & this Contains No Aspirin
 50 Capsules
 650 mg each
 (10-capsule strength)
 (N-1499 Tablets)
 PHARMACEUTICAL INFORMATION
 NDC 58163-3275-10
 NDC 58163-3275-10

Important: Read all product information before using. Keep the carton for important information.
 Do not use if foil liner seal is broken or missing.
Drug Facts
Active Ingredients (in each capsule) Purpose
 Acetaminophen USP, 650 mg Pain reducer/fever reducer
Dose
 in temporarily relieve minor aches and pains due to:
 in muscle aches
 in headache
 in other pain of arthritis
 in toothache
 in premenstrual and menstrual cramps
 in headache
 in the common cold
 in temporarily reduce fever
Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take in more than 6 capsules in 24 hours, which is the maximum daily amount
 in with other drugs containing acetaminophen
 in 3 or more alcoholic drinks every day while using this product
 Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
 in skin redness/itching
 in hives
 in rash
 If a skin reaction occurs, stop use and seek medical help right away.
 Do not use
 in with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 in if you are allergic to acetaminophen or any of the inactive

300000485
 529000028
 R-00

MADE IN INDIA
 8 40986 03612 5
 Lot
 Exp.

For Arthritis Pain

Item : 510 LOTS IMPROVED SUPP 051
 code # :
 date : 11/17/08 X 11/17/08 X 3/31/04
 lot # : 090111378
 material : 016 000
 view : PHT
 date : 01/1

R-00
 529000029
 300000487

Drug Facts (continued)
 Ingredients in this product:
 Ask a doctor before use if you have liver disease
 Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin
 Stop use and ask a doctor if:
 pain gets worse or lasts more than 10 days
 fever gets worse or lasts more than 3 days
 all new symptoms occur
 rash or swelling is present
 These could be signs of a serious condition.
 If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.
 Caution warnings: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)
 Caution: Medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions
 Do not take more than directed (see overdose warning)
 adults: take 2 capsules every 8 hours with water
 in swallow whole; do not crush, chew, split or disperse
 do not take more than 6 capsules in 24 hours
 do not use for more than 10 days unless directed by a doctor
 under 18 years of age: ask a doctor
Other information
 Store between 20-25°C (68-77°F)
 do not use if foil liner seal is broken or missing
Inactive ingredients
 hydroxyethyl cellulose, hydropropylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, p-

Drug Facts (continued)
 povidone, pregelatinized starch, sodium starch glycolate, stearic acid.
Questions or concerns?
 contact 1-877-776-2188 Mon-Fri 8:00 AM EST to 5:00 PM PST.
 **This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® 6 HR Arthritis Pain.
 Distributed By:
 Pharmacy Value Alliance, LLC
 407 East Lancaster Avenue,
 Myrtle, PA 19967
 If for any reason you are not satisfied with this product, please return it to the store where purchased for a full refund.


Arthritis Pain
Acetaminophen
 Extended-release Tablets USP, 650 mg
 Pain Reducer/Fever Reducer
 For the Temporary Relief of Minor Arthritis Pain
 Contains No Aspirin
 9 Hr
 650 mg each
 50 Capsules

Arthritis Pain
Acetaminophen
 Extended-release Tablets USP, 650 mg
 Pain Reducer/Fever Reducer
 For the Temporary Relief of Minor Arthritis Pain
 Contains No Aspirin
 8 Hr
 650 mg each
 50 Capsules
 CHERRY TO THE CORE ASSORTED OR TILLY'S™ 1 HR ARTHRIS PAIN

Important: Read all product information before using. Keep the carton for important information.
 Do not use if foil liner seal is broken or missing.
Drug Facts
Active ingredient (in each capsule) Purpose
 Acetaminophen USP 650 mg Pain reducer/fever reducer
Uses
 to temporarily relieve minor aches and pains due to:
 minor pain of arthritis
 muscular aches
 headache
 sinus and nasal congestion
 the common cold
 headache
 to temporarily reduce fever
Warnings
 Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 capsules in 24 hours, which is the maximum daily amount.
 Do not use with other drugs containing acetaminophen.
 3 or more alcoholic drinks every day while using this product.
 Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:
 skin redness
 skin itching
 If a skin reaction occurs, stop use and seek medical help right away.
 Do not use with any other drug containing acetaminophen (paracetamol or nonprescription) if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
 If you are allergic to acetaminophen or any of the inactive

300000487
 529000029
 R-00


MADE IN INDIA
 40986 03613 2
 Lot:
 Exp:



Don't modifications to side 2 incoming artwork:
 1) Updated adhesive deadener pattern



REINSERTION PRINTS ON HP FOR FRONT



acetaminophen tablet, film coated, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68016-183
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg

Inactive Ingredients

Ingredient Name	Strength
HYDROXYETHYL CELLULOSE (4000 MPAS AT 1%) (UNII: ZYD53NBL45)	
HYPROMELLOSE 2910 (6 MPAS) (UNII: 0WZ8WG20P6)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	white (White to off white colored)	Score	no score
Shape	OVAL (Capsule shaped, biconvex intact film coated tablets)	Size	19mm
Flavor		Imprint Code	G;650
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-183-05	50 in 1 BOTTLE; Type 0: Not a Combination Product	11/15/2019	
2	NDC:68016-183-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	11/15/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA211544	11/15/2019	

Labeler - PHARMACY VALUE ALLIANCE, LLC (101668460)