ACETAMINOPHEN- acetaminophen tablet Kroger Company

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Kroger Co. Acetaminophen Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- the common cold
- headache
- backache
- · minor pain of arthritis
- toothache
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

liver disease

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults and children 12
years and over

- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

children under 12 years ask a doctor

Other information

store at 20-25°C (68-77°F)

Inactive ingredients

carnauba wax, corn starch*, croscarmellose sodium*, hypromellose, polyethylene glycol, povidone, pregelatinized starch, sodium starch glycolate*, stearic acid

*may contain one or more of these ingredients

Questions or comments?

1-800-632-6900

Principal Display Panel

COMPARE TO the active ingredient of EXTRA STRENGTH TYLENOL® CAPLETS See back panel

OUR PHARMACIST RECOMMENDED

for adults

Extra Strength

actual size

Acetaminophen 500 mg

Pain Reliever/Fever Reducer

100 CAPLETS 500 mg EACH





Drug Facts

Active ingredient (in each caplet)

Acetaminophen 500 mg......Pain reliever/fever reducer

Purpose

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache minor pain of arthritis ■ backache

 - muscular aches ■ toothache
 - premenstrual and menstrual cramps
- temporarily reduces fever

Wamings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
 blisters
 rash If a skin reaction occurs, stop use and seek medical help right away.
- Do not use with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
 redness or swelling is present These could be signs of a serious condition.

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults and children 12 years and over	■ take 2 caplets every 6 hours while symptoms last ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor
children under 12 years	ask a doctor

Other information ■ store at 20-25°C (68-77°F)

Inactive ingredients camauta wax, com starch*, croscarmellose sodium*, hypromellose, polyethylene glycol, povidone, pregelatini zed starch, sodium starch glycolate*, stearic acid *may contain one or more of these ingredients

Questions or comments? 1-800-632-6900

"Tylenol" is a registered trademark of Johnson & Johnson Corporation, New Brunswick, NJ 08933. Johnson & Johnson Corporation is not affiliated with The Kroger Co. or this product.

DISTRIBUTED BY THE KROGER CO. CINCINNATI, OHIO 45202

QUALITY GUARANTEE 800-632-6900 www.kroger.com

GLUTEN FREE







ACETAMINOPHEN

acetaminophen tablet

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:30142-561	
Route of Administration	ORAL			

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	500 mg		

Inactive Ingredients			
Ingredient Name	Strength		
CARNAUBA WAX (UNII: R12CBM0EIZ)			
STARCH, CORN (UNII: O8232NY3SJ)			
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)			
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)			
STEARIC ACID (UNII: 4ELV7Z65AP)			
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)			

Product Characteristics				
Color	WHITE	Score	no score	
Shape	OVAL (caplet)	Size	16mm	
Flavor		Imprint Code	L484	
Contains				

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:30142-561-71	1 in 1 CARTON	07/15/1987	04/30/2020		
1		50 in 1 BOTTLE; Type 0: Not a Combination Product				

2	NDC:30142-561-78	1 in 1 CARTON	07/15/1987	
2		100 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:30142-561- 85	1 in 1 CARTON	07/15/1987	07/15/1987
3		250 in 1 BOTTLE; Type 0: Not a Combination Product		
4	NDC:30142-561- 90	500 in 1 BOTTLE; Type 0: Not a Combination Product	07/15/1987	03/01/2023
5	NDC:30142-561- 82	2 in 1 CARTON	07/15/1987	07/15/1987
5		100 in 1 BOTTLE; Type 0: Not a Combination Product		
6	NDC:30142-561- 76	1 in 1 CARTON	07/15/1987	11/30/2020
6		120 in 1 BOTTLE; Type 0: Not a Combination Product		
7	NDC:30142-561- 62	1 in 1 CARTON	05/10/2018	11/30/2022
7		24 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part343	07/15/1987		

Labeler - Kroger Company (006999528)

Revised: 11/2022 Kroger Company