#### QUALITY CHOICE MAXIMUM STRENGTH ORAL PAIN RELIEVING- benzocaine gel Chain Drug Marketing Association Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

#### QC Maximum Strength Oral Analgesic 0.42oz 95362 (2019) - ZDP

#### **Active ingredient Purpose**

Benzocaine 20%......Oral pain reliever

#### Uses

- for the temporarily relieves pain due to:
- toothache
- canker sores
- gum pain
- denture pain
- cheek bites

#### Warnings

Methemoglobinemia warning: Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

Allergy alert: do not use this product if your child has a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

#### Do not use

- more than directed
- for more than 7 days unless directed by a doctor
- for teething
- in children under 2 years of age

Stop use and ask a doctor if

- sore mouth symptoms do not improve in 7 days
- irritation, pain, or redness persists or worsens
- swelling, rash, or fever develops

**If pregnant or breast-feeding**, ask a health professional before use.

Keep out of reach of children. In cases of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

Adults and children 2 years of age and older.

product to the cavity and around gum surrounding the teeth. Use up to 4 times daily or as directed by a dentist or

doctor.

Children under 12 years of age.

Children under 2 years of age.

Should be supervised in the use of this product.

Do not use.

#### Other information

• store at room temperature 20° to 25°C (68° to 77°F)

- this preparation is intended for use in cases of toothache, only as a temporary expedient until a dentist or doctor can be consulted
- do not use continuously

#### Inactive ingredients

glycerin, peppermint oil, polyethylene glycol 400, polyethylene glycol 4000, saccharin sodium, sorbitol

Distributed By C.D.M.A., Inc.

43157 W 9 Mile Rd

Novi, MI 48375

www.qualitychoice.com

Made in China



#### QUALITY CHOICE MAXIMUM STRENGTH ORAL PAIN RELIEVING

benzocaine gel

Prod	luct	Informa	ation
1 1 0 0	uuu		alivii

Product Type HUMAN OTC DRUG Item Code (Source) NDC:63868-477

Route of Administration TOPICAL

#### **Active Ingredient/Active Moiety**

reave ingredient workly			
	Ingredient Name	Basis of Strength	Strength
	BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5)	BENZOCAINE	200 mg in 1 g

# Inactive Ingredients Ingredient Name Strength POLYETHYLENE GLYCOL 4000 (UNII: 4R4HFi6D95) POLYETHYLENE GLYCOL 4000 (UNII: B697894SGQ) SACCHARIN SODIUM (UNII: SB8ZUX40TY) GLYCERIN (UNII: PDC6A3C0OX)

SORBITOL (UNII: 506T60A25R)

SORBITOL (ONII. 300 100 A23K)

PEPPERMINT OIL (UNII: AV092KU4JH)

# Packaging#Item CodePackage DescriptionMarketing Start DateMarketing End Date1NDC:63868-477-111 in 1 CARTON08/09/2019

1 1.9 g in 1 TUBE; Type 0: Not a Combination Product

## **Marketing Information**

with hearing imorr	arricang mucron						
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date				
OTC monograph not final	part356	08/07/2019					

### Labeler - Chain Drug Marketing Association Inc (011920774)

Revised: 8/2019 Chain Drug Marketing Association Inc