

FS IODINE TINCTURE- iodine liquid
Genex Cooperative, Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

FS Iodine Tincture

FS Iodine Tincture

Iodine Tincture 7%™ is a topical antiseptic that should be used full strength for superficial cuts, wounds, abrasions, insect bites and bruises on the skin of animals.

DANGER

Hazard statement(s):

Toxic if swallowed. May cause eye irritation.

Precautionary statement(s):

Keep away from heat, sparks, open flames. No smoking. Do not eat, drink or smoke while using this product. Wear protective clothing, and eye and face protection.

Hazards not otherwise classified: Flammable

Not for use on burns, deep cuts, or body cavities.

First Aid:

External: : If irritation occurs, flush with plenty of cool, running water for at least 15 minutes while removing contaminated clothing and shoes. Wash clothing before reuse.

Eyes: Immediately flush with plenty of cool, running water. Remove contact lenses; then continue flushing for at least 15 minutes, holding eyelids apart.

Internal: Call a physician immediately. If swallowed, do not induce vomiting. Rinse mouth, then immediately drink one or two large glasses of water or milk. Never give anything by mouth to an unconscious person.

Inhalation of vapors: Should not occur with proper use of this product.

Most important symptoms/effects, acute and delayed: None known.

Immediate medical attention and special treatment needed, if any: If swallowed, give starch paste, milk, bread, egg white, or activated charcoal. A 5% solution of sodium thiosulfate (photographic ("hypo")) may be administered orally at a rate of 10 ml per kilogram of body weight.

DIRECTIONS:

Use full Strength for superficial cuts, wounds, abrasions, insect bites and bruises on the skin of animals. Apply iodine with a swab.

- If necessary, clip hair around the area being treated and clean with soap and water.
- Apply Iodine Tincture 7%™ only once daily. dilute product 3 to 1 if repeating application.
- Do not apply under bandage.
- Irritation may occur if used on tender skin areas. If redness, irritation, or swelling persists or increases, discontinue use and consult a veterinarian.

STORAGE:

- Store at 2-30 degrees C (36-86 degrees F).

- Keep container away from heat and out of sunlight

**FOR CHEMICAL EMERGENCIES
CALL CHEMTREC AT 800-424-9300**

FS Iodine Tincture



FS Iodine Tincture™
TOPICAL ANTISEPTIC

Active Ingredients:
Titratable Iodine.....7.00%
Isopropyl Alcohol.....65.00%

For Farm & Industrial Use Only

KEEP OUT OF REACH OF CHILDREN

DIRECTIONS:

Use full Strength for superficial cuts, wounds, abrasions, insect bites and bruises on the skin of animals. Apply iodine with a swab.

- If necessary, clip hair around the area being treated and clean with soap and water.
- Apply Iodine Tincture 7%™ only once daily, dilute product 3 to 1 if repeating application.
- Do not apply under bandage.
- Irritation may occur if used on tender skin areas. If redness, irritation, or swelling persists or increases, discontinue use and consult a veterinarian.

STORAGE:

- Store at 2-30 degrees C (36-86 degrees F).
- Keep container away from heat and out of sunlight



Manufactured by:
ProActive Solutions® USA, LLC • 1-800-279-7761
301 Bridge St. Green Bay, WI

FS Iodine Tincture™ Topical Antiseptic

Iodine Tincture 7%™ is a topical antiseptic that should be used full strength for superficial cuts, wounds, abrasions, insect bites and bruises on the skin of animals.

DANGER

Hazard statement(s):

Toxic if swallowed. May cause eye irritation.

Precautionary statement(s):

Keep away from heat, sparks, open flames. No smoking. Do not eat, drink or smoke while using this product. Wear protective clothing, and eye and face protection.

Hazards not otherwise classified: Flammable
Not for use on burns, deep cuts, or body cavities.

First Aid:

External: : If irritation occurs, flush with plenty of cool, running water for at least 15 minutes while removing contaminated clothing and shoes. Wash clothing before reuse.

Eyes: Immediately flush with plenty of cool, running water. Remove contact lenses; then continue flushing for at least 15 minutes, holding eyelids apart.

Internal: Call a physician immediately. If swallowed, do not induce vomiting. Rinse mouth, then immediately drink one or two large glasses of water or milk. Never give anything by mouth to an unconscious person.

Inhalation of vapors: Should not occur with proper use of this product.

Most important symptoms/effects, acute and delayed: None known.

Immediate medical attention and special treatment needed, if any: If swallowed, give starch paste, milk, bread, egg white, or activated charcoal. A 5% solution of sodium thiosulfate (photographic ("hypo")) may be administered orally at a rate of 10 ml per kilogram of body weight.

**FOR CHEMICAL EMERGENCIES
CALL CHEMTREC AT 800-424-9300**

UN 1293
TINCTURES, MEDICINAL, (IODINE TINCTURE 7% 3, II

Manufactured For:
Farm Systems
58 Interstate Dr NW
Melrose, MN 56362
+1-800-636-5681



FS IODINE TINCTURE

iodine liquid

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:86193-4680
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	7 kg in 100 kg

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:86193-4680-1	13.94 kg in 1 CASE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		01/23/2024	

Labeler - Genex Cooperative, Inc. (003437378)

Registrant - ProActive Solutions USA, LLC (029368225)

Establishment

Name	Address	ID/FEI	Business Operations
ProActive Solutions USA, LLC		029368225	api manufacture, manufacture, pack, label

Establishment

Name	Address	ID/FEI	Business Operations
Genex Cooperative, Inc.		003437378	repack

Revised: 1/2024

Genex Cooperative, Inc.